



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:03 AM UTC

PDB ID : 4PGH / pdb_00004pgh
Title : Caffeic acid O-methyltransferase from Sorghum bicolor
Authors : Green, A.R.; Lewis, K.M.; Kang, C.
Deposited on : 2014-05-02
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	93% 6%
1	B	358	2% 93% 7%
1	C	358	90% 9%
1	D	358	10% 87% 13%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10719 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caffeic acid O-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2616	1675	438	483	20			
1	B	357	Total	C	N	O	S	0	0	0
			2533	1608	436	469	20			
1	C	353	Total	C	N	O	S	0	0	0
			2612	1670	437	485	20			
1	D	358	Total	C	N	O	S	0	0	0
			2636	1688	443	485	20			

- Molecule 2 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	D	1	Total	C	N	O	S	0	0
			27	15	6	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total	O	0	0
			62	62		
3	B	43	Total	O	0	0
			43	43		
3	C	78	Total	O	0	0
			78	78		
3	D	31	Total	O	0	0
			31	31		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Caffeic acid O-methyltransferase

Chain A: 




- Molecule 1: Caffeic acid O-methyltransferase

Chain B: 




- Molecule 1: Caffeic acid O-methyltransferase


Chain C: 



- Molecule 1: Caffeic acid O-methyltransferase

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.78Å 63.90Å 152.47Å 90.00° 93.02° 90.00°	Depositor
Resolution (Å)	48.94 – 2.80 48.94 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.94-2.80) 97.2 (48.94-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.30 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.204 , 0.252 0.208 , 0.253	Depositor DCC
R_{free} test set	1999 reflections (5.43%)	wwPDB-VP
Wilson B-factor (Å ²)	58.7	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10719	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9957e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2680	0.72	0/3655
1	B	0.27	0/2595	0.71	0/3551
1	C	0.27	0/2676	0.71	2/3652 (0.1%)
1	D	0.27	0/2700	0.72	0/3684
All	All	0.27	0/10651	0.71	2/14542 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	306	VAL	CA-C-N	5.23	125.54	119.47
1	C	306	VAL	C-N-CA	5.23	125.54	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2616	0	2436	13	0
1	B	2533	0	2220	15	0
1	C	2612	0	2426	20	0
1	D	2636	0	2496	26	0
2	A	27	0	22	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	27	0	22	1	0
2	C	27	0	22	1	0
2	D	27	0	22	0	0
3	A	62	0	0	1	0
3	B	43	0	0	0	0
3	C	78	0	0	4	0
3	D	31	0	0	0	0
All	All	10719	0	9666	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:250:MET:O	1:D:280:ASN:ND2	2.23	0.71
1:C:229:ASP:O	3:C:553:HOH:O	2.11	0.69
1:C:248:GLY:N	3:C:553:HOH:O	2.24	0.67
1:D:127:MET:HE1	1:D:177:MET:HE2	1.79	0.64
1:D:306:VAL:HG22	1:D:308:LYS:H	1.63	0.64
1:D:76:ASP:OD2	1:D:101:ARG:NH2	2.30	0.61
1:B:263:LYS:NZ	2:B:401:SAM:OXT	2.34	0.60
1:C:345:GLY:HA3	1:C:361:ILE:HB	1.83	0.59
1:A:327:ARG:NH2	3:A:511:HOH:O	2.38	0.57
1:D:268:ASP:OD1	1:D:323:ASN:ND2	2.39	0.56
1:D:82:LEU:HB3	1:D:87:VAL:HB	1.89	0.55
1:A:262:MET:HG3	1:A:265:ILE:HB	1.89	0.54
1:B:203:VAL:HG22	1:B:226:VAL:HB	1.91	0.53
1:C:10:ALA:N	3:C:513:HOH:O	2.42	0.53
1:D:285:LEU:HD11	1:D:292:VAL:HG23	1.90	0.52
1:C:248:GLY:O	3:C:553:HOH:O	2.18	0.50
1:D:150:PRO:HG2	1:D:321:ALA:HA	1.92	0.50
1:D:240:PHE:HB2	1:D:243:VAL:HB	1.93	0.50
1:A:345:GLY:HA3	1:A:361:ILE:HB	1.94	0.50
1:D:249:ASP:OD1	1:D:249:ASP:N	2.44	0.50
1:B:82:LEU:HB3	1:B:87:VAL:HB	1.93	0.50
1:C:74:MET:HG2	1:D:320:LEU:HD11	1.93	0.50
1:B:251:PHE:HE1	1:B:273:HIS:HB3	1.76	0.49
1:B:81:LEU:O	1:B:84:SER:OG	2.30	0.49
1:C:145:LEU:HA	1:D:66:PRO:HB3	1.94	0.48
1:D:269:TRP:HB3	1:D:273:HIS:HB2	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:ASP:HA	1:B:261:LEU:HB3	1.93	0.48
1:A:82:LEU:HB3	1:A:87:VAL:HB	1.94	0.48
1:D:345:GLY:N	1:D:361:ILE:O	2.47	0.48
1:D:175:GLU:O	1:D:179:ASN:ND2	2.40	0.47
1:C:262:MET:HG3	1:C:265:ILE:HB	1.97	0.47
1:C:82:LEU:HB3	1:C:87:VAL:HB	1.95	0.47
1:C:150:PRO:HG2	1:C:321:ALA:HA	1.97	0.46
1:C:323:ASN:HA	1:C:324:PRO:HD3	1.81	0.46
1:D:262:MET:HE3	1:D:262:MET:HB2	1.80	0.46
1:A:16:ALA:HB1	1:B:108:VAL:HA	1.98	0.46
1:D:315:VAL:O	1:D:319:MET:HG2	2.16	0.45
1:C:189:LEU:HD21	1:C:216:ILE:HG12	1.97	0.45
1:A:170:ASN:O	1:A:174:ASN:ND2	2.29	0.45
1:B:125:ALA:O	1:B:129:GLN:NE2	2.49	0.45
1:A:94:ASP:CG	1:B:332:ARG:HH22	2.25	0.45
1:A:200:SER:OG	1:A:258:ASP:OD1	2.33	0.45
1:D:264:TRP:HB3	1:D:267:HIS:HE1	1.82	0.45
1:B:348:ALA:HA	1:B:358:ILE:HA	1.98	0.45
1:A:68:ASN:ND2	1:B:147:GLY:HA2	2.32	0.44
1:C:85:TYR:O	1:D:19:TYR:OH	2.32	0.44
1:C:262:MET:HE1	1:C:281:CYS:SG	2.58	0.44
1:D:57:GLU:OE2	1:D:102:ARG:NH2	2.50	0.44
1:B:62:LEU:HG	1:B:63:PRO:HD2	2.00	0.43
1:D:133:LEU:HD22	1:D:322:HIS:CD2	2.53	0.43
1:C:170:ASN:OD1	1:C:174:ASN:ND2	2.51	0.43
1:D:292:VAL:HB	1:D:360:PHE:HB2	2.01	0.42
1:A:27:SER:O	1:A:31:MET:HG2	2.19	0.42
1:A:74:MET:HG2	1:B:320:LEU:HD11	2.00	0.42
1:B:262:MET:HB2	1:B:294:VAL:HG22	2.00	0.42
1:B:300:PRO:HG2	1:B:309:ALA:HA	2.01	0.42
1:A:263:LYS:NZ	2:A:401:SAM:O	2.48	0.41
1:C:263:LYS:NZ	2:C:401:SAM:OXT	2.43	0.41
1:C:292:VAL:HB	1:C:360:PHE:HB2	2.02	0.41
1:D:257:GLY:O	1:D:285:LEU:HD23	2.21	0.41
1:A:262:MET:HE1	1:A:281:CYS:SG	2.61	0.40
1:C:140:LEU:HB3	1:D:35:ASN:CG	2.47	0.40
1:D:160:PHE:HA	1:D:163:HIS:CE1	2.56	0.40
1:C:68:ASN:ND2	1:D:147:GLY:HA2	2.35	0.40
1:C:299:LEU:HD23	1:C:312:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/358 (98%)	343 (97%)	9 (3%)	0	100	100
1	B	355/358 (99%)	347 (98%)	8 (2%)	0	100	100
1	C	351/358 (98%)	338 (96%)	13 (4%)	0	100	100
1	D	356/358 (99%)	342 (96%)	13 (4%)	1 (0%)	36	66
All	All	1414/1432 (99%)	1370 (97%)	43 (3%)	1 (0%)	48	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	63	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/287 (85%)	242 (100%)	1 (0%)	84	94
1	B	213/287 (74%)	212 (100%)	1 (0%)	81	93
1	C	244/287 (85%)	240 (98%)	4 (2%)	55	83
1	D	255/287 (89%)	253 (99%)	2 (1%)	73	90
All	All	955/1148 (83%)	947 (99%)	8 (1%)	73	90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	234	ILE
1	B	62	LEU
1	C	81	LEU
1	C	119	VAL
1	C	303	THR
1	C	306	VAL
1	D	53	LEU
1	D	279	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	323	ASN
1	B	180	HIS
1	B	323	ASN
1	C	322	HIS
1	D	22	GLN
1	D	232	HIS
1	D	267	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SAM	B	401	-	27,29,29	1.53	4 (14%)	34,42,42	2.50	12 (35%)
2	SAM	D	401	-	27,29,29	1.54	4 (14%)	34,42,42	2.46	13 (38%)
2	SAM	A	401	-	27,29,29	1.53	4 (14%)	34,42,42	2.43	11 (32%)
2	SAM	C	401	-	27,29,29	1.52	4 (14%)	34,42,42	2.37	11 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAM	B	401	-	-	7/17/33/33	0/3/3/3
2	SAM	D	401	-	-	4/17/33/33	0/3/3/3
2	SAM	A	401	-	-	4/17/33/33	0/3/3/3
2	SAM	C	401	-	-	8/17/33/33	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	SAM	C6-N6	3.77	1.43	1.34
2	A	401	SAM	C6-N6	3.76	1.43	1.34
2	B	401	SAM	C6-N6	3.66	1.43	1.34
2	C	401	SAM	C6-N6	3.66	1.43	1.34
2	C	401	SAM	C8-N9	-2.83	1.32	1.37
2	A	401	SAM	C8-N9	-2.80	1.32	1.37
2	B	401	SAM	C2'-C1'	-2.75	1.44	1.53
2	D	401	SAM	C2'-C1'	-2.69	1.45	1.53
2	D	401	SAM	C8-N9	-2.62	1.33	1.37
2	B	401	SAM	C8-N9	-2.55	1.33	1.37
2	C	401	SAM	C2'-C1'	-2.53	1.45	1.53
2	A	401	SAM	C2'-C1'	-2.49	1.45	1.53
2	B	401	SAM	O4'-C1'	2.24	1.47	1.42
2	A	401	SAM	O4'-C1'	2.13	1.46	1.42
2	C	401	SAM	O4'-C1'	2.09	1.46	1.42
2	D	401	SAM	C1'-N9	-2.01	1.40	1.46

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	SAM	CG-SD-C5'	7.24	121.13	103.43
2	A	401	SAM	CG-SD-C5'	7.08	120.73	103.43
2	D	401	SAM	CG-SD-C5'	6.89	120.27	103.43
2	C	401	SAM	CG-SD-C5'	6.78	120.01	103.43
2	C	401	SAM	C5-C4-N3	-5.16	119.61	126.72
2	A	401	SAM	C5-C4-N3	-5.11	119.68	126.72
2	B	401	SAM	C5-C4-N3	-5.07	119.74	126.72
2	D	401	SAM	C5-C4-N3	-4.66	120.30	126.72
2	B	401	SAM	N3-C4-N9	4.57	134.93	127.17
2	D	401	SAM	N3-C2-N1	-4.37	121.97	128.58
2	C	401	SAM	N3-C4-N9	4.28	134.45	127.17
2	A	401	SAM	N3-C4-N9	4.20	134.31	127.17
2	C	401	SAM	N3-C2-N1	-4.11	122.36	128.58
2	A	401	SAM	N3-C2-N1	-4.10	122.38	128.58
2	D	401	SAM	C4-N9-C8	3.98	109.91	105.74
2	D	401	SAM	N3-C4-N9	3.96	133.91	127.17
2	B	401	SAM	N3-C2-N1	-3.92	122.65	128.58
2	B	401	SAM	C4-N9-C8	3.91	109.84	105.74
2	D	401	SAM	C2-N3-C4	3.70	120.88	111.83
2	C	401	SAM	C2-N3-C4	3.63	120.70	111.83
2	D	401	SAM	N9-C8-N7	-3.63	108.78	113.94
2	A	401	SAM	C2-N3-C4	3.63	120.70	111.83
2	A	401	SAM	C4-N9-C8	3.33	109.23	105.74
2	B	401	SAM	C2-N3-C4	3.32	119.93	111.83
2	C	401	SAM	C4-N9-C8	3.31	109.22	105.74
2	B	401	SAM	N9-C8-N7	-3.28	109.29	113.94
2	B	401	SAM	C5-N7-C8	3.21	108.49	103.45
2	A	401	SAM	N9-C8-N7	-3.14	109.48	113.94
2	A	401	SAM	C4-C5-N7	-3.10	107.04	110.58
2	B	401	SAM	C4-C5-N7	-3.09	107.05	110.58
2	C	401	SAM	C4-C5-N7	-3.06	107.09	110.58
2	A	401	SAM	C5-N7-C8	3.05	108.25	103.45
2	C	401	SAM	N9-C8-N7	-3.03	109.64	113.94
2	D	401	SAM	C5-N7-C8	3.01	108.18	103.45
2	C	401	SAM	C5-N7-C8	2.97	108.12	103.45
2	D	401	SAM	C4-C5-N7	-2.85	107.33	110.58
2	B	401	SAM	C3'-C2'-C1'	2.51	106.22	101.46
2	D	401	SAM	CE-SD-C5'	2.49	120.09	100.54
2	A	401	SAM	CE-SD-C5'	2.42	119.55	100.54
2	D	401	SAM	C4'-O4'-C1'	-2.41	104.14	109.47
2	D	401	SAM	C6-C5-N7	2.40	136.72	132.09
2	C	401	SAM	CE-SD-C5'	2.40	119.39	100.54
2	B	401	SAM	CE-SD-C5'	2.35	119.01	100.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	401	SAM	C4-N9-C1'	-2.30	121.25	126.63
2	B	401	SAM	C4-N9-C1'	-2.20	121.50	126.63
2	A	401	SAM	C6-C5-N7	2.11	136.15	132.09
2	C	401	SAM	C6-C5-N7	2.01	135.97	132.09

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	SAM	N-CA-CB-CG
2	A	401	SAM	C-CA-CB-CG
2	A	401	SAM	CB-CG-SD-CE
2	A	401	SAM	C4'-C5'-SD-CE
2	B	401	SAM	CA-CB-CG-SD
2	B	401	SAM	CB-CG-SD-CE
2	C	401	SAM	N-CA-CB-CG
2	C	401	SAM	C-CA-CB-CG
2	C	401	SAM	CB-CG-SD-C5'
2	C	401	SAM	C4'-C5'-SD-CE
2	C	401	SAM	O4'-C4'-C5'-SD
2	C	401	SAM	C3'-C4'-C5'-SD
2	D	401	SAM	N-CA-CB-CG
2	D	401	SAM	C-CA-CB-CG
2	D	401	SAM	CB-CG-SD-CE
2	D	401	SAM	C4'-C5'-SD-CE
2	B	401	SAM	C2'-C1'-N9-C8
2	B	401	SAM	C3'-C4'-C5'-SD
2	B	401	SAM	C2'-C1'-N9-C4
2	B	401	SAM	O4'-C1'-N9-C8
2	C	401	SAM	OXT-C-CA-CB
2	B	401	SAM	OXT-C-CA-N
2	C	401	SAM	O-C-CA-CB


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	SAM	1	0
2	A	401	SAM	1	0
2	C	401	SAM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/358 (98%)	-0.21	1 (0%) 90 86	32, 53, 90, 108	0
1	B	357/358 (99%)	0.06	8 (2%) 62 52	29, 69, 103, 130	0
1	C	353/358 (98%)	-0.25	1 (0%) 90 86	37, 56, 93, 154	0
1	D	358/358 (100%)	0.57	36 (10%) 12 9	33, 95, 149, 166	0
All	All	1422/1432 (99%)	0.04	46 (3%) 50 40	29, 63, 127, 166	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	166	ASP	4.8
1	D	169	PHE	4.6
1	D	211	ALA	3.7
1	D	221	SER	3.7
1	D	170	ASN	3.7
1	B	345	GLY	3.6
1	D	303	THR	3.6
1	B	304	ASP	3.4
1	D	338	ALA	3.4
1	D	164	GLY	3.3
1	D	167	PRO	3.2
1	D	351	ILE	3.2
1	D	228	PHE	3.2
1	D	290	GLY	3.0
1	D	241	PRO	2.9
1	B	198	SER	2.9
1	D	343	PHE	2.8
1	D	215	ALA	2.7
1	D	262	MET	2.7
1	D	204	ASP	2.7
1	D	163	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	322	HIS	2.6
1	D	357	ALA	2.6
1	D	324	PRO	2.6
1	B	196	ASP	2.6
1	D	205	VAL	2.6
1	B	361	ILE	2.5
1	D	199	VAL	2.5
1	D	64	VAL	2.4
1	B	303	THR	2.3
1	D	280	ASN	2.3
1	D	281	CYS	2.2
1	D	157	MET	2.2
1	D	65	ALA	2.2
1	B	302	THR	2.2
1	B	197	GLU	2.2
1	C	304	ASP	2.2
1	D	354	ASN	2.1
1	A	324	PRO	2.1
1	D	209	ILE	2.1
1	D	255	PRO	2.1
1	D	323	ASN	2.1
1	D	191	PHE	2.1
1	D	63	PRO	2.0
1	D	341	ALA	2.0
1	D	213	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SAM	D	401	27/27	0.59	0.14	124,145,169,179	0
2	SAM	B	401	27/27	0.75	0.12	93,104,108,110	0
2	SAM	A	401	27/27	0.90	0.09	44,55,77,81	0
2	SAM	C	401	27/27	0.92	0.08	48,58,80,90	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around SAM B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAM A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.