



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:15 PM UTC

PDB ID : 2PH3 / pdb_00002ph3
Title : Crystal structure of 3-oxoacyl-[acyl carrier protein] reductase TTHA0415 from *Thermus thermophilus*
Authors : Swindell II, J.T.; Chen, L.; Zhu, J.; Ebihara, A.; Shinkai, A.; Kuramitsu, S.; Yokoyama, S.; Fu, Z.-Q.; Chrzas, J.; Rose, J.P.; Wang, B.C.; Southeast Collaboratory for Structural Genomics (SECSG); RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2007-04-10
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	 84% 12% ..
1	B	245	 82% 13% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-oxoacyl-[acyl carrier protein] reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	Total 1841	C 1155	N 335	O 345	S 6	0	0	0
1	B	237	Total 1780	C 1117	N 322	O 336	S 5	0	0	0


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	94	Total 94	O 94	0	0
2	B	94	Total 94	O 94	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-oxoacyl-[acyl carrier protein] reductase

Chain A:  84% 12% ..



- Molecule 1: 3-oxoacyl-[acyl carrier protein] reductase

Chain B:  82% 13% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	76.53Å 76.53Å 137.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.67 – 1.91 19.67 – 1.91	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.67-1.91) 99.9 (19.67-1.91)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	20.81 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.189 , 0.229 0.199 , 0.232	Depositor DCC
R_{free} test set	1763 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtrriage
Anisotropy	0.011	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 28.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.155 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3809	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.44	14/1863 (0.8%)	1.73	24/2517 (1.0%)
1	B	1.18	5/1797 (0.3%)	1.08	6/2426 (0.2%)
All	All	1.31	19/3660 (0.5%)	1.45	30/4943 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	7
1	B	0	1
All	All	1	8

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	189	MET	CA-CB	-19.81	1.20	1.53
1	A	130	ARG	CZ-NH2	-19.03	1.08	1.33
1	A	130	ARG	CD-NE	-16.52	1.23	1.46
1	A	130	ARG	CZ-NH1	-14.97	1.11	1.32
1	A	170	TYR	CB-CG	-12.85	1.23	1.51
1	A	209	ARG	C-N	12.77	1.50	1.33
1	A	54	VAL	CA-CB	-12.40	1.38	1.54
1	B	137	ILE	C-N	-11.94	1.17	1.33
1	A	2	ARG	C-O	8.33	1.34	1.24
1	A	171	ALA	C-N	8.27	1.45	1.33
1	A	2	ARG	N-CA	-7.76	1.35	1.46
1	B	170	TYR	CA-CB	-7.73	1.41	1.53
1	B	126	MET	C-N	-7.42	1.22	1.33
1	A	117	ARG	CD-NE	-6.53	1.37	1.46
1	A	208	GLY	C-N	6.22	1.41	1.33
1	A	3	LYS	CA-CB	-5.66	1.44	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	LYS	CB-CG	-5.25	1.36	1.52
1	B	138	THR	C-N	5.18	1.40	1.33
1	A	116	PHE	C-N	-5.01	1.27	1.33

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	ARG	N-CA-C	33.20	154.33	113.50
1	A	130	ARG	NE-CZ-NH1	31.92	153.42	121.50
1	A	130	ARG	NE-CZ-NH2	-30.94	91.35	119.20
1	A	3	LYS	CB-CA-C	20.55	144.01	109.48
1	A	3	LYS	N-CA-CB	-15.36	85.17	110.52
1	B	138	THR	O-C-N	-14.56	103.23	122.59
1	A	2	ARG	CA-C-N	11.99	140.52	122.65
1	A	2	ARG	C-N-CA	11.99	140.52	122.65
1	A	170	TYR	CA-CB-CG	11.72	135.00	113.90
1	A	170	TYR	CB-CA-C	11.43	129.87	109.99
1	A	4	ALA	N-CA-C	9.46	123.84	108.99
1	A	2	ARG	CB-CA-C	-8.15	94.45	109.29
1	A	3	LYS	CA-CB-CG	8.03	130.15	114.10
1	B	138	THR	CA-C-N	7.65	136.32	121.63
1	B	138	THR	C-N-CA	7.65	136.32	121.63
1	A	2	ARG	CA-C-O	7.45	127.33	119.14
1	B	189	MET	CB-CA-C	7.21	124.76	110.42
1	A	3	LYS	CA-C-N	6.54	133.43	121.85
1	A	3	LYS	C-N-CA	6.54	133.43	121.85
1	A	130	ARG	CD-NE-CZ	6.40	133.36	124.40
1	A	208	GLY	CA-C-N	5.79	131.96	121.89
1	A	208	GLY	C-N-CA	5.79	131.96	121.89
1	B	127	MET	CA-C-N	5.79	128.31	120.44
1	B	127	MET	C-N-CA	5.79	128.31	120.44
1	A	54	VAL	CA-CB-CG2	-5.70	100.72	110.40
1	A	2	ARG	NE-CZ-NH1	5.67	127.17	121.50
1	A	108	LEU	CA-CB-CG	-5.37	97.51	116.30
1	A	205	ILE	N-CA-C	5.32	113.13	107.76
1	A	54	VAL	CG1-CB-CG2	-5.18	99.40	110.80
1	A	170	TYR	N-CA-CB	-5.15	103.13	110.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	2	ARG	CA

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	116	PHE	Mainchain
1	A	130	ARG	Sidechain
1	A	170	TYR	Sidechain
1	A	2	ARG	Peptide,Mainchain
1	A	209	ARG	Mainchain
1	A	3	LYS	Peptide
1	B	170	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1841	0	1887	24	0
1	B	1780	0	1823	40	1
2	A	94	0	0	0	0
2	B	94	0	0	0	0
All	All	3809	0	3710	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:MET:HE1	1:B:134:ILE:HD11	1.35	1.07
1:A:172:GLN:CB	1:A:172:GLN:CD	2.50	0.85
1:B:185:ILE:HG22	1:B:210:PHE:CD1	2.12	0.85
1:B:104:TRP:CZ2	1:B:108:LEU:HD11	2.12	0.85
1:B:185:ILE:HG21	1:B:210:PHE:CE1	2.14	0.82
1:B:127:MET:O	1:B:130:ARG:NH1	2.12	0.81
1:B:185:ILE:CG2	1:B:210:PHE:CD1	2.64	0.80
1:B:104:TRP:CE2	1:B:108:LEU:CD1	2.69	0.75
1:B:104:TRP:CE2	1:B:108:LEU:HD12	2.22	0.74
1:A:2:ARG:HH22	1:A:52:PRO:HG2	1.52	0.73
1:B:11:ARG:HH21	1:B:38:LYS:NZ	1.87	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:VAL:HA	1:B:126:MET:HE3	1.73	0.69
1:B:188:GLU:HG3	1:B:190:THR:H	1.58	0.69
1:B:104:TRP:CZ2	1:B:108:LEU:CD1	2.76	0.69
1:B:119:THR:HG23	1:B:134:ILE:HD13	1.77	0.67
1:A:116:PHE:CD1	1:B:104:TRP:CE3	2.85	0.65
1:B:185:ILE:HG21	1:B:210:PHE:CD1	2.33	0.63
1:B:126:MET:HE1	1:B:134:ILE:CD1	2.21	0.61
1:A:113:SER:O	1:A:116:PHE:HB3	2.01	0.61
1:B:185:ILE:HG22	1:B:210:PHE:HD1	1.63	0.61
1:A:172:GLN:CB	1:A:172:GLN:NE2	2.65	0.60
1:B:189:MET:CB	1:B:189:MET:SD	2.91	0.58
1:B:11:ARG:HH21	1:B:38:LYS:HZ3	1.52	0.56
1:B:104:TRP:NE1	1:B:108:LEU:HD12	2.19	0.56
1:A:116:PHE:CE1	1:B:99:MET:SD	2.99	0.56
1:B:185:ILE:CG2	1:B:210:PHE:CE1	2.85	0.55
1:B:185:ILE:HG21	1:B:210:PHE:HE1	1.69	0.53
1:A:172:GLN:CB	1:A:172:GLN:HE21	2.21	0.53
1:B:122:ALA:C	1:B:126:MET:HE2	2.33	0.52
1:B:189:MET:O	1:B:192:ARG:HB2	2.10	0.51
1:A:104:TRP:CE2	1:A:108:LEU:HD13	2.46	0.51
1:A:116:PHE:HE1	1:B:99:MET:SD	2.34	0.51
1:B:166:VAL:HG12	1:B:177:VAL:HG21	1.91	0.51
1:A:4:ALA:HB2	1:A:83:THR:HB	1.94	0.49
1:A:2:ARG:HG3	1:A:3:LYS:HG3	1.95	0.49
1:B:35:ASN:OD1	1:B:37:GLU:HG3	2.13	0.48
1:A:138:THR:O	1:A:182:PRO:HD2	2.13	0.48
1:A:193:LEU:O	1:A:198:LYS:HE3	2.13	0.48
1:B:185:ILE:CG2	1:B:210:PHE:HD1	2.17	0.47
1:A:7:THR:HA	1:A:31:HIS:HB3	1.97	0.47
1:A:2:ARG:NH2	1:A:52:PRO:HG2	2.27	0.47
1:A:209:ARG:HG2	1:A:210:PHE:O	2.15	0.47
1:B:37:GLU:HG3	1:B:38:LYS:N	2.29	0.46
1:A:53:LEU:C	1:A:54:VAL:HG23	2.42	0.45
1:A:45:GLU:O	1:A:49:ARG:HG3	2.17	0.45
1:A:180:VAL:O	1:A:182:PRO:HD3	2.16	0.45
1:A:163:THR:HG23	1:A:177:VAL:HG12	1.98	0.44
1:A:145:GLY:O	1:B:165:ALA:HA	2.17	0.44
1:B:11:ARG:NH2	1:B:38:LYS:NZ	2.62	0.44
1:A:165:ALA:HA	1:B:145:GLY:O	2.18	0.43
1:B:35:ASN:OD1	1:B:37:GLU:CG	2.66	0.43
1:B:104:TRP:NE1	1:B:108:LEU:CD1	2.79	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:GLU:HG3	1:B:38:LYS:H	1.84	0.43
1:A:32:TYR:HE2	1:A:56:VAL:HG12	1.84	0.42
1:B:64:ALA:HA	1:B:117:ARG:NH2	2.36	0.41
1:B:11:ARG:HH21	1:B:38:LYS:HZ1	1.64	0.41
1:B:15:ARG:NH1	1:B:19:LEU:HD11	2.36	0.41
1:B:123:VAL:HA	1:B:126:MET:CE	2.46	0.41
1:A:127:MET:HE3	1:A:173:ARG:CD	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:ARG:NH1	1:B:191:GLU:OE2[3_775]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/245 (99%)	234 (96%)	9 (4%)	0	100	100
1	B	231/245 (94%)	220 (95%)	10 (4%)	1 (0%)	30	19
All	All	474/490 (97%)	454 (96%)	19 (4%)	1 (0%)	43	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	189	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/183 (100%)	179 (98%)	3 (2%)	55	44
1	B	176/183 (96%)	174 (99%)	2 (1%)	65	60
All	All	358/366 (98%)	353 (99%)	5 (1%)	59	50

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG
1	A	30	ILE
1	A	108	LEU
1	B	56	VAL
1	B	98	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	245	HIS
1	B	86	ASN
1	B	204	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	137:ILE	C	138:THR	N	1.17

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/245 (100%)	-1.46	0 100 100	8, 14, 29, 34	0
1	B	237/245 (96%)	-1.46	0 100 100	7, 15, 28, 37	0
All	All	482/490 (98%)	-1.46	0 100 100	7, 15, 29, 37	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.