



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:59 AM UTC

PDB ID : 1PI3 / pdb_00001pi3
Title : E28Q mutant Benzoylformate Decarboxylase From Pseudomonas Putida
Authors : Bera, A.K.; Hasson, M.S.
Deposited on : 2003-05-29
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	528	23% 91% 7% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Benzoylformate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	523	Total	C	N	O	S	0	7	0
			3937	2486	682	748	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	GLN	GLU	engineered mutation	UNP P20906


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		

- Molecule 4 is 2-{3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-4-METHYL-2-OXO-2,3-DIHYDRO-1,3-THIAZOL-5-YL}ETHYL TRIHYDROGEN DIPHOSPHATE (CCD ID: TZD) (formula: C₁₂H₁₈N₄O₈P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	27	12	4	8	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	504	Total	O	0	0
			504	504		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Benzoylformate decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	80.96Å 95.53Å 137.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.20 20.00 – 1.20	Depositor EDS
% Data completeness (in resolution range)	87.6 (20.00-1.20) 79.3 (20.00-1.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.46 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.121 , 0.136 0.169 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 67.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4472	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TZD, MG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	2/4069 (0.0%)	1.32	35/5560 (0.6%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	246	HIS	CG-ND1	-7.42	1.30	1.38
1	A	72	ALA	N-CA	-5.54	1.39	1.46

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	464	PHE	CA-CB-CG	10.98	124.78	113.80
1	A	382	GLN	OE1-CD-NE2	-10.00	112.60	122.60
1	A	294	ARG	CD-NE-CZ	8.97	136.96	124.40
1	A	443	GLN	OE1-CD-NE2	8.54	131.14	122.60
1	A	120	ARG	NE-CZ-NH1	-7.82	113.68	121.50
1	A	27	ASN	CA-CB-CG	7.72	120.32	112.60
1	A	40	ARG	CA-CB-CG	7.53	129.15	114.10
1	A	13	ARG	CA-CB-CG	7.50	129.09	114.10
1	A	420	ARG	NE-CZ-NH1	7.42	128.92	121.50
1	A	114	ASP	CA-CB-CG	6.97	119.57	112.60
1	A	327	ASN	CA-CB-CG	-6.95	105.65	112.60
1	A	420	ARG	NE-CZ-NH2	-6.85	113.03	119.20
1	A	246	HIS	ND1-CE1-NE2	-6.78	101.62	108.40
1	A	472	ASN	OD1-CG-ND2	6.35	128.95	122.60
1	A	40	ARG	NE-CZ-NH2	-6.11	113.70	119.20
1	A	247	PRO	CA-C-N	6.04	130.81	120.72
1	A	247	PRO	C-N-CA	6.04	130.81	120.72
1	A	247	PRO	O-C-N	-6.01	114.47	122.22
1	A	246	HIS	CG-ND1-CE1	5.91	119.34	109.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	513	ALA	CA-C-N	-5.72	113.49	122.65
1	A	513	ALA	C-N-CA	-5.72	113.49	122.65
1	A	71	SER	CA-C-N	5.66	130.25	120.58
1	A	71	SER	C-N-CA	5.66	130.25	120.58
1	A	120	ARG	CD-NE-CZ	5.55	132.17	124.40
1	A	388	ASN	CA-C-N	5.50	132.34	123.33
1	A	388	ASN	C-N-CA	5.50	132.34	123.33
1	A	162	ASP	CA-CB-CG	5.46	118.06	112.60
1	A	472	ASN	CA-CB-CG	-5.45	107.15	112.60
1	A	382	GLN	CG-CD-OE1	5.23	131.26	120.80
1	A	245	ARG	NE-CZ-NH2	-5.23	114.50	119.20
1	A	428	ASP	CA-CB-CG	5.20	117.80	112.60
1	A	280	TYR	CA-CB-CG	5.18	123.22	113.90
1	A	385	GLN	OE1-CD-NE2	5.09	127.69	122.60
1	A	294	ARG	NE-CZ-NH1	-5.05	116.45	121.50
1	A	246	HIS	ND1-CG-CD2	-5.04	101.06	106.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3937	0	3867	15	0
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	27	0	15	1	0
5	A	504	0	0	4	0
All	All	4472	0	3882	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:GLU:OE1	1:A:343:PRO:HD2	1.88	0.73
1:A:345:LYS:HE2	5:A:693:HOH:O	1.92	0.69
1:A:40:ARG:HG3	5:A:991:HOH:O	2.00	0.61
1:A:195:LYS:HG3	5:A:1010:HOH:O	2.05	0.57
1:A:439:TRP:CZ2	1:A:443:GLN:HG3	2.42	0.55
1:A:439:TRP:CE2	1:A:443:GLN:HG3	2.46	0.50
1:A:464:PHE:O	1:A:467:VAL:HG22	2.14	0.48
1:A:504:LYS:NZ	5:A:712:HOH:O	2.48	0.46
1:A:232:TRP:CZ3	1:A:263:LEU:HD21	2.50	0.46
1:A:232:TRP:CH2	1:A:263:LEU:HD11	2.52	0.44
1:A:119:PRO:HG3	1:A:155:TYR:CG	2.54	0.42
1:A:262:GLN:NE2	1:A:265:GLU:OE1	2.49	0.42
1:A:79[B]:MET:SD	1:A:119:PRO:HA	2.61	0.41
1:A:462:ARG:HH21	1:A:477:ASP:CG	2.29	0.41
1:A:377:THR:OG1	4:A:530:TZD:H5B1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	528/528 (100%)	521 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	416/414 (100%)	414 (100%)	2 (0%)	81 59

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	281	HIS
1	A	342	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	59	GLN
1	A	68	ASN
1	A	151	GLN
1	A	388	ASN
1	A	493	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TZD	A	530	3	27,28,28	2.08	4 (14%)	37,42,42	2.12	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TZD	A	530	3	-	1/17/17/17	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	530	TZD	C2-S1	-7.16	1.69	1.77
4	A	530	TZD	C2-N3	5.80	1.47	1.38
4	A	530	TZD	C5A-C5	2.92	1.56	1.50
4	A	530	TZD	C6'-C5'	2.45	1.42	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	530	TZD	C5-S1-C2	-6.50	88.52	91.64
4	A	530	TZD	OC2-C2-N3	-5.33	119.12	125.37
4	A	530	TZD	C4-C5-S1	4.49	114.73	110.63
4	A	530	TZD	S1-C2-N3	4.21	114.77	110.22
4	A	530	TZD	C4A-C4-N3	-2.66	118.38	122.27
4	A	530	TZD	C6'-N1'-C2'	2.46	120.11	116.07
4	A	530	TZD	C35-N3-C4	2.37	127.99	124.98
4	A	530	TZD	C5A-C5-S1	-2.06	114.87	119.29
4	A	530	TZD	C6'-C5'-C4'	-2.05	113.02	115.55

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	530	TZD	P2-O11-P1-O5G

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	530	TZD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	523/528 (99%)	1.43	120 (22%) 2 3	8, 13, 25, 45	23 (4%)

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	TYR	8.6
1	A	463	TRP	6.9
1	A	464	PHE	6.2
1	A	468	LEU	6.0
1	A	467	VAL	5.9
1	A	339	ALA	5.7
1	A	524	VAL	5.3
1	A	232	TRP	5.3
1	A	338	THR	5.2
1	A	343	PRO	4.7
1	A	466	GLY	4.6
1	A	344	ALA	4.6
1	A	465	ALA	4.5
1	A	353	LEU	4.4
1	A	341	PRO	4.4
1	A	346	VAL	4.4
1	A	263	LEU	4.2
1	A	254	PRO	3.9
1	A	469	GLU	3.8
1	A	514	LYS	3.8
1	A	333	SER	3.8
1	A	351	GLY	3.7
1	A	470	ALA	3.7
1	A	497	ALA	3.6
1	A	513	ALA	3.6
1	A	2	ALA	3.6
1	A	350	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	258	ALA	3.5
1	A	494	ALA	3.4
1	A	511	LEU	3.4
1	A	359	PHE	3.4
1	A	191	ASP	3.4
1	A	249	PHE	3.3
1	A	292	GLY	3.3
1	A	512	SER	3.3
1	A	328	LEU	3.3
1	A	345	LYS	3.3
1	A	169	PRO	3.2
1	A	336	LEU	3.2
1	A	262	GLN	3.2
1	A	223	LEU	3.1
1	A	247	PRO	3.1
1	A	329	VAL	3.1
1	A	17	ILE	3.0
1	A	340	ALA	3.0
1	A	472	ASN	3.0
1	A	194	VAL	3.0
1	A	260	ILE	3.0
1	A	335	GLN	2.9
1	A	337	PRO	2.8
1	A	192	ILE	2.8
1	A	224	ALA	2.8
1	A	495	LEU	2.7
1	A	36	PRO	2.7
1	A	291	PRO	2.7
1	A	197	LEU	2.7
1	A	505	GLY	2.7
1	A	521	VAL	2.7
1	A	248	CYS	2.6
1	A	188	GLN	2.6
1	A	222	MET	2.6
1	A	419	GLU	2.6
1	A	501	GLU	2.6
1	A	510	ALA	2.6
1	A	508	GLN	2.6
1	A	40	ARG	2.5
1	A	364	ASP	2.5
1	A	193	LEU	2.5
1	A	503	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	205	ILE	2.5
1	A	201	SER	2.5
1	A	227	LEU	2.5
1	A	461	LEU	2.5
1	A	366	ALA	2.5
1	A	288	TYR	2.5
1	A	342	GLU	2.4
1	A	384	TRP	2.4
1	A	504	LYS	2.4
1	A	259	ALA	2.4
1	A	190	LEU	2.4
1	A	507	LEU	2.4
1	A	334	ARG	2.4
1	A	354	HIS	2.4
1	A	332	SER	2.4
1	A	198	ASN	2.4
1	A	102	ALA	2.4
1	A	179	VAL	2.3
1	A	522	SER	2.3
1	A	502	GLN	2.3
1	A	229	ALA	2.3
1	A	35	PHE	2.3
1	A	362	LEU	2.3
1	A	355	PRO	2.3
1	A	331	GLU	2.3
1	A	199[A]	SER	2.2
1	A	498	ASP	2.2
1	A	500	LEU	2.2
1	A	330	GLU	2.2
1	A	370	ALA	2.2
1	A	269	VAL	2.2
1	A	324	ALA	2.2
1	A	509	GLU	2.2
1	A	285	PRO	2.2
1	A	479	PRO	2.2
1	A	515	GLY	2.1
1	A	358	VAL	2.1
1	A	290	LYS	2.1
1	A	226	ARG	2.1
1	A	496	LYS	2.1
1	A	397	PHE	2.1
1	A	322	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	326	ALA	2.1
1	A	231	VAL	2.1
1	A	265	GLU	2.1
1	A	367	PRO	2.1
1	A	230	PRO	2.1
1	A	471	GLU	2.0
1	A	387	LEU	2.0
1	A	200	ALA	2.0
1	A	168	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	533	1/1	0.90	0.20	18,18,18,18	1
4	TZD	A	530	27/27	0.98	0.08	9,12,15,17	1
3	CA	A	534	1/1	0.99	0.05	14,14,14,14	1
2	MG	A	531	1/1	0.99	0.04	9,9,9,9	1
3	CA	A	532	1/1	1.00	0.07	10,10,10,10	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.