



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:18 AM UTC

PDB ID : 8PI9 / pdb_00008pi9
Title : DNA binding domain of HNF-1A bound to P2-HNF4A promoter DNA variant (P2 -181G>A)
Authors : Kind, L.; Myllykoski, M.; Raasakka, A.; Kursula, P.
Deposited on : 2023-06-21
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	21	67% 33%
2	F	21	57% 43%
3	A	198	47% 34% 17%
3	B	198	4% 67% 23% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called Chains: E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	21	Total	C	N	O	P	0	0	0
			423	206	70	127	20			

- Molecule 2 is a DNA chain called Chains: F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	21	Total	C	N	O	P	0	0	0
			432	208	83	121	20			

- Molecule 3 is a protein called Hepatocyte nuclear factor 1-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	165	Total	C	N	O	S	0	0	0
			1359	842	262	251	4			
3	B	177	Total	C	N	O	S	0	0	0
			1443	896	274	269	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	GLY	-	expression tag	UNP P20823
B	82	GLY	-	expression tag	UNP P20823

i

- Molecule 1: Chains: E



- | | | | | |
|------|--|------|--|------|
| A401 | | G411 | | T421 |
| T402 | | A412 | | |
| A403 | | G413 | | |
| C404 | | | | |
| G405 | | | | |
| T406 | | | | |

- | | | | | | | | | | | | | | | | | | | |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| I242 | Q243 | Q244 | R244 | S247 | P248 | S249 | Q250 | A251 | L254 | T260 | E261 | V264 | Y265 | N266 | M270 | K273 | A276 | PHE | ARG | HIS | GLY | PRO | PRO | ILE | LEU | LYS | GLU | LEU | GLU | ASN | S93 | P94 | E95 | E96 | A97 | A98 | H99 | I100 | V103 | V104 | L107 | L108 | Q109 | E110 | V115 | K120 | Q125 | H126 | Q130 | R131 | T137 | G138 | L139 | N140 | Q141 | S142 | L144 | S145 | Q146 | H147 | L148 | N149 | K150 | G151 | T152 | P153 | K155 | T156 | Q157 |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|

-
- Figure 1: Amino acid composition of the 100 most abundant peptides in the 1000 most abundant proteins. The figure consists of two horizontal bar charts. The top chart shows the amino acid composition of the 100 most abundant peptides, with the y-axis labeled 'Amino acid' and the x-axis labeled 'Peptide'. The bottom chart shows the amino acid composition of the 1000 most abundant proteins, with the y-axis labeled 'Amino acid' and the x-axis labeled 'Protein'. Both charts use a color-coded system to represent different amino acid groups: Glycine (GLY), Proline (PRO), Prolines (P84, P85, P86, P87, P88, P89, P90, P91, P92, P93, P94, P95, P96, P97, P98, P99, P100), Serine (S121, S122, S123, S124, S125, S126, S127, S128, S129, S130, S131, S132, S133, S134, S135, S136, S137, S138, S139, S140, S141, S142, S143, S144, S145, S146, S147, S148, S149, S150, S151, S152, S153, S154, S155, S156, S157, S158, S159, S160, S161, S162, S163, S164, S165, S166, S167, S168, S169, S170, S171, S172, S173, S174, S175, S176, S177, S178, S179, S180, S181, S182, S183, S184, S185, S186, S187, S188, S189, S190, S191, S192, S193, S194, S195, S196, S197, S198, S199, S200), Alanine (A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A175, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A191, A192, A193, A194, A195, A196, A197, A198, A199, A200), Aspartic acid (D201, D202, D203, D204, D205, D206, D207, D208, D209, D210, D211, D212, D213, D214, D215, D216, D217, D218, D219, D220, D221, D222, D223, D224, D225, D226, D227, D228, D229, D230, D231, D232, D233, D234, D235, D236, D237, D238, D239, D240, D241, D242, D243, D244, D245, D246, D247, D248, D249, D250), Glutamic acid (E261, E262, E263, E264, E265, E266, E267, E268, E269, E270, E271, E272, E273, E274, E275, E276, E277, E278, E279, E280, E281, E282, E283, E284, E285, E286, E287, E288, E289, E290, E291, E292, E293, E294, E295, E296, E297, E298, E299, E300), Lysine (K155, K156, K157, K158, K159, K160, K161, K162, K163, K164, K165, K166, K167, K168, K169, K170, K171, K172, K173, K174, K175, K176, K177, K178, K179, K180, K181, K182, K183, K184, K185, K186, K187, K188, K189, K190, K191, K192, K193, K194, K195, K196, K197, K198, K199, K200), Threonine (T164, T165, T166, T167, T168, T169, T170, T171, T172, T173, T174, T175, T176, T177, T178, T179, T180, T181, T182, T183, T184, T185, T186, T187, T188, T189, T190, T191, T192, T193, T194, T195, T196, T197, T198, T199, T200), Valine (V233, V234, V235, V236, V237, V238, V239, V240, V241, V242, V243, V244, V245, V246, V247, V248, V249, V250), and Isoleucine (I242, I243, I244, I245, I246, I247, I248, I249, I250). The charts show the relative frequency of each amino acid in the peptides and proteins, with the top chart showing a higher frequency of proline and the bottom chart showing a higher frequency of leucine and isoleucine.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.01Å 53.06Å 202.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.64 – 2.80 47.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.5 (47.64-2.80) 84.9 (47.64-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.23 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.234 , 0.293 0.234 , 0.294	Depositor DCC
R_{free} test set	668 reflections (2.69%)	wwPDB-VP
Wilson B-factor (Å ²)	82.7	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 95.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3657	wwPDB-VP
Average B, all atoms (Å ²)	120.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.21	0/472	0.43	0/726
2	F	0.20	0/486	0.39	0/749
3	A	0.20	0/1386	0.43	0/1869
3	B	0.16	0/1471	0.36	0/1983
All	All	0.19	0/3815	0.40	0/5327

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	168	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	423	0	242	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	432	0	239	10	0
3	A	1359	0	1338	65	1
3	B	1443	0	1425	30	1
All	All	3657	0	3244	106	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:125:GLN:HE21	3:A:244:ARG:HA	1.36	0.91
3:A:96:GLU:HB3	3:A:99:HIS:ND1	1.93	0.83
3:B:142:SER:OG	3:B:146:GLN:NE2	2.22	0.73
3:A:168:ARG:HH12	3:A:172:GLU:H	1.36	0.72
3:A:247:SER:HG	3:A:249:SER:HG	1.35	0.70
3:B:136:THR:HG1	3:B:165:TRP:CD1	2.10	0.68
3:A:157:GLN:OE1	3:A:157:GLN:N	2.26	0.67
3:A:126:HIS:HD2	3:A:173:VAL:HG11	1.59	0.66
3:A:168:ARG:O	3:A:168:ARG:NH1	2.29	0.66
3:B:113:TRP:CE2	3:B:117:LYS:HE3	2.32	0.64
1:E:306:DT:OP1	3:B:155:LYS:NZ	2.26	0.63
3:A:94:PRO:HA	3:A:97:ALA:HB2	1.82	0.61
3:A:143:HIS:CE1	3:A:154:MET:HB3	2.36	0.61
3:A:120:LYS:NZ	3:A:149:ASN:OD1	2.32	0.61
1:E:318:DG:H2'	1:E:319:DT:H71	1.83	0.61
3:A:126:HIS:CD2	3:A:173:VAL:HG11	2.37	0.60
3:A:99:HIS:O	3:A:103:VAL:HG23	2.02	0.60
3:A:95:GLU:HG3	3:A:96:GLU:HG3	1.84	0.59
3:A:125:GLN:NE2	3:A:244:ARG:HA	2.14	0.59
3:A:266:ASN:O	3:A:270:ASN:ND2	2.36	0.58
3:B:233:VAL:HG22	3:B:264:VAL:HG21	1.85	0.58
1:E:321:DT:H3	2:F:401:DA:H62	1.50	0.57
3:A:228:GLU:OE1	3:A:228:GLU:N	2.34	0.57
3:A:125:GLN:HG2	3:A:126:HIS:ND1	2.19	0.56
2:F:404:DC:H2'	2:F:405:DG:C8	2.41	0.55
3:A:139:LEU:HD13	3:A:154:MET:HE1	1.88	0.55
3:A:227:GLU:HG3	3:A:228:GLU:OE1	2.06	0.55
2:F:402:DT:H2'	2:F:403:DA:C8	2.42	0.54
3:A:96:GLU:CB	3:A:99:HIS:ND1	2.69	0.54
3:A:270:ASN:H	3:A:270:ASN:HD22	1.55	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:90:GLU:HG2	3:B:175:GLN:OE1	2.07	0.54
3:A:96:GLU:HB3	3:A:99:HIS:CG	2.43	0.54
1:E:318:DG:O6	2:F:403:DA:N6	2.41	0.54
3:B:140:ASN:OD1	3:B:142:SER:N	2.41	0.53
3:A:103:VAL:HG12	3:A:107:LEU:HD11	1.91	0.53
3:A:100:GLN:O	3:A:104:VAL:HG12	2.09	0.53
3:B:87:LYS:HD2	3:B:90:GLU:OE1	2.08	0.53
3:A:169:LYS:O	3:A:173:VAL:HG23	2.10	0.52
3:A:276:ALA:HB3	3:B:150:LYS:HE2	1.91	0.52
3:B:119:VAL:O	3:B:123:LEU:HG	2.10	0.52
3:A:93:SER:O	3:A:93:SER:OG	2.27	0.52
3:A:131:ARG:NH2	3:A:202:ASN:OD1	2.40	0.52
3:A:225:SER:O	3:A:229:ARG:HG3	2.10	0.52
3:A:168:ARG:NH1	3:A:172:GLU:HB2	2.26	0.51
3:A:147:HIS:ND1	3:A:152:THR:O	2.42	0.51
3:A:238:ARG:O	3:A:242:ILE:HG13	2.11	0.51
3:A:130:GLN:NE2	3:A:145:SER:OG	2.34	0.50
3:A:270:ASN:HA	3:A:273:LYS:HG2	1.93	0.50
3:A:147:HIS:HD2	3:A:148:LEU:HD23	1.75	0.50
3:A:104:VAL:O	3:A:108:LEU:HG	2.12	0.50
3:A:140:ASN:OD1	3:A:142:SER:OG	2.25	0.50
3:A:125:GLN:HG2	3:A:126:HIS:CE1	2.47	0.49
3:A:137:THR:OG1	3:A:139:LEU:HG	2.12	0.49
3:B:242:ILE:HD12	3:B:243:GLN:N	2.27	0.49
2:F:412:DA:H2''	2:F:413:DG:C8	2.48	0.48
3:B:86:LEU:HD22	3:B:86:LEU:H	1.78	0.48
3:A:229:ARG:O	3:A:233:VAL:HG23	2.13	0.48
3:B:160:ALA:O	3:B:164:THR:OG1	2.30	0.48
3:A:125:GLN:NE2	3:A:243:GLN:O	2.47	0.47
3:B:112:PRO:HA	3:B:115:VAL:HG22	1.95	0.47
3:A:125:GLN:O	3:A:244:ARG:NH2	2.48	0.47
1:E:319:DT:H2''	1:E:320:DA:C8	2.50	0.46
3:B:165:TRP:CZ2	3:B:169:LYS:HD3	2.50	0.46
3:A:240:GLU:O	3:A:244:ARG:HG2	2.16	0.46
3:B:139:LEU:HD23	3:B:139:LEU:HA	1.65	0.46
1:E:321:DT:H3	2:F:401:DA:N6	2.14	0.46
3:A:229:ARG:HD2	3:A:261:GLU:OE1	2.14	0.46
3:B:92:LEU:HD23	3:B:92:LEU:HA	1.84	0.46
3:A:156:THR:O	3:A:160:ALA:N	2.46	0.45
2:F:411:DG:H2''	2:F:412:DA:C8	2.52	0.45
3:B:177:PHE:CE2	3:B:246:VAL:HG21	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:169:LYS:O	3:B:173:VAL:HG13	2.17	0.44
3:B:227:GLU:O	3:B:230:GLU:HG3	2.16	0.44
3:A:173:VAL:O	3:A:177:PHE:HD2	1.99	0.44
3:B:108:LEU:HD11	3:B:160:ALA:HB2	1.99	0.44
3:A:176:GLN:HG3	3:A:177:PHE:N	2.33	0.44
3:A:120:LYS:HA	3:A:120:LYS:HD2	1.83	0.44
1:E:314:DT:O3'	3:B:205:LYS:NZ	2.50	0.44
3:B:137:THR:HG23	3:B:161:ALA:HB1	2.00	0.44
2:F:406:DT:OP2	3:A:143:HIS:HE1	2.01	0.44
3:A:224:PRO:C	3:A:229:ARG:HH12	2.25	0.44
3:A:104:VAL:HA	3:A:107:LEU:HD12	1.99	0.43
3:A:273:LYS:N	3:A:273:LYS:HD2	2.33	0.43
3:B:204:PHE:HE2	3:B:206:TRP:CE2	2.37	0.43
2:F:403:DA:H2'	2:F:404:DC:C6	2.54	0.43
3:A:234:GLU:O	3:A:238:ARG:HD3	2.19	0.43
3:B:125:GLN:NE2	3:B:243:GLN:O	2.43	0.43
3:B:139:LEU:HD21	3:B:158:LYS:HD3	2.01	0.43
3:A:218:TYR:CZ	3:A:222:LYS:HD2	2.54	0.42
3:A:218:TYR:O	3:A:222:LYS:HD3	2.20	0.42
3:A:150:LYS:HD2	3:A:150:LYS:HA	1.58	0.41
3:A:233:VAL:HG12	3:A:237:ASN:OD1	2.20	0.41
3:A:260:THR:O	3:A:264:VAL:HG23	2.19	0.41
2:F:412:DA:H2''	2:F:413:DG:H8	1.83	0.41
3:A:96:GLU:HB3	3:A:99:HIS:CE1	2.54	0.41
3:A:219:GLU:HA	3:A:222:LYS:HE2	2.02	0.41
3:B:113:TRP:CZ2	3:B:117:LYS:HE3	2.56	0.41
3:A:115:VAL:HG21	3:A:163:TYR:OH	2.20	0.41
1:E:317:DC:H2'	1:E:318:DG:C8	2.55	0.41
3:A:165:TRP:CZ2	3:A:169:LYS:HD3	2.55	0.41
3:A:251:ALA:HA	3:A:254:LEU:HD13	2.02	0.41
3:B:229:ARG:O	3:B:233:VAL:HG23	2.21	0.41
3:B:120:LYS:HZ2	3:B:149:ASN:HA	1.84	0.40
3:B:229:ARG:NH2	3:B:261:GLU:OE2	2.51	0.40
3:A:148:LEU:HD21	3:A:162:LEU:HD11	2.04	0.40
3:A:103:VAL:O	3:A:107:LEU:HG	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:226:LYS:NZ	3:B:105:GLU:OE2[3_545]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	161/198 (81%)	151 (94%)	9 (6%)	1 (1%)	21	51
3	B	173/198 (87%)	168 (97%)	5 (3%)	0	100	100
All	All	334/396 (84%)	319 (96%)	14 (4%)	1 (0%)	36	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	233	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	145/172 (84%)	143 (99%)	2 (1%)	59	85
3	B	154/172 (90%)	153 (99%)	1 (1%)	78	92
All	All	299/344 (87%)	296 (99%)	3 (1%)	68	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	110	GLU
3	A	131	ARG
3	B	179	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
3	A	125	GLN
3	A	126	HIS
3	A	130	GLN
3	A	143	HIS
3	A	146	GLN
3	B	146	GLN
3	B	237	ASN
3	B	270	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	21/21 (100%)	-0.31	0 100 100	90, 109, 134, 155	0
2	F	21/21 (100%)	-0.14	0 100 100	94, 103, 142, 168	0
3	A	165/198 (83%)	0.33	2 (1%) 76 68	88, 126, 166, 216	0
3	B	177/198 (89%)	0.51	8 (4%) 38 30	85, 112, 148, 196	0
All	All	384/438 (87%)	0.35	10 (2%) 57 47	85, 118, 159, 216	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	122	TYR	3.1
3	A	107	LEU	2.9
3	B	84	PRO	2.9
3	B	146	GLN	2.6
3	B	152	THR	2.3
3	B	168	ARG	2.3
3	B	206	TRP	2.2
3	B	177	PHE	2.2
3	B	165	TRP	2.0
3	A	139	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.