



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 26, 2026 – 10:59 PM UTC

PDB ID : 2PJU / pdb_00002pju
Title : Crystal structure of propionate catabolism operon regulatory protein prpR
Authors : Ramagopal, U.A.; Toro, R.; Gilmore, M.; Wu, B.; Koss, J.; Groshong, C.; Gheyi, T.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-04-16
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	9% 71% 11% 17%
1	B	225	8% 73% 6% 18%
1	C	225	4% 69% 11% 16%
1	D	225	10% 75% 15% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6218 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Propionate catabolism operon regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	186	Total	C	N	O	S	0	1	0
			1440	920	242	274	4			
1	B	184	Total	C	N	O	S	0	0	0
			1417	906	239	268	4			
1	C	188	Total	C	N	O	S	0	1	0
			1462	936	248	274	4			
1	D	208	Total	C	N	O	S	0	0	0
			1627	1033	287	303	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP P77743
A	0	SER	-	cloning artifact	UNP P77743
A	1	LEU	-	cloning artifact	UNP P77743
A	96	PHE	ALA	engineered mutation	UNP P77743
A	216	GLU	-	cloning artifact	UNP P77743
A	217	GLY	-	cloning artifact	UNP P77743
A	218	HIS	-	cloning artifact	UNP P77743
A	219	HIS	-	cloning artifact	UNP P77743
A	220	HIS	-	cloning artifact	UNP P77743
A	221	HIS	-	cloning artifact	UNP P77743
A	222	HIS	-	cloning artifact	UNP P77743
A	223	HIS	-	cloning artifact	UNP P77743
B	-1	MET	-	cloning artifact	UNP P77743
B	0	SER	-	cloning artifact	UNP P77743
B	1	LEU	-	cloning artifact	UNP P77743
B	96	PHE	ALA	engineered mutation	UNP P77743
B	216	GLU	-	cloning artifact	UNP P77743
B	217	GLY	-	cloning artifact	UNP P77743
B	218	HIS	-	cloning artifact	UNP P77743
B	219	HIS	-	cloning artifact	UNP P77743
B	220	HIS	-	cloning artifact	UNP P77743

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	221	HIS	-	cloning artifact	UNP P77743
B	222	HIS	-	cloning artifact	UNP P77743
B	223	HIS	-	cloning artifact	UNP P77743
C	-1	MET	-	cloning artifact	UNP P77743
C	0	SER	-	cloning artifact	UNP P77743
C	1	LEU	-	cloning artifact	UNP P77743
C	96	PHE	ALA	engineered mutation	UNP P77743
C	216	GLU	-	cloning artifact	UNP P77743
C	217	GLY	-	cloning artifact	UNP P77743
C	218	HIS	-	cloning artifact	UNP P77743
C	219	HIS	-	cloning artifact	UNP P77743
C	220	HIS	-	cloning artifact	UNP P77743
C	221	HIS	-	cloning artifact	UNP P77743
C	222	HIS	-	cloning artifact	UNP P77743
C	223	HIS	-	cloning artifact	UNP P77743
D	-1	MET	-	cloning artifact	UNP P77743
D	0	SER	-	cloning artifact	UNP P77743
D	1	LEU	-	cloning artifact	UNP P77743
D	96	PHE	ALA	engineered mutation	UNP P77743
D	216	GLU	-	cloning artifact	UNP P77743
D	217	GLY	-	cloning artifact	UNP P77743
D	218	HIS	-	cloning artifact	UNP P77743
D	219	HIS	-	cloning artifact	UNP P77743
D	220	HIS	-	cloning artifact	UNP P77743
D	221	HIS	-	cloning artifact	UNP P77743
D	222	HIS	-	cloning artifact	UNP P77743
D	223	HIS	-	cloning artifact	UNP P77743


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	68	Total O 68 68	0	0
2	B	68	Total O 68 68	0	0
2	C	95	Total O 95 95	0	0
2	D	41	Total O 41 41	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Propionate catabolism operon regulatory protein



- Molecule 1: Propionate catabolism operon regulatory protein




- Molecule 1: Propionate catabolism operon regulatory protein



- Molecule 1: Propionate catabolism operon regulatory protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	35.66Å 58.55Å 103.58Å 85.03° 88.56° 86.94°	Depositor
Resolution (Å)	37.06 – 2.10 37.06 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.8 (37.06-2.10) 97.9 (37.06-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.201 , 0.252 0.229 , 0.270	Depositor DCC
R_{free} test set	2436 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6218	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4748e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/1462	0.96	1/1981 (0.1%)
1	B	0.80	2/1437 (0.1%)	0.94	0/1944
1	C	0.81	0/1484	0.98	0/2010
1	D	0.73	0/1657	0.94	2/2242 (0.1%)
All	All	0.79	2/6040 (0.0%)	0.95	3/8177 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	129	ARG	NE-CZ	6.43	1.40	1.33
1	B	129	ARG	CZ-NH1	5.10	1.39	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	151	ASN	N-CA-C	5.06	118.48	112.72
1	D	116	ILE	CA-C-N	5.05	124.66	119.05
1	D	116	ILE	C-N-CA	5.05	124.66	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1440	0	1465	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1417	0	1457	12	0
1	C	1462	0	1506	24	0
1	D	1627	0	1632	29	0
2	A	68	0	0	2	0
2	B	68	0	0	0	0
2	C	95	0	0	1	0
2	D	41	0	0	2	0
All	All	6218	0	6060	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:20:THR:HG23	1:C:21:ARG:H	1.36	0.90
1:C:21:ARG:HD3	2:C:303:HOH:O	1.82	0.79
1:D:90:GLY:HA2	1:D:119:LEU:HG	1.67	0.76
1:D:27:ARG:HG2	1:D:27:ARG:HH11	1.51	0.75
1:A:95:GLN:O	1:A:99:LYS:HD2	1.86	0.73
1:B:95:GLN:HE21	1:B:95:GLN:HA	1.55	0.71
1:B:35:HIS:CD2	1:B:35:HIS:H	2.08	0.71
1:B:133:ARG:HB3	1:B:144:GLN:HE21	1.56	0.71
1:C:27:ARG:HH11	1:C:27:ARG:HG2	1.55	0.70
1:D:116:ILE:O	1:D:120:VAL:HG23	1.92	0.70
1:A:96:PHE:CE1	1:A:178:SER:HB2	2.28	0.69
1:D:96:PHE:CE1	2:D:237:HOH:O	2.47	0.67
1:C:98:ALA:HA	1:C:103:LEU:CD2	2.25	0.67
1:D:137:THR:HG23	1:D:140:ASP:H	1.61	0.65
1:A:12:PRO:HB3	1:A:193:THR:HG21	1.79	0.64
1:C:20:THR:HG23	1:C:21:ARG:N	2.12	0.63
1:C:35:HIS:H	1:C:35:HIS:CD2	2.16	0.63
1:D:84:ILE:HD13	1:D:189:ALA:HB2	1.81	0.62
1:B:133:ARG:HB3	1:B:144:GLN:NE2	2.14	0.62
1:D:99:LYS:HG3	1:D:100:ALA:N	2.16	0.60
1:A:47:GLU:HB2	2:A:235:HOH:O	2.00	0.60
1:C:105:SER:HB3	1:C:154:GLU:HG3	1.84	0.60
1:C:19:VAL:HG23	1:C:20:THR:HG22	1.84	0.59
1:D:137:THR:HG22	1:D:140:ASP:CG	2.27	0.59
1:A:163:THR:HG21	1:A:175:PHE:HA	1.87	0.56
1:A:98:ALA:HB2	1:A:103:LEU:HD22	1.86	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ARG:HD2	1:A:144:GLN:OE1	2.06	0.55
1:C:98:ALA:HA	1:C:103:LEU:HD22	1.89	0.54
1:C:99:LYS:NZ	1:C:99:LYS:HB2	2.24	0.52
1:D:189:ALA:O	1:D:193:THR:HG23	2.09	0.52
1:A:30:SER:HB2	1:A:39:ILE:HD13	1.92	0.52
1:D:93:VAL:HG11	1:D:119:LEU:HD11	1.92	0.52
1:D:137:THR:CG2	1:D:140:ASP:H	2.23	0.51
1:C:141:ALA:O	1:C:145:ILE:HG23	2.11	0.50
1:D:89:SER:OG	1:D:92:ASP:HB2	2.11	0.50
1:C:20:THR:O	1:C:24:GLU:HG2	2.12	0.49
1:C:35:HIS:H	1:C:35:HIS:HD2	1.61	0.49
1:C:145:ILE:HD11	1:C:169:ALA:HB2	1.94	0.49
1:D:96:PHE:O	1:D:99:LYS:HG3	2.13	0.49
1:B:18:SER:O	1:B:43:GLN:HA	2.13	0.49
1:C:98:ALA:HB2	1:C:103:LEU:HD21	1.96	0.48
1:D:211:THR:HG23	1:D:214:VAL:H	1.78	0.48
1:C:196:SER:OG	1:D:199:HIS:HE1	1.96	0.48
1:D:28:ASP:O	1:D:31:LEU:HB2	2.14	0.48
1:C:130:LEU:HD22	1:C:132:GLN:HG2	1.96	0.47
1:C:125:THR:HG21	1:D:103:LEU:HB3	1.96	0.47
1:D:211:THR:CG2	1:D:214:VAL:HG23	2.45	0.47
1:A:151:ASN:HD22	1:A:151:ASN:C	2.23	0.47
1:B:95:GLN:HA	1:B:95:GLN:NE2	2.26	0.47
1:B:35:HIS:H	1:B:35:HIS:HD2	1.58	0.46
1:C:27:ARG:HH11	1:C:27:ARG:CG	2.26	0.46
1:D:104:THR:O	1:D:104:THR:HG22	2.16	0.46
1:A:96:PHE:HE1	2:A:246:HOH:O	1.98	0.45
1:C:130:LEU:HD22	1:C:132:GLN:CG	2.46	0.45
1:D:96:PHE:O	1:D:99:LYS:CG	2.64	0.45
1:C:96:PHE:CE2	1:C:178:SER:HB3	2.53	0.44
1:D:99:LYS:HG3	1:D:100:ALA:H	1.82	0.44
1:D:27:ARG:HH11	1:D:27:ARG:CG	2.26	0.43
1:A:195:MET:SD	1:B:192:MET:HG2	2.57	0.43
1:C:102:LYS:HG2	1:C:155:ALA:HB2	2.01	0.43
1:C:104:THR:O	1:C:129:ARG:NH1	2.52	0.42
1:D:96:PHE:HA	1:D:99:LYS:HG2	2.01	0.42
1:D:212:ARG:O	1:D:216:GLU:HB2	2.19	0.42
1:D:113:GLN:HG3	1:D:136:ILE:HD11	2.02	0.42
1:A:88:PRO:HG3	1:A:177:TYR:CD1	2.55	0.42
1:A:35:HIS:H	1:A:35:HIS:CD2	2.38	0.42
1:C:18:SER:O	1:C:43:GLN:HA	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:92:ASP:OD2	1:D:178:SER:OG	2.23	0.42
1:B:47:GLU:H	1:B:47:GLU:CD	2.28	0.41
1:A:88:PRO:HG3	1:A:177:TYR:CE1	2.55	0.41
1:B:123:GLN:HE21	1:B:123:GLN:HB3	1.59	0.41
1:D:96:PHE:HE1	2:D:237:HOH:O	1.93	0.41
1:A:85:LEU:CD2	1:A:87:LYS:HE2	2.50	0.41
1:D:102:LYS:O	1:D:107:ILE:HD11	2.20	0.41
1:B:192:MET:HB2	1:B:192:MET:HE3	1.83	0.41
1:D:191:ASP:OD1	1:D:194:ARG:NH2	2.52	0.41
1:B:99:LYS:HD3	1:B:99:LYS:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/225 (82%)	180 (97%)	5 (3%)	0	100	100
1	B	180/225 (80%)	176 (98%)	4 (2%)	0	100	100
1	C	187/225 (83%)	182 (97%)	5 (3%)	0	100	100
1	D	204/225 (91%)	196 (96%)	8 (4%)	0	100	100
All	All	756/900 (84%)	734 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/188 (81%)	147 (96%)	6 (4%)	28	31
1	B	151/188 (80%)	137 (91%)	14 (9%)	8	6
1	C	156/188 (83%)	142 (91%)	14 (9%)	9	6
1	D	172/188 (92%)	160 (93%)	12 (7%)	14	11
All	All	632/752 (84%)	586 (93%)	46 (7%)	13	10

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LEU
1	A	84	ILE
1	A	96	PHE
1	A	128	LEU
1	A	139	GLU
1	A	151	ASN
1	B	35	HIS
1	B	47	GLU
1	B	48	LYS
1	B	95	GLN
1	B	97	LEU
1	B	99	LYS
1	B	115	THR
1	B	120	VAL
1	B	123	GLN
1	B	124	LYS
1	B	130	LEU
1	B	148	LEU
1	B	192	MET
1	B	197	LEU
1	C	24	GLU
1	C	25	LEU
1	C	27	ARG
1	C	61	ARG
1	C	80	SER
1	C	85	LEU
1	C	99	LYS
1	C	103	LEU
1	C	123	GLN
1	C	124	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	129	ARG
1	C	130	LEU
1	C	145	ILE
1	C	154	GLU
1	D	27	ARG
1	D	31	LEU
1	D	47	GLU
1	D	70	SER
1	D	96	PHE
1	D	99	LYS
1	D	123	GLN
1	D	129	ARG
1	D	149	LYS
1	D	165	LEU
1	D	211	THR
1	D	215	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	HIS
1	A	151	ASN
1	B	35	HIS
1	B	43	GLN
1	B	95	GLN
1	B	123	GLN
1	B	144	GLN
1	C	35	HIS
1	C	113	GLN
1	C	123	GLN
1	D	43	GLN
1	D	199	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	186/225 (82%)	0.80	20 (10%)	11 11	20, 38, 48, 55	1 (0%)
1	B	184/225 (81%)	0.81	18 (9%)	13 13	32, 39, 47, 52	0
1	C	188/225 (83%)	0.77	8 (4%)	40 41	20, 38, 46, 53	1 (0%)
1	D	208/225 (92%)	0.87	22 (10%)	11 12	30, 39, 51, 58	0
All	All	766/900 (85%)	0.81	68 (8%)	15 16	20, 38, 49, 58	2 (0%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	19	VAL	4.4
1	D	202	HIS	4.0
1	D	117	PRO	3.9
1	D	100	ALA	3.8
1	A	126	PHE	3.8
1	C	198	ARG	3.6
1	D	204	ALA	3.4
1	B	124	LYS	3.4
1	A	127	ASN	3.2
1	D	205	THR	3.2
1	B	104	THR	3.1
1	A	96	PHE	3.0
1	A	88	PRO	3.0
1	A	120	VAL	3.0
1	D	125	THR	2.9
1	C	127	ASN	2.9
1	A	92[A]	ASP	2.8
1	C	19	VAL	2.8
1	A	150	ALA	2.8
1	B	80	SER	2.8
1	C	96	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	GLY	2.7
1	A	121	ALA	2.7
1	D	150	ALA	2.7
1	B	151	ASN	2.7
1	B	150	ALA	2.7
1	A	34	ASP	2.7
1	A	55	LYS	2.7
1	D	20	THR	2.6
1	B	34	ASP	2.6
1	D	196	SER	2.6
1	A	104	THR	2.6
1	A	194	ARG	2.6
1	A	58	ALA	2.6
1	C	117	PRO	2.6
1	A	24	GLU	2.6
1	D	58	ALA	2.6
1	D	215	LEU	2.6
1	B	96	PHE	2.5
1	B	140	ASP	2.5
1	B	121	ALA	2.5
1	D	94	LEU	2.5
1	D	91	TYR	2.4
1	C	126	PHE	2.4
1	A	196	SER	2.4
1	D	139	GLU	2.3
1	D	121	ALA	2.3
1	D	197	LEU	2.3
1	D	120	VAL	2.3
1	B	125	THR	2.3
1	B	154	GLU	2.2
1	C	20	THR	2.2
1	D	88	PRO	2.2
1	B	147	GLU	2.2
1	D	34	ASP	2.2
1	A	87	LYS	2.1
1	A	106	SER	2.1
1	C	130	LEU	2.1
1	B	123	GLN	2.1
1	D	199	HIS	2.1
1	A	86	ILE	2.1
1	A	124	LYS	2.1
1	B	148	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	19	VAL	2.1
1	A	80	SER	2.1
1	B	131	ASP	2.0
1	B	117	PRO	2.0
1	D	127	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.