



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:01 AM UTC

PDB ID : 1PKH / pdb_00001pkh
Title : STRUCTURAL BASIS FOR RECOGNITION AND CATALYSIS BY
THE BIFUNCTIONAL DCTP DEAMINASE AND DUTPASE FROM
METHANOCOCCUS JANNASCHII
Authors : Huffman, J.L.; Li, H.; White, R.H.; Tainer, J.A.
Deposited on : 2003-06-05
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	14% 75% 11% • 11%
1	B	204	9% 68% 15% • 14%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional deaminase/diphosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	182	Total	C	N	O	S	0	0	0
			1472	953	235	281	3			
1	B	175	Total	C	N	O	S	0	0	0
			1414	918	224	269	3			

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	182	Total O 182 182	0	0
3	B	179	Total O 179 179	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional deaminase/diphosphatase

Chain A: 



- Molecule 1: Bifunctional deaminase/diphosphatase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	110.70Å 110.70Å 110.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 1.42 45.00 – 1.42	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.00-1.42) 95.7 (45.00-1.42)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.53 (at 1.42Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.185 , 0.203 0.181 , 0.196	Depositor DCC
R_{free} test set	4251 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	15.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3283	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/1501	1.50	17/2029 (0.8%)
1	B	0.72	0/1442	1.49	16/1951 (0.8%)
All	All	0.71	0/2943	1.50	33/3980 (0.8%)

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	89	ASP	CA-CB-CG	-7.72	104.88	112.60
1	B	43	TYR	CA-C-N	-6.83	110.74	122.14
1	B	43	TYR	C-N-CA	-6.83	110.74	122.14
1	A	177	TYR	CA-C-O	-6.78	114.01	121.33
1	B	132	GLY	CA-C-N	6.71	130.37	120.90
1	B	132	GLY	C-N-CA	6.71	130.37	120.90
1	B	132	GLY	O-C-N	-6.60	114.73	122.45
1	A	128	HIS	O-C-N	-6.52	116.05	123.48
1	B	5	ASP	CA-CB-CG	6.08	118.68	112.60
1	A	127	SER	O-C-N	6.07	129.54	122.19
1	B	45	ASP	O-C-N	-5.98	116.24	122.94
1	A	16	ARG	NE-CZ-NH2	-5.96	113.83	119.20
1	B	133	TRP	N-CA-CB	5.77	118.86	109.69
1	A	16	ARG	NH1-CZ-NH2	5.66	126.65	119.30
1	B	82	PHE	CA-CB-CG	5.65	119.45	113.80
1	B	45	ASP	CA-CB-CG	5.58	118.18	112.60
1	A	117	ARG	CA-C-N	5.47	127.62	120.28
1	A	117	ARG	C-N-CA	5.47	127.62	120.28
1	A	5	ASP	CA-CB-CG	5.43	118.03	112.60
1	A	102	GLU	CA-C-N	5.39	130.15	122.72
1	A	102	GLU	C-N-CA	5.39	130.15	122.72
1	B	10	ASP	CA-CB-CG	5.38	117.98	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177	TYR	O-C-N	5.34	129.33	123.25
1	B	17	ILE	O-C-N	5.27	128.78	123.20
1	B	171	SER	CA-C-O	-5.26	115.50	120.34
1	A	133	TRP	CA-C-O	5.20	126.91	121.19
1	A	178	SER	CA-C-O	-5.16	114.73	120.66
1	A	159	GLN	O-C-N	5.14	128.93	122.86
1	B	158	ASN	CB-CG-ND2	5.14	124.11	116.40
1	A	179	GLU	CA-C-N	-5.06	115.87	123.00
1	A	179	GLU	C-N-CA	-5.06	115.87	123.00
1	B	80	ASN	CA-C-O	5.04	125.90	120.55
1	A	135	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1472	0	1487	9	0
1	B	1414	0	1431	20	0
2	A	12	0	18	0	0
2	B	24	0	36	10	0
3	A	182	0	0	3	0
3	B	179	0	0	6	0
All	All	3283	0	2972	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ARG:HD3	3:B:355:HOH:O	1.91	0.69
1:B:117:ARG:HD2	3:B:368:HOH:O	2.01	0.60
1:B:35:THR:HG21	2:B:210:EDO:H22	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:GLU:HG2	2:B:205:EDO:O2	2.03	0.59
1:A:62:LYS:HD3	1:A:89:ASP:OD1	2.03	0.58
1:B:60:LYS:HE2	3:B:338:HOH:O	2.07	0.55
1:A:62:LYS:HD3	1:A:89:ASP:CG	2.34	0.52
1:B:50:LEU:HD13	3:B:571:HOH:O	2.09	0.52
1:B:28:GLY:CA	2:B:206:EDO:H12	2.40	0.52
1:A:62:LYS:HG3	3:A:560:HOH:O	2.11	0.50
1:B:41:ILE:HD12	2:B:207:EDO:H12	1.95	0.49
1:B:43:TYR:HD2	1:B:48:TYR:HH	1.55	0.49
1:B:45:ASP:OD1	1:B:55:ASN:HB2	2.12	0.49
2:B:210:EDO:H11	3:B:501:HOH:O	2.12	0.49
1:B:33:ASP:OD2	2:B:206:EDO:H21	2.12	0.49
1:A:25:ASP:HB3	3:A:389:HOH:O	2.13	0.48
1:B:70:LEU:O	1:B:71:ASN:HB2	2.12	0.48
1:A:129:GLN:NE2	3:A:491:HOH:O	2.47	0.47
1:B:158:ASN:HD21	2:B:209:EDO:H22	1.78	0.47
1:B:158:ASN:ND2	2:B:209:EDO:H22	2.30	0.47
1:B:77:GLU:O	1:B:77:GLU:HG2	2.13	0.46
1:B:20:LYS:HE3	3:B:568:HOH:O	2.15	0.46
1:B:28:GLY:HA3	2:B:206:EDO:H12	1.98	0.45
1:B:59:ILE:HD12	1:B:90:TYR:CE1	2.53	0.43
1:B:35:THR:OG1	2:B:210:EDO:H22	2.19	0.43
1:A:179:GLU:C	1:A:180:ARG:HD3	2.44	0.42
1:A:117:ARG:NH1	1:A:160:ARG:HB2	2.34	0.42
1:A:128:HIS:CE1	1:A:132:GLY:HA2	2.55	0.41
1:B:42:ILE:CD1	1:B:92:VAL:HG21	2.51	0.41
1:A:113:GLN:OE1	1:A:131:ALA:O	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/204 (88%)	173 (96%)	7 (4%)	0	100	100
1	B	173/204 (85%)	167 (96%)	6 (4%)	0	100	100
All	All	353/408 (86%)	340 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/186 (89%)	156 (94%)	9 (6%)	19	1
1	B	159/186 (86%)	152 (96%)	7 (4%)	25	3
All	All	324/372 (87%)	308 (95%)	16 (5%)	22	2

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LYS
1	A	47	VAL
1	A	57	LYS
1	A	62	LYS
1	A	77	GLU
1	A	84	GLU
1	A	180	ARG
1	A	181	LYS
1	A	182	THR
1	B	53	GLU
1	B	57	LYS
1	B	76	GLU
1	B	77	GLU
1	B	115	GLN
1	B	128	HIS
1	B	129	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	63	ASN
1	A	101	ASN
1	A	113	GLN
1	A	129	GLN
1	A	158	ASN
1	B	55	ASN
1	B	101	ASN
1	B	113	GLN
1	B	128	HIS
1	B	158	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	210	-	3,3,3	0.55	0	2,2,2	0.58	0
2	EDO	B	207	-	3,3,3	0.53	0	2,2,2	0.89	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	206	-	3,3,3	0.42	0	2,2,2	0.33	0
2	EDO	B	209	-	3,3,3	0.62	0	2,2,2	1.16	0
2	EDO	B	205	-	3,3,3	0.79	0	2,2,2	0.74	0
2	EDO	A	207	-	3,3,3	0.56	0	2,2,2	0.73	0
2	EDO	B	208	-	3,3,3	0.58	0	2,2,2	0.56	0
2	EDO	A	205	-	3,3,3	0.65	0	2,2,2	0.74	0
2	EDO	B	206	-	3,3,3	0.71	0	2,2,2	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	210	-	-	0/1/1/1	-
2	EDO	B	207	-	-	0/1/1/1	-
2	EDO	A	206	-	-	0/1/1/1	-
2	EDO	B	209	-	-	1/1/1/1	-
2	EDO	B	205	-	-	1/1/1/1	-
2	EDO	A	207	-	-	0/1/1/1	-
2	EDO	B	208	-	-	1/1/1/1	-
2	EDO	A	205	-	-	1/1/1/1	-
2	EDO	B	206	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	205	EDO	O1-C1-C2-O2
2	B	205	EDO	O1-C1-C2-O2
2	B	209	EDO	O1-C1-C2-O2
2	B	206	EDO	O1-C1-C2-O2
2	B	208	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	210	EDO	3	0
2	B	207	EDO	1	0
2	B	209	EDO	2	0
2	B	205	EDO	1	0
2	B	206	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	182/204 (89%)	0.56	29 (15%) 5 5	12, 17, 44, 102	0
1	B	175/204 (85%)	0.38	19 (10%) 10 12	12, 16, 37, 62	0
All	All	357/408 (87%)	0.47	48 (13%) 7 8	12, 16, 41, 102	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	175	VAL	11.3
1	A	182	THR	5.0
1	A	180	ARG	4.3
1	A	90	TYR	4.1
1	A	72	TYR	3.9
1	A	177	TYR	3.6
1	B	72	TYR	3.5
1	B	76	GLU	3.5
1	A	181	LYS	3.4
1	A	130	THR	3.2
1	B	6	LYS	3.2
1	A	77	GLU	3.2
1	B	56	TYR	3.2
1	A	175	VAL	3.1
1	A	58	ARG	3.1
1	A	56	TYR	3.1
1	A	133	TRP	3.1
1	A	179	GLU	3.1
1	B	73	ASN	3.0
1	B	174	ASP	3.0
1	A	73	ASN	2.9
1	A	62	LYS	2.9
1	A	78	LYS	2.9
1	A	1	MET	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	71	ASN	2.9
1	B	130	THR	2.9
1	B	77	GLU	2.9
1	A	81	TYR	2.8
1	A	84	GLU	2.6
1	B	43	TYR	2.5
1	A	128	HIS	2.5
1	B	53	GLU	2.4
1	A	57	LYS	2.4
1	A	59	ILE	2.4
1	A	176	GLY	2.4
1	B	71	ASN	2.4
1	A	129	GLN	2.3
1	B	57	LYS	2.2
1	B	59	ILE	2.2
1	A	178	SER	2.1
1	B	44	ASP	2.1
1	A	55	ASN	2.1
1	A	131	ALA	2.1
1	B	42	ILE	2.1
1	B	133	TRP	2.1
1	A	79	ILE	2.0
1	B	129	GLN	2.0
1	B	74	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	B	205	4/4	0.73	0.18	24,31,35,48	0
2	EDO	B	206	4/4	0.79	0.19	37,39,40,55	0
2	EDO	B	210	4/4	0.82	0.24	53,56,58,69	0
2	EDO	B	209	4/4	0.83	0.20	26,34,38,38	0
2	EDO	A	205	4/4	0.85	0.18	37,41,42,42	0
2	EDO	A	207	4/4	0.85	0.15	37,38,45,59	0
2	EDO	A	206	4/4	0.91	0.10	23,24,26,31	0
2	EDO	B	207	4/4	0.93	0.12	25,29,31,37	0
2	EDO	B	208	4/4	0.93	0.11	25,27,27,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.