



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 11:02 PM UTC

PDB ID : 4PKD / pdb_00004pkd
Title : U1-70k in complex with U1 snRNA stem-loops 1 and U1-A RRM in complex with stem-loop 2
Authors : Oubridge, C.; Kondo, Y.; van Roon, A.M.; Nagai, K.
Deposited on : 2014-05-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)
RNA backbone	3983	1003 (2.78-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	V	55	
2	B	279	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called U1 snRNA stem-loops 1 and 2 (55-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	V	55	Total	C	N	O	P	0	0	0
			1177	522	205	393	57			

- Molecule 2 is a protein called U1 small nuclear ribonucleoprotein A,U1 small nuclear ribonucleoprotein 70 kDa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	0	0
			1954	1226	368	352	8			


There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	GLY	ALA	engineered mutation	UNP Q06AA4
B	112	GLY	-	linker	UNP Q06AA4
B	113	SER	-	linker	UNP Q06AA4
B	114	GLY	-	linker	UNP Q06AA4
B	115	SER	-	linker	UNP Q06AA4
B	116	GLY	-	linker	UNP Q06AA4
B	117	SER	-	linker	UNP Q06AA4
B	118	GLY	-	linker	UNP Q06AA4
B	119	SER	-	linker	UNP Q06AA4
B	120	GLY	-	linker	UNP Q06AA4
B	121	SER	-	linker	UNP Q06AA4
B	122	GLY	-	linker	UNP Q06AA4
B	123	SER	-	linker	UNP Q06AA4
B	124	GLY	-	linker	UNP Q06AA4

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	V	2	Total Mg 2 2	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C N 5 3 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	V	44	Total O 44 44	0	0
5	B	69	Total O 69 69	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: U1 snRNA stem-loops 1 and 2 (55-MER)



- Molecule 2: U1 small nuclear ribonucleoprotein A, U1 small nuclear ribonucleoprotein 70 kDa



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.22Å 66.60Å 93.73Å 90.00° 110.95° 90.00°	Depositor
Resolution (Å)	87.00 – 2.50 87.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (87.00-2.50) 99.1 (87.00-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.197 , 0.258 0.201 , 0.255	Depositor DCC
R_{free} test set	812 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.507	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3252	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, IMD, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	V	0.79	1/1278 (0.1%)	1.02	4/1988 (0.2%)
2	B	1.53	15/1991 (0.8%)	1.28	9/2664 (0.3%)
All	All	1.30	16/3269 (0.5%)	1.18	13/4652 (0.3%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	241	VAL	C-O	-8.03	1.15	1.24
2	B	207	LYS	CA-C	7.00	1.60	1.53
2	B	169	THR	C-O	-6.54	1.16	1.24
2	B	257	LEU	N-CA	-6.28	1.38	1.46
2	B	194	ILE	C-O	-6.00	1.17	1.24
1	V	100	A	O3'-P	-5.86	1.52	1.61
2	B	175	VAL	CA-C	5.65	1.59	1.52
2	B	54	GLN	CA-C	5.61	1.59	1.52
2	B	137	ARG	C-O	-5.61	1.17	1.24
2	B	168	LYS	C-O	5.60	1.31	1.24
2	B	187	GLU	C-O	-5.55	1.17	1.24
2	B	148	GLU	C-O	-5.33	1.17	1.24
2	B	190	VAL	C-O	-5.27	1.17	1.24
2	B	230	ASP	C-O	-5.20	1.17	1.23
2	B	245	ARG	CA-C	-5.03	1.46	1.52
2	B	193	PRO	CA-C	5.02	1.58	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	161	ASN	N-CA-C	6.91	119.66	111.71
2	B	256	ARG	CG-CD-NE	-6.35	98.03	112.00
1	V	71	C	C2'-C3'-O3'	-6.12	104.52	113.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8	PRO	N-CA-C	6.09	120.79	111.11
2	B	251	GLY	N-CA-C	5.99	123.32	115.47
2	B	253	ARG	CA-C-N	-5.96	114.49	120.21
2	B	253	ARG	C-N-CA	-5.96	114.49	120.21
2	B	182	SER	N-CA-C	5.74	117.21	111.07
2	B	186	ARG	N-CA-C	5.50	117.71	111.11
1	V	102	G	C2'-C3'-O3'	-5.39	105.61	113.70
1	V	73	C	C2'-C3'-O3'	-5.33	105.71	113.70
1	V	63	C	C3'-C2'-O2'	5.20	118.50	110.70
2	B	98	LYS	N-CA-C	5.06	118.72	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	V	1177	0	592	34	0
2	B	1954	0	1939	20	0
3	B	1	0	0	0	0
3	V	2	0	0	0	0
4	B	5	0	5	0	0
5	B	69	0	0	2	0
5	V	44	0	0	0	0
All	All	3252	0	2536	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:V:73:C:H2'	1:V:74:C:H6	1.44	0.81
1:V:73:C:H2'	1:V:74:C:C6	2.16	0.80
1:V:87:U:H6	1:V:87:U:H5''	1.46	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:V:73:C:C2	1:V:74:C:C5	2.70	0.78
2:B:33:ILE:HG23	2:B:72:MET:HE2	1.70	0.72
2:B:127:THR:OG1	2:B:130:GLU:HG3	1.96	0.65
2:B:92:ASP:O	2:B:96:LYS:HG3	1.97	0.62
1:V:73:C:O2	1:V:74:C:C5	2.53	0.62
2:B:75:PHE:HB3	2:B:82:MET:HE3	1.83	0.61
1:V:73:C:C2'	1:V:74:C:C6	2.83	0.60
1:V:72:U:H5''	1:V:73:C:OP1	2.01	0.60
1:V:111:C:H6	1:V:111:C:H5''	1.67	0.59
1:V:82:G:O2'	1:V:83:G:H5'	2.02	0.59
1:V:73:C:C2'	1:V:74:C:H6	2.13	0.58
1:V:82:G:C2'	1:V:83:G:H5'	2.34	0.58
2:B:151:LEU:HD12	2:B:257:LEU:HD13	1.85	0.58
1:V:73:C:H3'	1:V:74:C:C5	2.39	0.57
1:V:73:C:C2	1:V:74:C:C6	2.94	0.54
1:V:73:C:H2'	1:V:74:C:O4'	2.07	0.54
1:V:111:C:H5''	1:V:111:C:C6	2.41	0.54
2:B:140:ILE:O	2:B:144:GLN:HG3	2.07	0.54
2:B:33:ILE:O	2:B:36:GLN:HG2	2.09	0.52
1:V:73:C:N3	1:V:74:C:C4	2.79	0.51
1:V:60:G:O2'	1:V:61:A:O4'	2.29	0.51
2:B:33:ILE:CG2	2:B:72:MET:HE2	2.38	0.51
2:B:150:GLU:HA	2:B:153:MET:CE	2.41	0.51
1:V:112:C:O2'	1:V:113:U:H5'	2.12	0.50
2:B:66:THR:O	2:B:70:ARG:HG2	2.12	0.50
1:V:87:U:H5''	1:V:87:U:C6	2.37	0.49
2:B:93:ILE:O	2:B:97:MET:HG3	2.14	0.47
1:V:73:C:O2'	1:V:74:C:H5'	2.14	0.47
1:V:72:U:C5'	1:V:73:C:OP1	2.62	0.47
1:V:87:U:H6	1:V:87:U:C5'	2.21	0.46
1:V:60:G:HO2'	1:V:61:A:H8	1.64	0.46
1:V:74:C:H6	1:V:74:C:O5'	1.99	0.46
2:B:75:PHE:HB3	2:B:82:MET:CE	2.46	0.45
2:B:141:GLU:HA	2:B:144:GLN:CD	2.41	0.45
1:V:112:C:C2'	1:V:113:U:H5'	2.48	0.44
1:V:73:C:H3'	1:V:74:C:H5	1.83	0.44
2:B:75:PHE:HD2	2:B:82:MET:HE2	1.82	0.43
2:B:89:THR:OG1	5:B:468:HOH:O	2.21	0.43
1:V:60:G:H8	1:V:60:G:OP2	2.02	0.43
1:V:73:C:C2	1:V:74:C:C4	3.06	0.43
2:B:150:GLU:HA	2:B:153:MET:HE3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:V:101:A:H2'	1:V:102:G:O4'	2.17	0.42
2:B:199:MET:HE2	5:B:431:HOH:O	2.20	0.42
2:B:174:ARG:O	2:B:237:ARG:HD3	2.20	0.42
1:V:85:G:C2'	1:V:86:A:H5'	2.50	0.41
2:B:154:TRP:CZ2	2:B:156:PRO:HG3	2.55	0.41
2:B:50:LYS:HG2	2:B:51:MET:SD	2.60	0.41
1:V:73:C:O2	1:V:73:C:H5'	2.20	0.40
1:V:73:C:H3'	1:V:74:C:C6	2.55	0.40
1:V:78:U:H2'	1:V:79:C:O4'	2.21	0.40
1:V:87:U:C6	1:V:87:U:C5'	3.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	236/279 (85%)	230 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	202/237 (85%)	189 (94%)	13 (6%)	16	33

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	8	PRO
2	B	39	GLN
2	B	61	GLU
2	B	103	GLU
2	B	135	LYS
2	B	147	VAL
2	B	151	LEU
2	B	189	GLU
2	B	205	SER
2	B	232	LYS
2	B	233	LYS
2	B	237	ARG
2	B	253	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	145	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	V	53/55 (96%)	13 (24%)	0

All (13) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	V	72	U
1	V	73	C
1	V	75	G
1	V	81	A
1	V	85	G
1	V	86	A
1	V	87	U
1	V	88	A
1	V	93	G
1	V	94	A
1	V	100	A
1	V	106	G

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	V	113	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	IMD	B	302	-	5,5,5	0.90	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	B	302	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	V	54/55 (98%)	-0.01	1 (1%) 66 62	27, 56, 87, 135	0
2	B	240/279 (86%)	-0.03	3 (1%) 75 71	26, 45, 78, 94	0
All	All	294/334 (88%)	-0.03	4 (1%) 73 70	26, 45, 82, 135	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	126	GLU	2.2
2	B	152	LYS	2.2
1	V	79	C	2.1
2	B	93	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	IMD	B	302	5/5	0.85	0.14	74,74,77,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	V	202	1/1	0.91	0.08	32,32,32,32	0
3	MG	V	201	1/1	0.95	0.06	22,22,22,22	0
3	MG	B	301	1/1	0.96	0.08	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.