



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:41 AM UTC

PDB ID : 8PKW / pdb_00008pkw
Title : Kelch domain of KEAP1 in complex with a ortho-dimethylbenzene linked cyclic peptide 5 (ortho-WRCDPETaEC).
Authors : Braun, M.B.; Bischof, L.; Hartmann, M.D.
Deposited on : 2023-06-27
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	2% 87% 5% 8%
1	B	316	19% 83% 8% 9%
2	P	12	8% 92% 8%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	18	0
			2334	1451	420	444	19			
1	B	287	Total	C	N	O	S	0	4	0
			2124	1319	381	408	16			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	309	GLY	-	expression tag	UNP Q14145
A	310	HIS	-	expression tag	UNP Q14145
A	311	MET	-	expression tag	UNP Q14145
A	319	SER	CYS	conflict	UNP Q14145
A	540	ALA	GLU	conflict	UNP Q14145
A	542	ALA	GLU	conflict	UNP Q14145
A	613	SER	CYS	conflict	UNP Q14145
A	622	SER	CYS	conflict	UNP Q14145
A	624	SER	-	expression tag	UNP Q14145
B	309	GLY	-	expression tag	UNP Q14145
B	310	HIS	-	expression tag	UNP Q14145
B	311	MET	-	expression tag	UNP Q14145
B	319	SER	CYS	conflict	UNP Q14145
B	540	ALA	GLU	conflict	UNP Q14145
B	542	ALA	GLU	conflict	UNP Q14145
B	613	SER	CYS	conflict	UNP Q14145
B	622	SER	CYS	conflict	UNP Q14145
B	624	SER	-	expression tag	UNP Q14145

- Molecule 2 is a protein called CP5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	12	Total	C	N	O	S	0	1	1
			83	46	15	19	3			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Na	0	0
			2	2		
7	B	1	Total	Na	0	0
			1	1		

- Molecule 8 is (2-methylphenyl)methanol (CCD ID: ZK2) (formula: C₈H₁₀O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	P	1	Total	C	0	0
			8	8		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	257	Total	O	0	0
			257	257		
9	B	106	Total	O	0	0
			106	106		
9	P	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 2: CP5



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.68Å 68.97Å 77.97Å 90.00° 117.86° 90.00°	Depositor
Resolution (Å)	42.13 – 1.54 42.13 – 1.54	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.13-1.54) 100.0 (42.13-1.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.54Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.158 , 0.199 0.164 , 0.203	Depositor DCC
R_{free} test set	2255 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4957	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZK2, SO4, EDO, NA, DAL, ACE, CL, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/2396	0.49	0/3270
1	B	0.20	0/2176	0.45	0/2974
2	P	0.25	0/73	0.44	0/95
All	All	0.23	0/4645	0.47	0/6339

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2334	0	2190	12	0
1	B	2124	0	1916	17	0
2	P	83	0	65	1	0
3	A	10	0	0	0	0
3	B	5	0	0	0	0
4	A	1	0	0	0	0
5	A	8	0	12	0	0
6	A	6	0	8	0	0
6	B	6	0	8	0	0
7	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	1	0	0	0	0
8	P	8	0	0	0	0
9	A	257	0	0	0	0
9	B	106	0	0	0	0
9	P	6	0	0	0	0
All	All	4957	0	4199	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:441:GLU:HB3	1:B:452:LEU:HD23	1.60	0.82
1:A:466:ALA:HB1	1:A:514[A]:VAL:HG23	1.70	0.74
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.69	0.74
1:A:447:ARG:HG2	1:A:447:ARG:HH11	1.57	0.68
1:B:454:ALA:H	1:B:495:ASN:HD21	1.47	0.62
1:A:466:ALA:HB1	1:A:514[B]:VAL:HG13	1.82	0.60
1:B:425:ILE:HB	1:B:443:TYR:HB3	1.85	0.57
1:A:557:LEU:HD23	1:A:557:LEU:H	1.70	0.56
1:A:518:CYS:HB3	1:A:536:ARG:HG3	1.91	0.53
1:A:515:LEU:HD22	1:A:566:ILE:HG13	1.91	0.51
1:A:447:ARG:HG2	1:A:447:ARG:NH1	2.25	0.50
1:B:398:PRO:HD2	1:B:610:MET:HE2	1.94	0.49
1:A:554:ARG:HD2	1:A:568[B]:VAL:HG11	1.94	0.49
1:B:365:LEU:H	1:B:365:LEU:HD23	1.80	0.47
1:A:411[B]:VAL:HG11	1:A:437:HIS:CE1	2.51	0.46
1:B:557:LEU:H	1:B:557:LEU:HD23	1.81	0.46
1:B:416:ILE:HD11	1:B:427:ALA:HB1	1.98	0.46
1:A:365:LEU:HD23	1:A:365:LEU:H	1.80	0.45
1:B:472:LEU:HB3	1:B:490:TYR:HB3	1.98	0.45
1:B:369:VAL:HG23	1:B:607:ALA:HB1	2.00	0.43
1:B:415:ARG:NH2	2:P:4:GLU:OE2	2.45	0.42
1:B:562:HIS:HB3	1:B:567:TYR:CE2	2.54	0.41
1:A:604:VAL:HG23	1:A:606[A]:VAL:HG13	2.01	0.41
1:B:327:LEU:CD1	1:B:611:GLU:HA	2.50	0.41
1:A:373:LEU:HB3	1:A:395[A]:CYS:SG	2.60	0.41
1:B:562:HIS:HB3	1:B:567:TYR:HE2	1.86	0.41
1:B:597:MET:HE1	1:B:601:ARG:HG2	2.03	0.41
1:B:327:LEU:HD11	1:B:611:GLU:HA	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:369:VAL:HG21	1:B:609:THR:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/316 (98%)	301 (98%)	7 (2%)	0	100	100
1	B	289/316 (92%)	280 (97%)	9 (3%)	0	100	100
2	P	10/12 (83%)	10 (100%)	0	0	100	100
All	All	607/644 (94%)	591 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/258 (95%)	245 (100%)	0	100	100
1	B	207/258 (80%)	207 (100%)	0	100	100
2	P	9/9 (100%)	9 (100%)	0	100	100
All	All	461/525 (88%)	461 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	337	GLN
1	B	414	ASN
1	B	495	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	701	-	4,4,4	0.18	0	6,6,6	0.25	0
6	GOL	A	705	-	5,5,5	0.88	0	5,5,5	1.19	0
6	GOL	B	703	-	5,5,5	0.76	0	5,5,5	1.13	0
3	SO4	A	701	-	4,4,4	0.24	0	6,6,6	0.18	0
3	SO4	A	702	-	4,4,4	0.23	0	6,6,6	0.07	0
5	EDO	A	706	-	3,3,3	0.45	0	2,2,2	0.34	0
5	EDO	A	704	-	3,3,3	0.42	0	2,2,2	0.43	0
8	ZK2	P	101	2	8,8,9	0.26	0	10,10,11	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	705	-	-	0/4/4/4	-
6	GOL	B	703	-	-	0/4/4/4	-
5	EDO	A	706	-	-	0/1/1/1	-
5	EDO	A	704	-	-	1/1/1/1	-
8	ZK2	P	101	2	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	704	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/316 (92%)	-0.00	7 (2%) 59 69	13, 32, 58, 107	18 (6%)
1	B	287/316 (90%)	1.04	61 (21%) 2 3	23, 57, 91, 115	4 (1%)
2	P	9/12 (75%)	0.78	1 (11%) 10 14	25, 40, 69, 80	1 (11%)
All	All	588/644 (91%)	0.52	69 (11%) 9 12	13, 39, 86, 115	23 (3%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	VAL	4.7
2	P	-1	TRP	4.6
1	B	608	VAL	4.5
1	B	352	TRP	4.4
1	B	515	LEU	4.3
1	B	612	PRO	4.3
1	B	598	THR	4.2
1	B	346	ASN	4.2
1	B	516	HIS	4.1
1	B	327	LEU	3.9
1	B	517	ASN	3.6
1	B	421	ILE	3.6
1	B	353	LEU	3.6
1	B	561	VAL	3.5
1	B	348	SER	3.4
1	B	594	VAL	3.4
1	B	588	THR	3.4
1	A	615	LYS	3.3
1	A	326	ARG	3.3
1	B	518	CYS	3.3
1	A	613	SER	3.3
1	B	399	MET	3.3
1	B	373	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	400	THR	3.2
1	B	610	MET	3.1
1	B	611	GLU	3.0
1	B	452	LEU	2.9
1	B	328	ILE	2.9
1	B	369	VAL	2.7
1	B	403	TRP	2.7
1	B	347	PRO	2.6
1	A	610	MET	2.6
1	A	614	ARG	2.6
1	B	491	TYR	2.6
1	B	468	LEU	2.6
1	B	595	THR	2.5
1	B	375	TYR	2.5
1	B	349	ASP	2.5
1	B	420	VAL	2.5
1	B	351	THR	2.5
1	B	422	ASP	2.5
1	B	593	GLU	2.4
1	B	453	VAL	2.4
1	B	395	CYS	2.4
1	B	371	GLY	2.4
1	B	356	ALA	2.3
1	B	445	PRO	2.3
1	B	586	PRO	2.3
1	B	350	GLY	2.3
1	B	494	ARG	2.3
1	B	402	GLN	2.2
1	B	446	GLU	2.2
1	B	355	LEU	2.2
1	B	563	GLN	2.2
1	B	344	ALA	2.2
1	B	596	ARG	2.2
1	B	374	LEU	2.1
1	B	423	GLY	2.1
1	B	562	HIS	2.1
1	B	370	VAL	2.1
1	B	514	VAL	2.1
1	B	592	SER	2.1
1	B	607	ALA	2.1
1	B	495	ASN	2.1
1	B	567	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	540	ALA	2.0
1	B	425	ILE	2.0
1	B	552	HIS	2.0
1	B	591	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DAL	P	6	5/6	0.97	0.08	37,38,39,43	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	702	5/5	0.85	0.10	132,132,133,134	0
8	ZK2	P	101	8/9	0.89	0.15	63,72,78,78	0
5	EDO	A	706	4/4	0.91	0.14	68,73,73,74	0
6	GOL	B	703	6/6	0.93	0.15	38,40,41,41	6
5	EDO	A	704	4/4	0.94	0.11	41,53,59,64	0
7	NA	A	707	1/1	0.95	0.17	63,63,63,63	0
7	NA	A	708	1/1	0.95	0.14	67,67,67,67	0
6	GOL	A	705	6/6	0.95	0.11	35,54,61,66	0
3	SO4	B	701	5/5	0.96	0.12	39,40,45,49	5
4	CL	A	703	1/1	0.97	0.17	80,80,80,80	0
7	NA	B	702	1/1	0.99	0.06	46,46,46,46	0
3	SO4	A	701	5/5	0.99	0.06	33,34,40,42	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ZK2 P 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.