



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:37 AM UTC

PDB ID : 3PL2 / pdb_00003pl2
Title : Crystal structure of a 5-keto-2-deoxygluconokinase (NCgl0155, Cgl0158) from *Corynebacterium glutamicum* ATCC 13032 KITASATO at 1.89 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-11-12
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	
1	B	319	
1	C	319	
1	D	319	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sugar kinase, ribokinase family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	Se	0	8	0
			2428	1539	409	470	4	6			
1	B	310	Total	C	N	O	S	Se	0	14	0
			2443	1547	408	478	4	6			
1	C	310	Total	C	N	O	S	Se	0	12	0
			2445	1547	413	475	4	6			
1	D	309	Total	C	N	O	S	Se	0	7	0
			2408	1523	407	467	4	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8NTZ3
B	0	GLY	-	expression tag	UNP Q8NTZ3
C	0	GLY	-	expression tag	UNP Q8NTZ3
D	0	GLY	-	expression tag	UNP Q8NTZ3

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	128	Total	O	0	0
			128	128		
3	B	164	Total	O	0	0
			164	164		
3	C	162	Total	O	0	0
			162	162		
3	D	123	Total	O	0	0
			123	123		

- Molecule 1: Sugar kinase, ribokinase family



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	45.91Å 79.49Å 81.70Å 101.86° 95.72° 92.11°	Depositor
Resolution (Å)	37.80 – 1.89 37.80 – 1.89	Depositor EDS
% Data completeness (in resolution range)	97.0 (37.80-1.89) 96.9 (37.80-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.180 , 0.225 (Not available) , 0.244	Depositor DCC
R_{free} test set	4384 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10314	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/2505	0.98	1/3401 (0.0%)
1	B	0.95	0/2533	0.99	0/3438
1	C	0.91	0/2534	0.98	0/3442
1	D	0.93	0/2480	0.99	1/3366 (0.0%)
All	All	0.92	0/10052	0.99	2/13647 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	23	LEU	N-CA-C	-5.13	105.14	111.40
1	A	180	SER	N-CA-C	5.03	112.75	108.22

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	171	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2428	0	2369	10	0
1	B	2443	0	2387	9	0
1	C	2445	0	2386	6	0
1	D	2408	0	2357	12	0
2	B	13	0	5	0	0
3	A	128	0	0	0	0
3	B	164	0	0	0	0
3	C	162	0	0	0	0
3	D	123	0	0	0	0
All	All	10314	0	9504	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:27[B]:VAL:HG13	1:D:32:VAL:HG22	1.70	0.73
1:B:280:LEU:O	1:B:284:LEU:HD13	1.96	0.65
1:D:27[A]:VAL:HG23	1:D:32:VAL:HG22	1.83	0.58
1:C:156:GLU:O	1:C:160:THR:HG23	2.06	0.56
1:A:238:VAL:HG22	1:A:284:LEU:HD11	1.89	0.54
1:C:265:ALA:HB2	1:C:294:VAL:HG21	1.90	0.54
1:A:249:VAL:HG23	1:A:284:LEU:HD23	1.89	0.54
1:C:21[B]:TYR:CE1	1:C:38:TYR:CE2	2.95	0.53
1:B:265:ALA:HB2	1:B:294:VAL:HG21	1.91	0.52
1:A:245:GLU:HG3	1:A:247:VAL:HG23	1.92	0.51
1:A:247:VAL:HG11	1:A:281:GLU:HA	1.94	0.49
1:A:265:ALA:HB2	1:A:294:VAL:HG21	1.93	0.49
1:D:27[B]:VAL:HG13	1:D:32:VAL:CG2	2.41	0.49
1:B:176[B]:MSE:HE3	1:B:176[B]:MSE:H	1.80	0.46
1:C:199:VAL:HG22	1:C:230:ILE:HD12	1.98	0.46
1:B:241[B]:MSE:HG2	1:B:246:THR:HG23	1.97	0.45
1:D:27[B]:VAL:CG1	1:D:32:VAL:HG22	2.41	0.45
1:D:299:GLU:HB2	1:D:303:ALA:HB2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:265:ALA:HB2	1:D:294:VAL:HG21	1.99	0.44
1:D:132:VAL:CG1	1:D:157:ILE:HG23	2.48	0.44
1:A:228:LEU:HD11	1:A:280:LEU:HD11	1.99	0.43
1:D:22:PRO:HB3	1:D:32:VAL:HG11	2.00	0.43
1:D:176[C]:MSE:HB2	1:D:176[C]:MSE:HE2	1.72	0.43
1:D:161:ARG:NH2	1:D:164:ARG:HD3	2.34	0.42
1:C:245:GLU:HG2	1:C:247:VAL:HG23	2.01	0.42
1:B:224[A]:ARG:HG3	1:B:224[A]:ARG:NH1	2.35	0.42
1:B:249:VAL:HG12	1:B:250:PRO:O	2.20	0.42
1:B:224[A]:ARG:HG3	1:B:224[A]:ARG:HH11	1.85	0.41
1:D:24:GLN:O	1:D:27[B]:VAL:HG12	2.20	0.41
1:B:165:ARG:HA	1:B:166:HIS:HA	1.92	0.41
1:B:174[B]:ARG:HE	1:B:176[B]:MSE:SE	2.54	0.41
1:A:52:ARG:HG3	1:A:304:MSE:HB2	2.02	0.40
1:A:165:ARG:HA	1:A:166:HIS:HA	1.96	0.40
1:A:130:ASP:OD1	1:A:133[A]:ARG:NH2	2.54	0.40
1:A:297[A]:ARG:HD2	1:A:303:ALA:HB1	2.04	0.40
1:C:45:ASN:HB3	1:C:304:MSE:HE3	2.03	0.40
1:D:24:GLN:HB2	1:D:27[B]:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/319 (99%)	311 (98%)	5 (2%)	1 (0%)	36	29
1	B	322/319 (101%)	316 (98%)	5 (2%)	1 (0%)	36	29
1	C	321/319 (101%)	317 (99%)	3 (1%)	1 (0%)	36	29
1	D	315/319 (99%)	311 (99%)	3 (1%)	1 (0%)	36	29
All	All	1275/1276 (100%)	1255 (98%)	16 (1%)	4 (0%)	36	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	172	ASP
1	B	172	ASP
1	C	172	ASP
1	D	172	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/255 (101%)	257 (100%)	1 (0%)	84	87
1	B	263/255 (103%)	258 (98%)	5 (2%)	50	47
1	C	262/255 (103%)	258 (98%)	4 (2%)	57	56
1	D	257/255 (101%)	252 (98%)	5 (2%)	50	47
All	All	1040/1020 (102%)	1025 (99%)	15 (1%)	59	59

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LYS
1	B	37	LYS
1	B	176[A]	MSE
1	B	176[B]	MSE
1	B	212	THR
1	B	281	GLU
1	C	23	LEU
1	C	130	ASP
1	C	203	GLU
1	C	227	GLU
1	D	138	LEU
1	D	160	THR
1	D	202	LYS
1	D	233	GLN
1	D	254	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	C	233	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CIT	B	400	-	12,12,12	1.05	0	17,17,17	1.40	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	400	-	-	6/16/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	400	CIT	O4-C5-O3	-3.22	115.05	123.33
2	B	400	CIT	O6-C6-C3	2.73	118.37	113.14

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	400	CIT	C4-C3-C6-O6
2	B	400	CIT	O7-C3-C6-O6
2	B	400	CIT	C4-C3-C6-O5
2	B	400	CIT	C2-C3-C6-O6
2	B	400	CIT	C2-C3-C6-O5
2	B	400	CIT	O7-C3-C6-O5

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/319 (95%)	0.35	10 (3%) 49 52	15, 36, 64, 84	7 (2%)
1	B	306/319 (95%)	0.03	8 (2%) 57 61	16, 29, 49, 61	12 (3%)
1	C	306/319 (95%)	0.18	7 (2%) 61 65	16, 33, 58, 74	11 (3%)
1	D	305/319 (95%)	0.31	11 (3%) 46 49	20, 36, 63, 77	5 (1%)
All	All	1223/1276 (95%)	0.22	36 (2%) 53 57	15, 34, 60, 84	35 (2%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	314	LEU	4.7
1	A	38	TYR	4.4
1	C	253	PHE	4.3
1	C	235	PRO	4.3
1	D	252	PHE	4.3
1	D	38	TYR	3.9
1	B	235	PRO	3.6
1	A	253	PHE	3.5
1	D	234	GLY	3.5
1	A	314	LEU	3.3
1	D	237	GLY	3.2
1	B	253	PHE	3.1
1	D	253	PHE	3.1
1	A	252	PHE	3.0
1	A	312	ALA	2.8
1	A	310	VAL	2.7
1	A	315	ASN	2.6
1	D	21	TYR	2.5
1	A	235	PRO	2.5
1	C	234	GLY	2.5
1	D	6	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	280	LEU	2.3
1	D	254	VAL	2.3
1	B	252	PHE	2.3
1	B	6	SER	2.3
1	D	238	VAL	2.2
1	C	7	THR	2.2
1	C	314	LEU	2.2
1	A	254	VAL	2.2
1	B	297	ARG	2.2
1	B	315	ASN	2.1
1	C	6	SER	2.1
1	B	234	GLY	2.1
1	B	300	CYS	2.1
1	D	251	PRO	2.1
1	C	78[A]	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIT	B	400	13/13	0.91	0.09	32,38,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.