



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 11:50 AM UTC

PDB ID : 3PMA / pdb_00003pma
Title : 2.2 Angstrom crystal structure of the complex between Bovine Thrombin and
Sucrose Octasulfate
Authors : Wright, H.T.; Scarsdale, J.N.; Desai, B.J.
Deposited on : 2010-11-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	29	3% 90% 10%
1	C	29	17% 76% 24%
2	B	259	3% 81% 9% 8%
2	D	259	5% 91% 8%
3	E	2	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	2	 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thrombin light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	29	Total	C	N	O	S	0	0	0
			243	153	38	51	1			
1	C	29	Total	C	N	O	S	0	0	0
			231	146	36	48	1			

- Molecule 2 is a protein called Thrombin heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	0	0
			1883	1207	333	331	12			
2	D	259	Total	C	N	O	S	0	2	0
			2050	1308	366	363	13			

- Molecule 3 is an oligosaccharide called 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	O	S	0	0	0
			55	12	35	8			
3	F	2	Total	C	O	S	0	0	0
			55	12	35	8			

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Na	0	0
			1	1		

- Molecule 5 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			13	6	7		
5	D	1	Total	C	O	0	0
			13	6	7		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	14	Total	O	0	0
			14	14		
6	B	120	Total	O	0	0
			120	120		
6	C	6	Total	O	0	0
			6	6		
6	D	113	Total	O	0	0
			113	113		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Thrombin light chain





- Molecule 1: Thrombin light chain



- Molecule 2: Thrombin heavy chain



- Molecule 2: Thrombin heavy chain



- Molecule 3: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain E:  100%

GU41
YYJ2

- Molecule 3: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain F:  100%

GU41
YYJ2

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.56Å 87.56Å 191.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.80 – 2.20 27.80 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (27.80-2.20) 99.7 (27.80-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.91 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.204 , 0.245 0.218 , 0.255	Depositor DCC
R_{free} test set	1938 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4798	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, YYJ, GU4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/246	0.72	0/328
1	C	0.55	0/234	0.69	0/314
2	B	0.49	0/1930	0.72	1/2613 (0.0%)
2	D	0.46	0/2103	0.73	1/2854 (0.0%)
All	All	0.48	0/4513	0.72	2/6109 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	60(B)	PRO	N-CA-C	6.01	118.03	110.70
2	B	60(B)	PRO	N-CA-C	5.34	117.22	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	243	0	233	2	0
1	C	231	0	212	3	0
2	B	1883	0	1857	12	0
2	D	2050	0	1995	12	0
3	E	55	0	6	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	55	0	6	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	B	13	0	5	0	0
5	D	13	0	5	2	0
6	A	14	0	0	0	0
6	B	120	0	0	1	0
6	C	6	0	0	0	0
6	D	113	0	0	0	0
All	All	4798	0	4319	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:67:ARG:HG2	2:D:70:LYS:HD2	1.71	0.71
2:B:32:MET:HE3	2:B:67:ARG:HB3	1.71	0.70
2:D:60(G):ASN:HD21	5:D:4:CIT:H22	1.66	0.60
2:D:91:HIS:CE1	2:D:101:ARG:HD3	2.38	0.58
3:E:1:GU4:H5	3:E:2:YYJ:O5	2.04	0.58
2:D:17:VAL:HG12	2:D:18:GLU:HG2	1.89	0.55
2:B:243:ASP:O	2:B:244:ARG:HG2	2.08	0.53
2:B:224:LYS:HE2	6:B:300:HOH:O	2.09	0.53
2:D:60(G):ASN:HD21	5:D:4:CIT:C2	2.23	0.50
2:B:123:LEU:O	2:D:246:GLY:HA2	2.12	0.50
2:B:236:LYS:HD2	2:B:237:TRP:N	2.27	0.49
2:B:91:HIS:CE1	2:B:101:ARG:HD3	2.48	0.49
2:B:235:LYS:HE2	2:D:245:LEU:O	2.12	0.49
1:C:14(K):ILE:O	1:C:14(L):GLU:HB3	2.13	0.48
2:B:125:ASP:HB2	3:E:2:YYJ:O3S6	2.13	0.48
2:B:16:ILE:N	2:B:194:ASP:OD2	2.47	0.48
1:A:14(K):ILE:O	1:A:14(L):GLU:HB2	2.16	0.46
1:C:14:ASP:CG	1:C:14(C):GLU:HG3	2.41	0.45
2:B:85:LEU:HD22	2:B:106:LEU:HB3	1.98	0.44
2:B:211:GLY:HA2	2:B:229:THR:O	2.17	0.44
2:D:105:LEU:HD13	2:D:241:VAL:HG22	1.99	0.44
2:D:105:LEU:HD13	2:D:241:VAL:CG2	2.49	0.43
1:A:6:LEU:HD12	2:B:25:GLY:HA3	2.01	0.42
2:D:244:ARG:NH2	3:E:1:GU4:H62	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:36(A):SER:HA	2:D:37:PRO:HA	1.89	0.41
1:C:14(H):GLU:O	1:C:14(K):ILE:HG12	2.20	0.41
2:D:172:THR:HG21	2:D:176:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/29 (93%)	25 (93%)	2 (7%)	0	100	100
1	C	27/29 (93%)	26 (96%)	1 (4%)	0	100	100
2	B	232/259 (90%)	223 (96%)	9 (4%)	0	100	100
2	D	259/259 (100%)	251 (97%)	8 (3%)	0	100	100
All	All	545/576 (95%)	525 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	27/27 (100%)	27 (100%)	0	100	100
1	C	24/27 (89%)	22 (92%)	2 (8%)	10	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	199/226 (88%)	188 (94%)	11 (6%)	19	24
2	D	214/226 (95%)	206 (96%)	8 (4%)	30	41
All	All	464/506 (92%)	443 (96%)	21 (4%)	24	33

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	36	LYS
2	B	36(A)	SER
2	B	46	LEU
2	B	67	ARG
2	B	106	LEU
2	B	126	LYS
2	B	182	CYS
2	B	224	LYS
2	B	234	LEU
2	B	235	LYS
2	B	241	VAL
1	C	1(A)	ASP
1	C	1	CYS
2	D	38	GLN
2	D	46	LEU
2	D	50	ARG
2	D	67	ARG
2	D	106	LEU
2	D	164	GLU
2	D	195	SER
2	D	241	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	143	ASN
2	D	60(G)	ASN
2	D	143	ASN
2	D	179	ASN
2	D	239	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GU4	E	1	3	27,27,28	1.91	4 (14%)	32,43,45	1.54	7 (21%)
3	YYJ	E	2	3	27,28,28	2.07	5 (18%)	33,46,46	1.20	3 (9%)
3	GU4	F	1	3	27,27,28	1.86	6 (22%)	32,43,45	1.63	7 (21%)
3	YYJ	F	2	3	27,28,28	2.01	5 (18%)	33,46,46	1.31	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GU4	E	1	3	-	5/21/38/41	0/1/1/1
3	YYJ	E	2	3	-	9/23/42/42	0/1/1/1
3	GU4	F	1	3	-	8/21/38/41	0/1/1/1
3	YYJ	F	2	3	-	7/23/42/42	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	YYJ	O6-S6	-5.18	1.43	1.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	1	GU4	O6-S6	-5.00	1.43	1.56
3	F	2	YYJ	O6-S6	-4.92	1.43	1.56
3	F	2	YYJ	O1-S1	-4.84	1.43	1.56
3	E	2	YYJ	O1-S1	-4.82	1.44	1.56
3	E	1	GU4	O6-S6	-4.66	1.44	1.56
3	E	2	YYJ	O3-S3	-4.32	1.44	1.57
3	E	1	GU4	O4-S4	-4.16	1.44	1.57
3	E	1	GU4	O3-S3	-4.13	1.44	1.57
3	E	2	YYJ	O2-C2	4.02	1.47	1.40
3	F	2	YYJ	O4-S4	-4.01	1.45	1.57
3	F	1	GU4	O3-S3	-3.87	1.45	1.57
3	F	2	YYJ	O3-S3	-3.81	1.45	1.57
3	E	1	GU4	O2-S2	-3.81	1.45	1.57
3	F	2	YYJ	O2-C2	3.71	1.47	1.40
3	F	1	GU4	O4-S4	-3.71	1.46	1.57
3	E	2	YYJ	O4-S4	-3.58	1.46	1.57
3	F	1	GU4	O2-S2	-3.07	1.48	1.57
3	F	1	GU4	C3-C4	2.08	1.56	1.52
3	F	1	GU4	C1-C2	2.03	1.54	1.51

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	GU4	O2-C2-C3	4.92	112.10	106.65
3	F	2	YYJ	C3-O3-S3	4.40	123.12	117.69
3	E	1	GU4	O2-C2-C3	4.36	111.48	106.65
3	E	2	YYJ	C3-O3-S3	-4.19	112.52	117.69
3	E	1	GU4	O6-C6-C5	3.24	113.34	107.57
3	F	1	GU4	O6-C6-C5	3.23	113.32	107.57
3	F	1	GU4	O10-S2-O2	2.84	112.90	106.37
3	E	1	GU4	O5-C5-C4	2.53	114.61	110.06
3	E	1	GU4	O24-S4-O4	2.43	111.96	106.37
3	E	2	YYJ	C4-O4-S4	2.43	124.87	119.04
3	E	1	GU4	C3-O3-S3	-2.42	113.25	119.04
3	E	1	GU4	O10-S2-O2	2.40	111.89	106.37
3	F	1	GU4	C3-O3-S3	-2.30	113.52	119.04
3	F	2	YYJ	O1S4-S4-O4	2.26	111.57	106.37
3	F	1	GU4	O5-C5-C4	2.25	114.11	110.06
3	E	1	GU4	C1-O5-C5	2.22	115.16	112.19
3	F	1	GU4	O24-S4-O4	2.17	111.36	106.37
3	F	2	YYJ	O1S3-S3-O3	2.12	111.25	106.37
3	F	1	GU4	C1-O5-C5	2.09	114.99	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	YYJ	O1S4-S4-O4	2.02	111.00	106.37

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1	GU4	O5-C5-C6-O6
3	E	1	GU4	C4-O4-S4-O25
3	E	2	YYJ	O1-C1-C2-O5
3	E	2	YYJ	C5-C4-O4-S4
3	E	2	YYJ	C4-C5-C6-O6
3	E	2	YYJ	O5-C5-C6-O6
3	E	2	YYJ	C6-O6-S6-O1S6
3	F	1	GU4	O5-C5-C6-O6
3	F	1	GU4	C4-C5-C6-O6
3	F	2	YYJ	O1-C1-C2-C3
3	F	2	YYJ	O1-C1-C2-O2
3	F	2	YYJ	O1-C1-C2-O5
3	F	2	YYJ	C2-C1-O1-S1
3	F	2	YYJ	C4-C3-O3-S3
3	F	2	YYJ	C4-C5-C6-O6
3	F	2	YYJ	O5-C5-C6-O6
3	E	1	GU4	C4-O4-S4-O26
3	E	2	YYJ	C6-O6-S6-O2S6
3	E	2	YYJ	C6-O6-S6-O3S6
3	E	1	GU4	C4-O4-S4-O24
3	F	1	GU4	C4-O4-S4-O25
3	F	1	GU4	C4-O4-S4-O24
3	F	1	GU4	C4-O4-S4-O26
3	E	2	YYJ	O1-C1-C2-O2
3	E	1	GU4	C4-C5-C6-O6
3	F	1	GU4	C3-O3-S3-O29
3	F	1	GU4	C2-O2-S2-O10
3	F	1	GU4	C2-O2-S2-O12
3	E	2	YYJ	C3-C4-O4-S4

There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1	GU4	2	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	2	YYJ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	CIT	D	4	-	12,12,12	1.05	0	17,17,17	1.76	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	CIT	B	3	-	12,12,12	1.10	0	17,17,17	1.60	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CIT	D	4	-	-	5/16/16/16	-
5	CIT	B	3	-	-	6/16/16/16	-

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	4	CIT	O6-C6-C3	4.23	121.26	113.14
5	B	3	CIT	O6-C6-C3	3.58	120.01	113.14
5	B	3	CIT	O2-C1-C2	2.56	122.45	114.35
5	B	3	CIT	O2-C1-O1	-2.45	117.04	123.33
5	D	4	CIT	O2-C1-C2	2.43	122.04	114.35
5	D	4	CIT	C4-C3-C2	2.41	115.49	109.31
5	B	3	CIT	O4-C5-O3	-2.15	117.79	123.33
5	D	4	CIT	O2-C1-O1	-2.11	117.91	123.33
5	D	4	CIT	O4-C5-O3	-2.08	117.97	123.33
5	B	3	CIT	O6-C6-O5	-2.03	117.35	123.86

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	3	CIT	C2-C3-C6-O5
5	B	3	CIT	C2-C3-C6-O6
5	B	3	CIT	O7-C3-C6-O5
5	B	3	CIT	O7-C3-C6-O6
5	D	4	CIT	C2-C3-C4-C5
5	D	4	CIT	C6-C3-C4-C5
5	D	4	CIT	O7-C3-C4-C5
5	D	4	CIT	C1-C2-C3-O7
5	B	3	CIT	C1-C2-C3-C4
5	B	3	CIT	C1-C2-C3-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	D	4	CIT	C1-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	4	CIT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	29/29 (100%)	0.58	1 (3%) 48 45	26, 42, 63, 68	0
1	C	29/29 (100%)	0.78	5 (17%) 4 3	28, 44, 63, 69	0
2	B	238/259 (91%)	0.18	8 (3%) 48 45	28, 40, 64, 80	0
2	D	259/259 (100%)	0.35	14 (5%) 31 28	28, 44, 69, 85	2 (0%)
All	All	555/576 (96%)	0.31	28 (5%) 34 31	26, 42, 66, 85	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	148	TRP	3.9
2	B	145	ARG	3.4
2	B	243	ASP	3.3
2	D	149(C)	VAL	3.2
1	A	1(C)	GLU	3.1
2	D	149(A)	THR	3.1
2	D	149	THR	3.0
2	B	79	VAL	3.0
2	D	84[A]	MET	3.0
1	C	1(C)	GLU	2.9
2	B	242	ILE	2.7
2	D	247	SER	2.7
2	D	78	LYS	2.6
2	D	147	THR	2.6
2	D	149(D)	ALA	2.5
2	D	112	ILE	2.3
2	B	70	LYS	2.3
1	C	14(K)	ILE	2.3
2	B	236	LYS	2.3
1	C	14(L)	GLU	2.3
2	D	76	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	80	GLU	2.2
1	C	1(B)	ALA	2.1
1	C	1(A)	ASP	2.1
2	D	82	ILE	2.1
2	B	126	LYS	2.1
2	D	65	LEU	2.1
2	D	75	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	YYJ	F	2	28/28	0.27	0.19	93,100,100,100	28
3	GU4	E	1	27/28	0.62	0.21	86,89,100,100	27
3	YYJ	E	2	28/28	0.75	0.17	58,79,88,88	28
3	GU4	F	1	27/28	0.77	0.20	68,82,89,90	22

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CIT	D	4	13/13	0.65	0.15	78,84,87,88	0
5	CIT	B	3	13/13	0.74	0.19	46,63,68,69	0
4	NA	D	2	1/1	0.97	0.04	34,34,34,34	0
4	NA	B	1	1/1	0.99	0.03	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.