



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:10 PM UTC

PDB ID : 3PMK / pdb\_00003pmk  
Title : Crystal structure of the Vesicular Stomatitis Virus RNA free nucleoprotein/p  
hosphoprotein complex  
Authors : Leyrat, C.; Yabukarski, F.; Tarbouriech, N.; Ruigrok, R.W.H.; Jamin, M.  
Deposited on : 2010-11-17  
Resolution : 3.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3685 (3.08-3.00)
Clashscore	190562	4007 (3.08-3.00)
Ramachandran outliers	187476	3834 (3.08-3.00)
Sidechain outliers	187428	3836 (3.08-3.00)
RSRZ outliers	180081	3684 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	
1	B	404	
1	C	404	
1	D	404	
1	E	404	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	N	68	
2	O	68	
2	P	68	
2	Q	68	
2	R	68	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleocapsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3191	2030	534	608	19	0	0	0
1	B	404	3191	2030	534	608	19	0	0	0
1	C	404	3191	2030	534	608	19	0	0	0
1	D	404	3192	2030	534	609	19	0	0	0
1	E	404	3191	2030	534	608	19	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	GLY	-	expression tag	UNP B7UCZ2
A	20	ALA	-	expression tag	UNP B7UCZ2
A	21	MET	-	expression tag	UNP B7UCZ2
B	19	GLY	-	expression tag	UNP B7UCZ2
B	20	ALA	-	expression tag	UNP B7UCZ2
B	21	MET	-	expression tag	UNP B7UCZ2
C	19	GLY	-	expression tag	UNP B7UCZ2
C	20	ALA	-	expression tag	UNP B7UCZ2
C	21	MET	-	expression tag	UNP B7UCZ2
D	19	GLY	-	expression tag	UNP B7UCZ2
D	20	ALA	-	expression tag	UNP B7UCZ2
D	21	MET	-	expression tag	UNP B7UCZ2
E	19	GLY	-	expression tag	UNP B7UCZ2
E	20	ALA	-	expression tag	UNP B7UCZ2
E	21	MET	-	expression tag	UNP B7UCZ2

- Molecule 2 is a protein called Phosphoprotein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	N	30	Total	C	N	O	0	0	0
			246	151	44	51			
2	O	29	Total	C	N	O	0	0	0
			238	146	42	50			
2	P	29	Total	C	N	O	0	0	0
			240	148	43	49			
2	Q	30	Total	C	N	O	0	0	0
			247	152	44	51			
2	R	29	Total	C	N	O	0	0	0
			240	148	43	49			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	61	LEU	-	expression tag	UNP B7UCZ3
N	62	GLU	-	expression tag	UNP B7UCZ3
N	63	HIS	-	expression tag	UNP B7UCZ3
N	64	HIS	-	expression tag	UNP B7UCZ3
N	65	HIS	-	expression tag	UNP B7UCZ3
N	66	HIS	-	expression tag	UNP B7UCZ3
N	67	HIS	-	expression tag	UNP B7UCZ3
N	68	HIS	-	expression tag	UNP B7UCZ3
O	61	LEU	-	expression tag	UNP B7UCZ3
O	62	GLU	-	expression tag	UNP B7UCZ3
O	63	HIS	-	expression tag	UNP B7UCZ3
O	64	HIS	-	expression tag	UNP B7UCZ3
O	65	HIS	-	expression tag	UNP B7UCZ3
O	66	HIS	-	expression tag	UNP B7UCZ3
O	67	HIS	-	expression tag	UNP B7UCZ3
O	68	HIS	-	expression tag	UNP B7UCZ3
P	61	LEU	-	expression tag	UNP B7UCZ3
P	62	GLU	-	expression tag	UNP B7UCZ3
P	63	HIS	-	expression tag	UNP B7UCZ3
P	64	HIS	-	expression tag	UNP B7UCZ3
P	65	HIS	-	expression tag	UNP B7UCZ3
P	66	HIS	-	expression tag	UNP B7UCZ3
P	67	HIS	-	expression tag	UNP B7UCZ3
P	68	HIS	-	expression tag	UNP B7UCZ3
Q	61	LEU	-	expression tag	UNP B7UCZ3
Q	62	GLU	-	expression tag	UNP B7UCZ3
Q	63	HIS	-	expression tag	UNP B7UCZ3
Q	64	HIS	-	expression tag	UNP B7UCZ3
Q	65	HIS	-	expression tag	UNP B7UCZ3
Q	66	HIS	-	expression tag	UNP B7UCZ3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
Q	67	HIS	-	expression tag	UNP B7UCZ3
Q	68	HIS	-	expression tag	UNP B7UCZ3
R	61	LEU	-	expression tag	UNP B7UCZ3
R	62	GLU	-	expression tag	UNP B7UCZ3
R	63	HIS	-	expression tag	UNP B7UCZ3
R	64	HIS	-	expression tag	UNP B7UCZ3
R	65	HIS	-	expression tag	UNP B7UCZ3
R	66	HIS	-	expression tag	UNP B7UCZ3
R	67	HIS	-	expression tag	UNP B7UCZ3
R	68	HIS	-	expression tag	UNP B7UCZ3


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	51	Total O 51 51	0	0
3	B	71	Total O 71 71	0	0
3	C	62	Total O 62 62	0	0
3	D	51	Total O 51 51	0	0
3	E	47	Total O 47 47	0	0
3	N	7	Total O 7 7	0	0
3	O	7	Total O 7 7	0	0
3	P	7	Total O 7 7	0	0
3	Q	4	Total O 4 4	0	0
3	R	4	Total O 4 4	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleocapsid protein




- Molecule 1: Nucleocapsid protein




- Molecule 1: Nucleocapsid protein









- Molecule 2: Phosphoprotein



- Molecule 2: Phosphoprotein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.56Å 171.97Å 239.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.97 – 3.03 59.97 – 3.03	Depositor EDS
% Data completeness (in resolution range)	89.0 (59.97-3.03) 89.0 (59.97-3.03)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.244 , 0.278 0.241 , 0.273	Depositor DCC
$R_{free}$ test set	2742 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtrriage
Anisotropy	0.222	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	17478	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3265	0.80	0/4417
1	B	0.41	0/3265	0.82	3/4417 (0.1%)
1	C	0.41	0/3265	0.82	0/4417
1	D	0.41	0/3266	0.80	3/4417 (0.1%)
1	E	0.42	0/3265	0.82	3/4417 (0.1%)
2	N	0.41	0/247	0.79	0/328
2	O	0.43	0/239	0.75	0/319
2	P	0.38	0/241	0.79	0/320
2	Q	0.43	0/248	0.81	0/330
2	R	0.38	0/241	0.76	0/320
All	All	0.41	0/17542	0.81	9/23702 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	112	ALA	N-CA-C	7.47	120.78	110.24
1	B	149	MET	CA-C-N	5.61	125.27	119.05
1	B	149	MET	C-N-CA	5.61	125.27	119.05
1	D	149	MET	CA-C-N	5.40	125.22	119.28
1	D	149	MET	C-N-CA	5.40	125.22	119.28

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3191	0	3131	47	0
1	B	3191	0	3131	37	0
1	C	3191	0	3131	51	0
1	D	3192	0	3131	51	0
1	E	3191	0	3131	49	0
2	N	246	0	244	1	0
2	O	238	0	233	6	0
2	P	240	0	239	1	0
2	Q	247	0	246	4	0
2	R	240	0	239	2	0
3	A	51	0	0	0	0
3	B	71	0	0	0	0
3	C	62	0	0	0	0
3	D	51	0	0	0	0
3	E	47	0	0	0	0
3	N	7	0	0	0	0
3	O	7	0	0	0	0
3	P	7	0	0	0	0
3	Q	4	0	0	0	0
3	R	4	0	0	0	0
All	All	17478	0	16856	236	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 236 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:23:ASP:HB3	1:E:293:LYS:HE2	1.38	1.04
1:B:117:LEU:H	1:B:118:PRO:HD3	1.22	1.00
1:C:408:ARG:HG3	1:C:408:ARG:HH11	1.33	0.93
1:A:367:ASN:HD22	1:A:367:ASN:H	1.17	0.87
2:Q:16:ARG:HH11	2:Q:16:ARG:HG3	1.40	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/404 (100%)	378 (94%)	15 (4%)	9 (2%)	5	23
1	B	402/404 (100%)	364 (90%)	29 (7%)	9 (2%)	5	23
1	C	402/404 (100%)	373 (93%)	25 (6%)	4 (1%)	12	41
1	D	402/404 (100%)	369 (92%)	23 (6%)	10 (2%)	4	20
1	E	402/404 (100%)	363 (90%)	32 (8%)	7 (2%)	7	29
2	N	28/68 (41%)	27 (96%)	1 (4%)	0	100	100
2	O	27/68 (40%)	24 (89%)	1 (4%)	2 (7%)	1	3
2	P	27/68 (40%)	26 (96%)	1 (4%)	0	100	100
2	Q	28/68 (41%)	26 (93%)	2 (7%)	0	100	100
2	R	27/68 (40%)	24 (89%)	3 (11%)	0	100	100
All	All	2147/2360 (91%)	1974 (92%)	132 (6%)	41 (2%)	6	27

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	LEU
1	B	47	SER
1	B	117	LEU
1	B	176	PRO
1	D	110	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/344 (100%)	329 (96%)	15 (4%)	25	56
1	B	344/344 (100%)	334 (97%)	10 (3%)	37	67
1	C	344/344 (100%)	326 (95%)	18 (5%)	21	51
1	D	344/344 (100%)	333 (97%)	11 (3%)	34	64
1	E	344/344 (100%)	328 (95%)	16 (5%)	23	55
2	N	26/61 (43%)	24 (92%)	2 (8%)	12	37
2	O	25/61 (41%)	25 (100%)	0	100	100
2	P	25/61 (41%)	24 (96%)	1 (4%)	28	59
2	Q	26/61 (43%)	25 (96%)	1 (4%)	29	60
2	R	25/61 (41%)	24 (96%)	1 (4%)	28	59
All	All	1847/2025 (91%)	1772 (96%)	75 (4%)	27	58

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	88	TRP
2	N	18	ASP
1	E	114	ASP
1	E	292	VAL
1	B	361	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	170	GLN
1	E	203	HIS
1	D	208	HIS
1	D	367	ASN
1	E	315	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/404 (100%)	0.35	31 (7%) 19 10	13, 23, 44, 51	0
1	B	404/404 (100%)	0.38	30 (7%) 20 10	15, 25, 47, 57	0
1	C	404/404 (100%)	0.27	28 (6%) 23 11	12, 21, 44, 49	0
1	D	404/404 (100%)	0.34	28 (6%) 23 11	13, 26, 50, 58	0
1	E	404/404 (100%)	0.44	31 (7%) 19 10	14, 29, 53, 59	0
2	N	30/68 (44%)	0.76	3 (10%) 12 6	23, 35, 57, 72	0
2	O	29/68 (42%)	1.25	6 (20%) 2 2	22, 38, 72, 87	0
2	P	29/68 (42%)	0.81	5 (17%) 4 2	22, 34, 57, 61	0
2	Q	30/68 (44%)	1.17	7 (23%) 2 1	22, 37, 86, 98	0
2	R	29/68 (42%)	0.98	6 (20%) 2 2	21, 35, 65, 78	0
All	All	2167/2360 (91%)	0.40	175 (8%) 18 9	12, 25, 50, 98	0

The worst 5 of 175 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	360	SER	6.1
1	C	358	ASP	5.7
1	E	43	ASN	5.4
1	C	365	THR	4.8
1	E	113	LEU	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.