



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 03:35 AM UTC

PDB ID : 6PMA / pdb_00006pma
Title : TRK-A IN COMPLEX WITH LIGAND
Authors : Subramanian, G.; Brown, D.G.
Deposited on : 2019-07-01
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2381 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called High affinity nerve growth factor receptor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	290	2323	1482	417	407	1	16	77	0	0

- Molecule 2 is N-[2-chloro-5-(trifluoromethyl)phenyl]-2-[1-(4-methoxyphenyl)-4-oxo-1,4-dihydro-5H-pyrazolo[3,4-d]pyrimidin-5-yl]acetamide (CCD ID: OQS) (formula: C₂₁H₁₅ClF₃N₅O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	F	N	O		
2	A	1	33	21	1	3	5	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: High affinity nerve growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	52.09Å 52.09Å 227.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.16 – 2.53 45.16 – 2.53	Depositor EDS
% Data completeness (in resolution range)	99.2 (45.16-2.53) 99.2 (45.16-2.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.204 , 0.266 0.209 , 0.271	Depositor DCC
R_{free} test set	906 reflections (7.41%)	wwPDB-VP
Wilson B-factor (Å ²)	52.0	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.075 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2381	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OQS, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.30	19/2369 (0.8%)	1.94	32/3204 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	761	ARG	NE-CZ	71.67	2.11	1.33
1	A	508	ARG	NE-CZ	-37.47	0.91	1.33
1	A	780	ARG	CD-NE	-26.80	1.08	1.46
1	A	607	ASP	CB-CG	-23.00	0.94	1.52
1	A	521	PHE	CG-CD2	-17.46	1.02	1.38
1	A	523	LYS	CE-NZ	-16.30	1.00	1.49
1	A	686	ARG	CB-CG	15.75	1.99	1.52
1	A	570	GLN	CG-CD	-14.29	1.16	1.52
1	A	521	PHE	CG-CD1	13.75	1.67	1.38
1	A	767	GLU	CG-CD	11.86	1.81	1.52
1	A	547	LYS	CG-CD	-9.64	1.23	1.52
1	A	559	ARG	CB-CG	-8.93	1.25	1.52
1	A	747	GLU	CG-CD	-7.55	1.33	1.52
1	A	585	LEU	CG-CD1	-6.20	1.32	1.52
1	A	585	LEU	CG-CD2	5.68	1.71	1.52
1	A	583	ARG	NE-CZ	-5.39	1.27	1.33
1	A	650	ASP	C-O	5.29	1.28	1.23
1	A	555	GLN	CB-CG	5.14	1.67	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	554	ARG	CB-CG	-5.12	1.37	1.52

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	583	ARG	NE-CZ-NH2	26.44	143.00	119.20
1	A	761	ARG	CD-NE-CZ	-24.45	90.17	124.40
1	A	583	ARG	NE-CZ-NH1	-24.23	97.27	121.50
1	A	686	ARG	CA-CB-CG	-18.60	76.90	114.10
1	A	607	ASP	CA-CB-CG	15.42	128.02	112.60
1	A	583	ARG	CD-NE-CZ	15.28	145.80	124.40
1	A	548	GLU	CG-CD-OE2	15.20	153.35	118.40
1	A	548	GLU	CG-CD-OE1	-15.11	83.64	118.40
1	A	570	GLN	CB-CG-CD	14.76	137.69	112.60
1	A	761	ARG	NE-CZ-NH1	-14.28	107.22	121.50
1	A	747	GLU	CB-CG-CD	12.71	134.21	112.60
1	A	780	ARG	CD-NE-CZ	12.34	141.68	124.40
1	A	521	PHE	CB-CG-CD1	-11.28	101.53	120.70
1	A	570	GLN	CG-CD-NE2	10.45	132.07	116.40
1	A	523	LYS	CD-CE-NZ	9.97	143.79	111.90
1	A	686	ARG	CB-CG-CD	-9.52	89.40	111.30
1	A	535	GLU	CA-CB-CG	-8.50	97.10	114.10
1	A	554	ARG	CA-CB-CG	8.01	130.11	114.10
1	A	570	GLN	CG-CD-OE1	-7.75	105.30	120.80
1	A	508	ARG	NE-CZ-NH1	-7.61	113.89	121.50
1	A	767	GLU	CB-CG-CD	-7.61	99.67	112.60
1	A	521	PHE	CG-CD1-CE1	-7.43	108.07	120.70
1	A	558	GLN	CG-CD-NE2	6.18	125.68	116.40
1	A	744	ARG	CB-CA-C	6.09	119.63	109.89
1	A	558	GLN	CG-CD-OE1	-5.66	109.48	120.80
1	A	628	LEU	CA-C-N	5.31	127.34	120.44
1	A	628	LEU	C-N-CA	5.31	127.34	120.44
1	A	560	GLU	CB-CG-CD	5.19	121.42	112.60
1	A	678	THR	CA-CB-OG1	-5.19	101.82	109.60
1	A	556	ASP	CA-C-N	5.13	127.11	120.44
1	A	556	ASP	C-N-CA	5.13	127.11	120.44
1	A	532	LEU	N-CA-C	-5.10	105.37	112.45

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	508	ARG	Sidechain
1	A	521	PHE	Sidechain
1	A	583	ARG	Sidechain
1	A	761	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2323	0	2304	21	0
2	A	33	0	0	0	0
3	A	25	0	0	0	0
All	All	2381	0	2304	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:769:GLN:HA	1:A:769:GLN:OE1	1.83	0.78
1:A:560:GLU:HB3	1:A:587:MET:HE1	1.90	0.53
1:A:617:VAL:HG22	1:A:662:LEU:HD11	1.90	0.53
1:A:680:TYR:HB2	1:A:688:MET:HE2	1.92	0.52
1:A:506:LYS:HD3	1:A:508:ARG:HE	1.75	0.51
1:A:757:TYR:CE2	1:A:761:ARG:HD2	2.46	0.49
1:A:568:GLN:O	1:A:569:HIS:HB2	2.12	0.49
1:A:683:VAL:HG13	1:A:687:THR:HB	1.95	0.49
1:A:533:LEU:O	1:A:536:GLN:O	2.33	0.47
1:A:581:GLU:O	1:A:581:GLU:HG3	2.14	0.47
1:A:618:ALA:HB1	1:A:619:PRO:HD2	1.97	0.47
1:A:617:VAL:HG12	1:A:618:ALA:O	2.16	0.46
1:A:695:PRO:HB3	1:A:711:TRP:CD2	2.51	0.46
1:A:581:GLU:O	1:A:581:GLU:CG	2.66	0.44
1:A:648:HIS:O	1:A:649:ARG:HB2	2.18	0.43
1:A:710:VAL:O	1:A:763:CYS:HB3	2.19	0.43
1:A:752:CYS:SG	1:A:760:MET:HE3	2.59	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:683:VAL:O	1:A:685:GLY:N	2.53	0.41
1:A:728:TRP:HB3	1:A:731:LEU:HD12	2.01	0.41
1:A:565:THR:HA	1:A:568:GLN:NE2	2.35	0.41
1:A:593:ARG:NH1	1:A:660:GLN:HG3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/311 (91%)	267 (94%)	12 (4%)	4 (1%)	9	16

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	643	GLY
1	A	684	GLY
1	A	547	LYS
1	A	685	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/260 (93%)	230 (95%)	12 (5%)	22	42

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	508	ARG
1	A	536	GLN
1	A	558	GLN
1	A	581	GLU
1	A	596	ASP
1	A	599	ARG
1	A	739	CYS
1	A	755	GLU
1	A	761	ARG
1	A	773	SER
1	A	780	ARG
1	A	795	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	677	1	8,9,10	0.72	0	7,12,14	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	677	1	-	3/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	677	SEP	CB-OG-P-O3P
1	A	677	SEP	CB-OG-P-O1P
1	A	677	SEP	CB-OG-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OQS	A	801	-	36,36,36	0.69	1 (2%)	53,53,53	1.08	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OQS	A	801	-	-	0/20/20/20	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	OQS	C7-C19	-2.19	1.39	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	OQS	C6-N2-N1	4.31	108.03	104.59
2	A	801	OQS	C7-C6-N2	-2.07	109.88	112.45
2	A	801	OQS	C19-N1-N2	-2.06	108.58	111.12


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/311 (92%)	0.12	11 (3%) 44 40	23, 42, 73, 93	24 (8%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	500	ALA	3.9
1	A	520	ALA	3.4
1	A	612	ALA	3.3
1	A	501	CYS	3.2
1	A	521	PHE	3.0
1	A	537	ASP	2.6
1	A	519	GLY	2.5
1	A	684	GLY	2.3
1	A	582	GLY	2.1
1	A	607	ASP	2.1
1	A	613	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	677	10/11	0.84	0.12	37,46,73,80	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OQS	A	801	33/33	0.94	0.07	31,33,47,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.