



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:17 AM UTC

PDB ID : 1PNO / pdb_00001pno
Title : Crystal structure of R. rubrum transhydrogenase domain III bound to NADP
Authors : Sundaresan, V.; Yamaguchi, M.; Chartron, J.; Stout, C.D.
Deposited on : 2003-06-12
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	17% 82% 16% .
1	B	180	16% 83% 14% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3018 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD(P) transhydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	0	0
			1359	857	229	261	12			
1	B	177	Total	C	N	O	S	0	0	0
			1339	846	224	257	12			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	285	SER	-	cloning artifact	UNP Q59765
A	286	GLY	-	cloning artifact	UNP Q59765
A	287	HIS	-	cloning artifact	UNP Q59765
A	288	ILE	-	cloning artifact	UNP Q59765
A	289	GLU	-	cloning artifact	UNP Q59765
A	290	GLY	-	cloning artifact	UNP Q59765
A	291	ARG	-	cloning artifact	UNP Q59765
A	292	HIS	-	cloning artifact	UNP Q59765
A	293	MET	-	cloning artifact	UNP Q59765
B	285	SER	-	cloning artifact	UNP Q59765
B	286	GLY	-	cloning artifact	UNP Q59765
B	287	HIS	-	cloning artifact	UNP Q59765
B	288	ILE	-	cloning artifact	UNP Q59765
B	289	GLU	-	cloning artifact	UNP Q59765
B	290	GLY	-	cloning artifact	UNP Q59765
B	291	ARG	-	cloning artifact	UNP Q59765
B	292	HIS	-	cloning artifact	UNP Q59765
B	293	MET	-	cloning artifact	UNP Q59765

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total	O	0	0
			120	120		
3	B	104	Total	O	0	0
			104	104		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD(P) transhydrogenase subunit beta



- Molecule 1: NAD(P) transhydrogenase subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	117.95Å 117.95Å 211.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.50 – 2.10 28.50 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.7 (28.50-2.10) 97.8 (28.50-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.209 , 0.230 0.210 , 0.232	Depositor DCC
R_{free} test set	1496 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3018	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1384	0.86	2/1869 (0.1%)
1	B	0.39	0/1363	0.87	2/1841 (0.1%)
All	All	0.39	0/2747	0.87	4/3710 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	341	VAL	N-CA-C	6.02	117.14	108.48
1	A	395	THR	N-CA-C	-5.31	106.74	114.12
1	B	292	HIS	N-CA-C	5.07	116.90	109.24
1	A	396	ASN	N-CA-C	5.02	116.67	109.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1359	0	1340	18	0
1	B	1339	0	1325	20	0
2	A	48	0	25	0	0
2	B	48	0	25	0	0
3	A	120	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	104	0	0	1	0
All	All	3018	0	2715	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:289:GLU:HG3	1:B:292:HIS:NE2	1.95	0.79
1:B:324:GLN:HE21	1:B:324:GLN:H	1.38	0.71
1:B:346:HIS:CE1	1:B:348:VAL:HG22	2.27	0.69
1:A:442:ARG:HG3	1:A:442:ARG:HH11	1.57	0.69
1:A:289:GLU:HG3	1:A:292:HIS:NE2	2.12	0.65
1:A:459:ILE:HG22	1:A:463:MET:HE3	1.79	0.64
1:B:293:MET:HE3	1:B:446:MET:HE2	1.81	0.62
1:B:324:GLN:H	1:B:324:GLN:NE2	1.98	0.61
1:B:373:LEU:C	1:B:373:LEU:HD23	2.26	0.59
1:A:442:ARG:HG3	1:A:442:ARG:NH1	2.20	0.56
1:A:459:ILE:CG2	1:A:463:MET:HE3	2.35	0.56
1:A:414:ASP:HB3	1:A:417:LYS:HD3	1.87	0.56
1:A:373:LEU:HD23	1:A:373:LEU:C	2.33	0.53
1:A:346:HIS:CE1	1:A:348:VAL:HG22	2.44	0.52
1:B:453:LYS:O	1:B:457:GLU:HG3	2.08	0.52
1:B:403:PRO:HA	1:B:408:TYR:CD2	2.46	0.51
1:B:376:ILE:HD13	1:B:379:SER:HB3	1.93	0.50
1:A:403:PRO:HA	1:A:408:TYR:CG	2.47	0.49
1:B:374:GLU:HB2	3:B:77:HOH:O	2.12	0.49
1:A:416:GLU:HG2	1:A:439:LEU:HD11	1.95	0.48
1:A:437:ASN:OD1	1:A:439:LEU:HB2	2.13	0.48
1:A:456:THR:O	1:A:460:VAL:HG23	2.15	0.47
1:B:299:ASP:O	1:B:303:ILE:HG13	2.15	0.47
1:B:425:LYS:O	1:B:449:PHE:HA	2.16	0.46
1:A:422:LEU:HD12	1:A:446:MET:HB3	1.96	0.46
1:B:289:GLU:HG3	1:B:292:HIS:CE1	2.52	0.45
1:B:376:ILE:HD13	1:B:376:ILE:O	2.17	0.45
1:B:289:GLU:CG	1:B:292:HIS:NE2	2.73	0.45
1:B:398:ALA:C	1:B:400:LYS:H	2.24	0.45
1:A:388:VAL:CG1	1:A:391:ALA:HB3	2.48	0.43
1:B:325:HIS:O	1:B:329:GLU:HG3	2.19	0.42
1:B:289:GLU:CD	1:B:292:HIS:HE2	2.28	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:LYS:O	1:A:449:PHE:HA	2.20	0.41
1:A:402:ASP:O	1:A:408:TYR:HB2	2.20	0.41
1:A:298:GLU:OE1	1:A:298:GLU:N	2.54	0.40
1:B:398:ALA:C	1:B:400:LYS:N	2.79	0.40
1:A:404:SER:O	1:B:353:PRO:HB3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/180 (99%)	172 (97%)	6 (3%)	0	100	100
1	B	175/180 (97%)	164 (94%)	11 (6%)	0	100	100
All	All	353/360 (98%)	336 (95%)	17 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/142 (100%)	134 (94%)	8 (6%)	19	18
1	B	140/142 (99%)	135 (96%)	5 (4%)	31	34
All	All	282/284 (99%)	269 (95%)	13 (5%)	24	25

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	400	LYS
1	A	412	ILE
1	A	416	GLU
1	A	422	LEU
1	A	439	LEU
1	A	442	ARG
1	A	454	LYS
1	A	464	ASN
1	B	324	GLN
1	B	334	LEU
1	B	375	GLU
1	B	376	ILE
1	B	436	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	458	GLN
1	A	461	GLN
1	B	324	GLN
1	B	458	GLN
1	B	461	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	B	500	-	50,52,52	2.68	17 (34%)	71,80,80	2.57	32 (45%)
2	NAP	A	500	-	50,52,52	2.68	17 (34%)	71,80,80	2.50	31 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	500	-	-	14/35/67/67	0/5/5/5
2	NAP	A	500	-	-	11/35/67/67	0/5/5/5

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAP	PN-O3	9.56	1.69	1.59
2	B	500	NAP	PN-O3	9.54	1.69	1.59
2	A	500	NAP	C5A-C4A	6.83	1.51	1.39
2	B	500	NAP	C5A-C4A	6.69	1.51	1.39
2	B	500	NAP	C2N-N1N	6.51	1.42	1.35
2	A	500	NAP	C2N-N1N	6.29	1.41	1.35
2	A	500	NAP	O4D-C1D	5.65	1.48	1.40
2	B	500	NAP	O4D-C1D	5.51	1.48	1.40
2	A	500	NAP	C6N-N1N	4.38	1.45	1.35
2	B	500	NAP	C6N-N1N	4.36	1.45	1.35
2	A	500	NAP	C1B-N9A	4.11	1.57	1.46
2	B	500	NAP	C1B-N9A	3.80	1.56	1.46
2	B	500	NAP	C3B-C2B	-3.57	1.45	1.53
2	A	500	NAP	C3B-C2B	-3.52	1.45	1.53
2	A	500	NAP	O4B-C1B	3.26	1.49	1.42
2	B	500	NAP	C2N-C3N	3.19	1.44	1.39
2	A	500	NAP	C4N-C3N	3.14	1.44	1.39
2	B	500	NAP	C4N-C3N	3.09	1.44	1.39
2	B	500	NAP	O4B-C1B	3.09	1.49	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAP	C2N-C3N	2.81	1.43	1.39
2	B	500	NAP	P2B-O3X	-2.66	1.44	1.54
2	A	500	NAP	P2B-O3X	-2.63	1.45	1.54
2	B	500	NAP	PA-O3	2.55	1.62	1.59
2	B	500	NAP	C2A-N1A	2.37	1.38	1.33
2	B	500	NAP	O2B-C2B	-2.37	1.36	1.44
2	A	500	NAP	O2B-C2B	-2.36	1.36	1.44
2	A	500	NAP	C2A-N1A	2.33	1.38	1.33
2	B	500	NAP	C3B-C4B	-2.30	1.47	1.53
2	B	500	NAP	C2B-C1B	-2.30	1.47	1.53
2	A	500	NAP	C2B-C1B	-2.29	1.47	1.53
2	A	500	NAP	C3B-C4B	-2.25	1.47	1.53
2	B	500	NAP	C8A-N9A	2.24	1.41	1.37
2	A	500	NAP	C8A-N9A	2.23	1.41	1.37
2	A	500	NAP	O3B-C3B	2.05	1.48	1.43

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	NAP	C4D-O4D-C1D	-7.86	102.73	109.92
2	A	500	NAP	C4D-O4D-C1D	-7.44	103.11	109.92
2	A	500	NAP	P2B-O2B-C2B	7.07	142.32	123.43
2	B	500	NAP	P2B-O2B-C2B	6.99	142.10	123.43
2	B	500	NAP	C6N-N1N-C2N	-6.91	116.00	121.88
2	A	500	NAP	C6N-N1N-C2N	-6.70	116.18	121.88
2	B	500	NAP	O4D-C4D-C3D	4.04	113.18	105.15
2	A	500	NAP	C6A-C5A-N7A	4.02	139.84	132.09
2	B	500	NAP	O2B-C2B-C3B	3.91	125.72	111.68
2	B	500	NAP	C6A-C5A-N7A	3.90	139.61	132.09
2	B	500	NAP	C2B-C3B-C4B	3.89	110.36	101.99
2	A	500	NAP	O4D-C4D-C3D	3.86	112.81	105.15
2	A	500	NAP	O5D-C5D-C4D	3.84	122.08	108.99
2	B	500	NAP	C4A-N9A-C8A	3.69	109.61	105.74
2	B	500	NAP	O5B-C5B-C4B	3.66	121.45	108.99
2	B	500	NAP	C6A-C5A-C4A	-3.66	112.19	117.18
2	A	500	NAP	O2B-C2B-C3B	3.65	124.77	111.68
2	A	500	NAP	C2B-C3B-C4B	3.65	109.84	101.99
2	A	500	NAP	C6A-C5A-C4A	-3.64	112.21	117.18
2	A	500	NAP	C4A-N9A-C8A	3.64	109.56	105.74
2	B	500	NAP	O5D-C5D-C4D	3.63	121.36	108.99
2	B	500	NAP	N3A-C2A-N1A	-3.63	123.09	128.58
2	B	500	NAP	C2D-C3D-C4D	-3.57	95.72	102.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	NAP	O5B-C5B-C4B	3.49	120.87	108.99
2	A	500	NAP	C3N-C7N-N7N	-3.38	113.58	117.74
2	A	500	NAP	N3A-C2A-N1A	-3.34	123.52	128.58
2	A	500	NAP	C5A-N7A-C8A	3.33	108.68	103.45
2	A	500	NAP	C2D-C3D-C4D	-3.31	96.22	102.61
2	A	500	NAP	C4B-O4B-C1B	-3.24	102.31	109.47
2	B	500	NAP	C4B-O4B-C1B	-3.19	102.43	109.47
2	B	500	NAP	C5A-N7A-C8A	3.16	108.42	103.45
2	B	500	NAP	C3N-C7N-N7N	-3.00	114.04	117.74
2	B	500	NAP	O4B-C4B-C5B	2.91	118.65	109.33
2	A	500	NAP	O4B-C4B-C5B	2.91	118.64	109.33
2	B	500	NAP	C2B-C1B-N9A	2.86	118.46	113.75
2	B	500	NAP	C5N-C6N-N1N	2.84	124.25	120.38
2	B	500	NAP	O2N-PN-O3	-2.75	99.84	107.27
2	A	500	NAP	C5N-C6N-N1N	2.58	123.91	120.38
2	A	500	NAP	O2N-PN-O3	-2.52	100.45	107.27
2	A	500	NAP	N9A-C8A-N7A	-2.48	110.42	113.94
2	B	500	NAP	N9A-C8A-N7A	-2.47	110.43	113.94
2	B	500	NAP	O4D-C4D-C5D	2.47	117.25	109.33
2	A	500	NAP	PN-O5D-C5D	2.44	135.34	121.35
2	A	500	NAP	O4D-C4D-C5D	2.42	117.07	109.33
2	B	500	NAP	PN-O5D-C5D	2.38	135.00	121.35
2	A	500	NAP	O5D-PN-O1N	2.37	118.35	108.94
2	A	500	NAP	O2D-C2D-C3D	2.36	119.39	111.82
2	A	500	NAP	C2B-C1B-N9A	2.34	117.61	113.75
2	B	500	NAP	C2A-N1A-C6A	2.31	122.52	118.73
2	A	500	NAP	C4A-C5A-N7A	-2.30	107.95	110.58
2	B	500	NAP	O2D-C2D-C3D	2.28	119.13	111.82
2	A	500	NAP	O3D-C3D-C4D	2.27	117.61	111.08
2	B	500	NAP	C5A-C4A-N9A	-2.27	103.33	105.81
2	B	500	NAP	O5D-PN-O1N	2.24	117.83	108.94
2	A	500	NAP	C2N-C3N-C4N	2.23	120.85	118.26
2	A	500	NAP	C5A-C4A-N9A	-2.21	103.40	105.81
2	B	500	NAP	PA-O5B-C5B	2.20	133.96	121.35
2	B	500	NAP	O3D-C3D-C4D	2.20	117.39	111.08
2	B	500	NAP	C2N-C3N-C4N	2.11	120.71	118.26
2	B	500	NAP	O4B-C1B-N9A	-2.11	104.04	108.09
2	A	500	NAP	C2A-N1A-C6A	2.08	122.15	118.73
2	B	500	NAP	C4A-C5A-N7A	-2.08	108.20	110.58
2	A	500	NAP	N3A-C4A-N9A	2.03	130.61	127.17

There are no chirality outliers.


All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	500	NAP	O4D-C1D-N1N-C2N
2	A	500	NAP	O4D-C1D-N1N-C6N
2	A	500	NAP	C2D-C1D-N1N-C2N
2	A	500	NAP	C2D-C1D-N1N-C6N
2	B	500	NAP	C5B-O5B-PA-O1A
2	B	500	NAP	O4D-C1D-N1N-C2N
2	B	500	NAP	O4D-C1D-N1N-C6N
2	B	500	NAP	C2D-C1D-N1N-C2N
2	B	500	NAP	C2D-C1D-N1N-C6N
2	A	500	NAP	C1B-C2B-O2B-P2B
2	B	500	NAP	C1B-C2B-O2B-P2B
2	B	500	NAP	O4B-C4B-C5B-O5B
2	B	500	NAP	C3B-C4B-C5B-O5B
2	A	500	NAP	C3B-C2B-O2B-P2B
2	B	500	NAP	C3B-C2B-O2B-P2B
2	B	500	NAP	O4D-C4D-C5D-O5D
2	A	500	NAP	O4B-C4B-C5B-O5B
2	A	500	NAP	O4D-C4D-C5D-O5D
2	A	500	NAP	C3B-C4B-C5B-O5B
2	B	500	NAP	C5B-O5B-PA-O2A
2	B	500	NAP	C5B-O5B-PA-O3
2	B	500	NAP	PN-O3-PA-O1A
2	A	500	NAP	C2B-C1B-N9A-C8A
2	A	500	NAP	PN-O3-PA-O2A
2	B	500	NAP	PN-O3-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/180 (100%)	0.95	31 (17%) 4 4	31, 50, 79, 87	0
1	B	177/180 (98%)	0.76	29 (16%) 4 4	28, 47, 81, 97	0
All	All	357/360 (99%)	0.86	60 (16%) 4 4	28, 49, 80, 97	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	288	ILE	9.7
1	A	285	SER	4.9
1	A	287	HIS	4.8
1	A	374	GLU	4.7
1	B	289	GLU	4.7
1	A	288	ILE	4.5
1	A	286	GLY	4.4
1	B	401	THR	4.3
1	B	405	SER	4.2
1	B	375	GLU	4.2
1	B	403	PRO	3.9
1	A	368	ASP	3.9
1	A	442	ARG	3.6
1	B	290	GLY	3.5
1	A	401	THR	3.5
1	A	464	ASN	3.3
1	A	289	GLU	3.3
1	B	464	ASN	3.2
1	B	404	SER	3.1
1	B	462	ALA	3.0
1	B	406	PRO	3.0
1	A	409	GLY	2.9
1	A	290	GLY	2.9
1	A	375	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	408	TYR	2.9
1	B	402	ASP	2.8
1	B	417	LYS	2.8
1	A	461	GLN	2.8
1	B	418	ALA	2.7
1	B	408	TYR	2.7
1	B	461	GLN	2.6
1	A	381	GLN	2.6
1	A	398	ALA	2.6
1	A	462	ALA	2.6
1	A	402	ASP	2.6
1	A	438	GLU	2.5
1	B	378	SER	2.5
1	A	416	GLU	2.5
1	A	337	GLU	2.4
1	B	407	ILE	2.4
1	A	308	SER	2.4
1	A	397	PRO	2.4
1	B	438	GLU	2.4
1	A	412	ILE	2.3
1	B	443	ASN	2.3
1	A	443	ASN	2.2
1	B	381	GLN	2.2
1	A	291	ARG	2.2
1	B	414	ASP	2.2
1	B	374	GLU	2.2
1	B	441	PHE	2.1
1	B	416	GLU	2.1
1	B	436	GLU	2.1
1	A	407	ILE	2.1
1	A	454	LYS	2.1
1	B	400	LYS	2.1
1	B	298	GLU	2.0
1	A	384	ASP	2.0
1	A	414	ASP	2.0
1	B	291	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAP	A	500	48/48	0.96	0.09	32,39,43,46	0
2	NAP	B	500	48/48	0.97	0.08	34,43,46,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.