



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:18 AM UTC

PDB ID : 4PNS / pdb_00004pns
Title : Crystal Structure of human Tankyrase 2 in complex with INH2BP.
Authors : Qiu, W.; Lam, R.; Romanov, V.; Gordon, R.; Gebremeskel, S.; Vodsedalek, J.; Thompson, C.; Beletskaya, I.; Battaile, K.P.; Pai, E.F.; Chirgadze, N.Y.
Deposited on : 2014-05-25
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	4% 82% 13%
1	B	227	6% 84% 12%
1	C	227	7% 80% 11% 8%
1	D	227	10% 76% 7% 16%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	2	0
			1591	1000	298	282	11			
1	B	200	Total	C	N	O	S	0	1	0
			1625	1027	299	288	11			
1	C	208	Total	C	N	O	S	0	0	0
			1673	1055	308	299	11			
1	D	190	Total	C	N	O	S	0	0	0
			1538	970	286	271	11			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	938	MET	-	initiating methionine	UNP Q9H2K2
A	939	GLY	-	expression tag	UNP Q9H2K2
A	940	SER	-	expression tag	UNP Q9H2K2
A	941	SER	-	expression tag	UNP Q9H2K2
A	942	HIS	-	expression tag	UNP Q9H2K2
A	943	HIS	-	expression tag	UNP Q9H2K2
A	944	HIS	-	expression tag	UNP Q9H2K2
A	945	HIS	-	expression tag	UNP Q9H2K2
A	946	HIS	-	expression tag	UNP Q9H2K2
A	947	HIS	-	expression tag	UNP Q9H2K2
A	948	SER	-	expression tag	UNP Q9H2K2
A	949	SER	-	expression tag	UNP Q9H2K2
A	950	GLY	-	expression tag	UNP Q9H2K2
A	951	ARG	-	expression tag	UNP Q9H2K2
A	952	GLU	-	expression tag	UNP Q9H2K2
A	953	ASN	-	expression tag	UNP Q9H2K2
A	954	LEU	-	expression tag	UNP Q9H2K2
A	955	TYR	-	expression tag	UNP Q9H2K2
A	956	PHE	-	expression tag	UNP Q9H2K2
A	957	GLN	-	expression tag	UNP Q9H2K2
A	958	GLY	-	expression tag	UNP Q9H2K2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	938	MET	-	initiating methionine	UNP Q9H2K2
B	939	GLY	-	expression tag	UNP Q9H2K2
B	940	SER	-	expression tag	UNP Q9H2K2
B	941	SER	-	expression tag	UNP Q9H2K2
B	942	HIS	-	expression tag	UNP Q9H2K2
B	943	HIS	-	expression tag	UNP Q9H2K2
B	944	HIS	-	expression tag	UNP Q9H2K2
B	945	HIS	-	expression tag	UNP Q9H2K2
B	946	HIS	-	expression tag	UNP Q9H2K2
B	947	HIS	-	expression tag	UNP Q9H2K2
B	948	SER	-	expression tag	UNP Q9H2K2
B	949	SER	-	expression tag	UNP Q9H2K2
B	950	GLY	-	expression tag	UNP Q9H2K2
B	951	ARG	-	expression tag	UNP Q9H2K2
B	952	GLU	-	expression tag	UNP Q9H2K2
B	953	ASN	-	expression tag	UNP Q9H2K2
B	954	LEU	-	expression tag	UNP Q9H2K2
B	955	TYR	-	expression tag	UNP Q9H2K2
B	956	PHE	-	expression tag	UNP Q9H2K2
B	957	GLN	-	expression tag	UNP Q9H2K2
B	958	GLY	-	expression tag	UNP Q9H2K2
C	938	MET	-	initiating methionine	UNP Q9H2K2
C	939	GLY	-	expression tag	UNP Q9H2K2
C	940	SER	-	expression tag	UNP Q9H2K2
C	941	SER	-	expression tag	UNP Q9H2K2
C	942	HIS	-	expression tag	UNP Q9H2K2
C	943	HIS	-	expression tag	UNP Q9H2K2
C	944	HIS	-	expression tag	UNP Q9H2K2
C	945	HIS	-	expression tag	UNP Q9H2K2
C	946	HIS	-	expression tag	UNP Q9H2K2
C	947	HIS	-	expression tag	UNP Q9H2K2
C	948	SER	-	expression tag	UNP Q9H2K2
C	949	SER	-	expression tag	UNP Q9H2K2
C	950	GLY	-	expression tag	UNP Q9H2K2
C	951	ARG	-	expression tag	UNP Q9H2K2
C	952	GLU	-	expression tag	UNP Q9H2K2
C	953	ASN	-	expression tag	UNP Q9H2K2
C	954	LEU	-	expression tag	UNP Q9H2K2
C	955	TYR	-	expression tag	UNP Q9H2K2
C	956	PHE	-	expression tag	UNP Q9H2K2
C	957	GLN	-	expression tag	UNP Q9H2K2
C	958	GLY	-	expression tag	UNP Q9H2K2

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	938	MET	-	initiating methionine	UNP Q9H2K2
D	939	GLY	-	expression tag	UNP Q9H2K2
D	940	SER	-	expression tag	UNP Q9H2K2
D	941	SER	-	expression tag	UNP Q9H2K2
D	942	HIS	-	expression tag	UNP Q9H2K2
D	943	HIS	-	expression tag	UNP Q9H2K2
D	944	HIS	-	expression tag	UNP Q9H2K2
D	945	HIS	-	expression tag	UNP Q9H2K2
D	946	HIS	-	expression tag	UNP Q9H2K2
D	947	HIS	-	expression tag	UNP Q9H2K2
D	948	SER	-	expression tag	UNP Q9H2K2
D	949	SER	-	expression tag	UNP Q9H2K2
D	950	GLY	-	expression tag	UNP Q9H2K2
D	951	ARG	-	expression tag	UNP Q9H2K2
D	952	GLU	-	expression tag	UNP Q9H2K2
D	953	ASN	-	expression tag	UNP Q9H2K2
D	954	LEU	-	expression tag	UNP Q9H2K2
D	955	TYR	-	expression tag	UNP Q9H2K2
D	956	PHE	-	expression tag	UNP Q9H2K2
D	957	GLN	-	expression tag	UNP Q9H2K2
D	958	GLY	-	expression tag	UNP Q9H2K2

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is 6-amino-5-iodo-2H-chromen-2-one (CCD ID: 32Y) (formula: C₉H₆INO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	I	N	O	0	0
			13	9	1	1	2		
3	B	1	Total	C	I	N	O	0	0
			13	9	1	1	2		
3	C	1	Total	C	I	N	O	0	0
			13	9	1	1	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	241	Total	O	0	0
			241	241		
5	B	242	Total	O	0	0
			242	242		
5	C	250	Total	O	0	0
			250	250		
5	D	203	Total	O	0	0
			203	203		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tankyrase-2




• Molecule 1: Tankyrase-2




• Molecule 1: Tankyrase-2



• Molecule 1: Tankyrase-2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.92Å 79.50Å 153.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.25 – 1.65 44.25 – 1.65	Depositor EDS
% Data completeness (in resolution range)	92.2 (44.25-1.65) 92.2 (44.25-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.65Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.10.0, BUSTER 2.10.0	Depositor
R, R_{free}	0.188 , 0.209 0.189 , 0.202	Depositor DCC
R_{free} test set	1020 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 54.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7418	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 32Y, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/1638 (0.1%)	1.10	2/2200 (0.1%)
1	B	0.81	1/1670 (0.1%)	1.06	3/2239 (0.1%)
1	C	0.79	0/1718	1.15	11/2310 (0.5%)
1	D	0.80	0/1580	1.09	2/2122 (0.1%)
All	All	0.80	2/6606 (0.0%)	1.10	18/8871 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1115	MET	SD-CE	8.62	2.01	1.79
1	A	994	ILE	CG1-CD1	-5.35	1.30	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1052	GLY	N-CA-C	7.41	121.62	112.73
1	D	969	GLU	CB-CG-CD	6.75	124.07	112.60
1	B	969	GLU	CB-CG-CD	6.43	123.53	112.60
1	C	1116	ALA	N-CA-C	-6.40	105.61	113.41
1	A	1116	ALA	N-CA-C	-6.33	105.69	113.41
1	C	960	PRO	N-CA-CB	6.24	109.80	103.25
1	C	969	GLU	CB-CG-CD	5.95	122.71	112.60
1	C	1051	ILE	N-CA-C	5.79	120.23	111.89
1	C	1083	VAL	N-CA-C	5.46	116.12	110.82
1	A	969	GLU	CB-CG-CD	5.44	121.84	112.60
1	B	990	ASN	CA-CB-CG	5.40	118.00	112.60
1	B	1116	ALA	N-CA-C	-5.37	106.40	113.12
1	C	1036	VAL	CA-C-N	5.21	128.03	120.79
1	C	1036	VAL	C-N-CA	5.21	128.03	120.79
1	C	1150	GLU	N-CA-C	5.17	117.00	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1116	ALA	N-CA-C	-5.13	106.71	113.12
1	C	959	SER	CA-C-N	5.04	126.14	119.84
1	C	959	SER	C-N-CA	5.04	126.14	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1591	0	1526	7	0
1	B	1625	0	1553	3	0
1	C	1673	0	1595	18	0
1	D	1538	0	1469	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	13	0	6	0	0
3	B	13	0	6	0	0
3	C	13	0	6	0	0
4	C	12	0	16	1	0
5	A	241	0	0	4	0
5	B	242	0	0	0	0
5	C	250	0	0	4	0
5	D	203	0	0	3	0
All	All	7418	0	6177	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1023:HIS:HD2	1:C:1025:ASN:H	1.13	0.91

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1105:LYS:H	1:C:1123:HIS:HD2	1.21	0.87
1:C:1023:HIS:CD2	1:C:1025:ASN:H	2.01	0.75
1:A:1143[B]:ARG:NH2	5:A:1400:HOH:O	2.27	0.68
1:C:1023:HIS:HD2	1:C:1025:ASN:N	1.93	0.60
1:A:969:GLU:HG3	1:A:994:ILE:HD12	1.84	0.59
1:C:972:MET:HG2	1:C:994:ILE:HD11	1.86	0.58
1:C:1083:VAL:HG13	1:C:1092:CYS:SG	2.43	0.58
1:B:1057:ALA:O	1:D:1117:HIS:HE1	1.87	0.58
1:A:1057:ALA:O	1:C:1117:HIS:HE1	1.87	0.57
1:C:1054:MET:HB2	4:C:1203:GOL:H11	1.87	0.57
1:C:1083:VAL:CG1	1:C:1092:CYS:SG	2.93	0.57
1:C:1082:PRO:HD2	5:C:1519:HOH:O	2.04	0.56
1:C:1083:VAL:HG12	5:C:1519:HOH:O	2.05	0.56
1:D:1117:HIS:HD2	5:D:1350:HOH:O	1.89	0.55
1:D:972:MET:HG2	1:D:994:ILE:HD11	1.89	0.54
1:A:1110:PHE:HD2	1:A:1130:SER:HB3	1.74	0.53
1:C:1023:HIS:HE1	1:D:1022:ASN:O	1.92	0.53
1:B:972:MET:HG2	1:B:994:ILE:HD11	1.90	0.52
1:C:1011:HIS:HE1	5:C:1526:HOH:O	1.93	0.51
1:D:1054:MET:HE3	1:D:1055:PHE:CE2	2.45	0.51
1:A:1143[B]:ARG:CZ	5:A:1400:HOH:O	2.58	0.49
1:C:1036:VAL:HG11	1:C:1152:LEU:CD1	2.42	0.49
1:C:1036:VAL:HG11	1:C:1152:LEU:HD13	1.96	0.48
1:C:1117:HIS:HD2	5:C:1347:HOH:O	1.96	0.47
1:D:1011:HIS:HE1	5:D:1358:HOH:O	2.01	0.44
1:B:1115:MET:SD	1:B:1119:PRO:HG3	2.57	0.44
1:D:1115:MET:HE1	1:D:1119:PRO:HG3	1.99	0.43
1:A:1082:PRO:HD2	5:A:1476:HOH:O	2.18	0.43
1:A:1011:HIS:HE1	5:A:1409:HOH:O	2.01	0.42
1:C:1019:GLU:OE1	1:C:1123:HIS:HE1	2.02	0.42
1:D:975:THR:HB	1:D:1064:ASN:HA	2.01	0.42
1:C:1036:VAL:CG1	1:C:1152:LEU:HD13	2.50	0.41
1:D:970:GLU:OE1	1:D:974:SER:OG	2.38	0.40
1:D:984:HIS:HD2	5:D:1304:HOH:O	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/227 (86%)	192 (98%)	3 (2%)	0	100	100
1	B	193/227 (85%)	191 (99%)	2 (1%)	0	100	100
1	C	206/227 (91%)	204 (99%)	2 (1%)	0	100	100
1	D	186/227 (82%)	181 (97%)	5 (3%)	0	100	100
All	All	780/908 (86%)	768 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/192 (86%)	165 (99%)	1 (1%)	78	68
1	B	169/192 (88%)	166 (98%)	3 (2%)	51	30
1	C	174/192 (91%)	169 (97%)	5 (3%)	37	14
1	D	160/192 (83%)	157 (98%)	3 (2%)	50	28
All	All	669/768 (87%)	657 (98%)	12 (2%)	51	30

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1132	ASN
1	B	962	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	990	ASN
1	B	1128	ARG
1	C	957	GLN
1	C	1037	ASN
1	C	1042	LYS
1	C	1083	VAL
1	C	1100	ARG
1	D	1051	ILE
1	D	1054	MET
1	D	1113	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	990	ASN
1	A	998	GLN
1	A	1011	HIS
1	A	1023	HIS
1	A	1095	GLN
1	B	984	HIS
1	B	990	ASN
1	B	1011	HIS
1	B	1041	HIS
1	C	984	HIS
1	C	990	ASN
1	C	1011	HIS
1	C	1023	HIS
1	C	1041	HIS
1	C	1064	ASN
1	C	1109	GLN
1	C	1117	HIS
1	C	1123	HIS
1	D	990	ASN
1	D	998	GLN
1	D	1011	HIS
1	D	1023	HIS
1	D	1095	GLN
1	D	1117	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GOL	C	1204	-	5,5,5	0.12	0	5,5,5	0.22	0
3	32Y	C	1202	-	13,14,14	2.22	6 (46%)	18,20,20	1.47	3 (16%)
3	32Y	A	1202	-	13,14,14	2.09	5 (38%)	18,20,20	1.25	2 (11%)
3	32Y	B	1202	-	13,14,14	1.76	4 (30%)	18,20,20	1.42	3 (16%)
4	GOL	C	1203	-	5,5,5	0.23	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	1204	-	-	0/4/4/4	-
3	32Y	A	1202	-	-	-	0/2/2/2
3	32Y	C	1202	-	-	-	0/2/2/2
3	32Y	B	1202	-	-	-	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	1203	-	-	2/4/4/4	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1202	32Y	O06-C07	4.16	1.45	1.38
3	A	1202	32Y	O06-C07	3.93	1.44	1.38
3	C	1202	32Y	C03-C02	3.81	1.48	1.40
3	B	1202	32Y	C12-I13	-3.79	2.02	2.09
3	A	1202	32Y	O06-C05	3.18	1.43	1.38
3	A	1202	32Y	C12-I13	-2.96	2.04	2.09
3	A	1202	32Y	C03-C02	2.84	1.46	1.40
3	C	1202	32Y	C04-C05	2.73	1.45	1.39
3	B	1202	32Y	C03-C02	2.56	1.46	1.40
3	C	1202	32Y	O06-C05	2.52	1.42	1.38
3	B	1202	32Y	C04-C05	2.50	1.44	1.39
3	A	1202	32Y	C04-C05	2.43	1.44	1.39
3	B	1202	32Y	O06-C07	2.38	1.42	1.38
3	C	1202	32Y	C09-C10	2.30	1.38	1.34
3	C	1202	32Y	C11-C05	2.20	1.45	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1202	32Y	O06-C05-C11	3.08	122.98	120.09
3	A	1202	32Y	C05-O06-C07	-2.99	117.95	121.67
3	C	1202	32Y	O06-C05-C11	2.88	122.79	120.09
3	B	1202	32Y	O06-C07-O08	-2.63	112.99	116.40
3	C	1202	32Y	O06-C07-O08	-2.58	113.05	116.40
3	A	1202	32Y	O06-C05-C11	2.52	122.45	120.09
3	C	1202	32Y	C05-O06-C07	-2.46	118.61	121.67
3	B	1202	32Y	C11-C10-C09	-2.13	118.67	121.05

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	C	1203	GOL	O1-C1-C2-C3
4	C	1203	GOL	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1203	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/227 (86%)	0.17	10 (5%) 33 37	11, 25, 50, 67	2 (1%)
1	B	200/227 (88%)	0.22	14 (7%) 22 25	14, 24, 47, 67	1 (0%)
1	C	208/227 (91%)	0.14	17 (8%) 17 19	14, 23, 45, 62	0
1	D	190/227 (83%)	0.59	23 (12%) 8 9	17, 30, 55, 71	0
All	All	795/908 (87%)	0.28	64 (8%) 18 20	11, 25, 50, 71	3 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	958	GLY	6.4
1	A	1052	GLY	6.1
1	D	1136	LEU	5.5
1	A	1049	ALA	5.2
1	B	1050	TYR	5.1
1	B	1052	GLY	4.8
1	B	1049	ALA	4.7
1	B	954	LEU	4.7
1	D	1129	PRO	4.7
1	D	1113	MET	4.6
1	D	1051	ILE	4.5
1	C	965	PHE	4.2
1	A	965	PHE	4.2
1	D	1054	MET	4.1
1	A	1131	VAL	4.0
1	C	955	TYR	3.8
1	C	1052	GLY	3.7
1	D	1050	TYR	3.6
1	D	965	PHE	3.4
1	D	1110	PHE	3.4
1	B	1083	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	974	SER	3.3
1	A	962	ASP	3.2
1	C	1051	ILE	3.1
1	C	953	ASN	3.1
1	D	1083	VAL	3.0
1	D	1112	ALA	2.8
1	C	960	PRO	2.8
1	D	1055	PHE	2.8
1	D	1071	TYR	2.7
1	B	1133	GLY	2.7
1	D	1114	LYS	2.7
1	B	965	PHE	2.6
1	A	1132	ASN	2.6
1	B	962	ASP	2.6
1	C	1050	TYR	2.6
1	D	1053	GLY	2.5
1	B	963	LYS	2.5
1	A	1110	PHE	2.4
1	B	1053	GLY	2.4
1	C	1053	GLY	2.4
1	D	1049	ALA	2.4
1	A	966	GLN	2.4
1	D	1075	ILE	2.4
1	A	1008	ARG	2.4
1	C	954	LEU	2.4
1	C	962	ASP	2.4
1	C	1083	VAL	2.4
1	A	1133	GLY	2.3
1	B	1115	MET	2.3
1	C	963	LYS	2.3
1	D	1001	CYS	2.3
1	C	959	SER	2.3
1	C	958	GLY	2.3
1	D	1115	MET	2.3
1	B	1160	PRO	2.2
1	C	1021	HIS	2.2
1	C	957	GLN	2.2
1	D	1052	GLY	2.2
1	D	979	HIS	2.2
1	C	1085	LYS	2.1
1	D	982	GLY	2.1
1	D	983	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	955	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	32Y	B	1202	13/13	0.73	0.17	28,37,59,90	0
3	32Y	A	1202	13/13	0.74	0.18	32,35,57,86	0
4	GOL	C	1203	6/6	0.75	0.18	39,43,49,55	0
3	32Y	C	1202	13/13	0.78	0.14	20,23,44,79	0
4	GOL	C	1204	6/6	0.91	0.10	21,28,34,39	0
2	ZN	D	1200	1/1	0.99	0.04	40,40,40,40	0
2	ZN	B	1201	1/1	0.99	0.02	32,32,32,32	0
2	ZN	A	1201	1/1	1.00	0.02	31,31,31,31	0
2	ZN	C	1201	1/1	1.00	0.04	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 32Y B 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around 32Y A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 32Y C 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.