



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 12:11 AM UTC

PDB ID : 7PN9 / pdb_00007pn9
Title : Evolved unspecific peroxygenase with A77L mutation in complex with lauric acid
Authors : Fernandez-Garcia, A.; Sanz-Aparicio, J.
Deposited on : 2021-09-06
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	32% 95%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	DAO	A	408	-	-	-	X

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aromatic peroxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C	N	O	S	0	6	0
			2566	1624	447	486	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ALA	VAL	engineered mutation	UNP B9W4V6
A	67	PHE	LEU	engineered mutation	UNP B9W4V6
A	75	ILE	VAL	engineered mutation	UNP B9W4V6
A	77	LEU	ALA	engineered mutation	UNP B9W4V6
A	248	VAL	ILE	engineered mutation	UNP B9W4V6
A	311	LEU	PHE	engineered mutation	UNP B9W4V6

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $\text{C}_{34}\text{H}_{32}\text{FeN}_4\text{O}_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0


- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg		
			1	1	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl		
			1	1	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O		
			14	8	1	5	0	0
5	A	1	Total	C	N	O		
			14	8	1	5	0	0
5	A	1	Total	C	N	O		
			14	8	1	5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is LAURIC ACID (CCD ID: DAO) (formula: $C_{12}H_{24}O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			14	12	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	234	Total	O	0	0
			234	234		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aromatic peroxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.09Å 57.78Å 60.86Å 90.00° 109.50° 90.00°	Depositor
Resolution (Å)	48.16 – 1.85 48.16 – 1.85	Depositor EDS
% Data completeness (in resolution range)	91.7 (48.16-1.85) 91.7 (48.16-1.85)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.26 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.247 , 0.277 0.255 , 0.285	Depositor DCC
R_{free} test set	1386 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	9.1	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	2926	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAO, PO4, GOL, MG, NAG, CL, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	0/2636	1.26	1/3586 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	131	ASP	CA-CB-CG	6.10	118.70	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2566	0	2445	11	0
2	A	43	0	30	1	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	56	0	52	0	0
6	A	5	0	0	0	0
7	A	6	0	8	0	0
8	A	14	0	23	4	0
9	A	234	0	0	2	0
All	All	2926	0	2558	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:THR:HA	8:A:408:DAO:H112	1.51	0.93
1:A:195:GLY:HA3	8:A:408:DAO:H92	1.78	0.65
1:A:195:GLY:HA3	8:A:408:DAO:C9	2.37	0.54
1:A:48:LEU:HD21	1:A:59:ILE:HA	1.96	0.48
1:A:61:ASN:O	1:A:65[B]:GLU:HG3	2.17	0.45
1:A:319:CYS:HA	9:A:687:HOH:O	2.17	0.44
2:A:401:HEM:HBC2	2:A:401:HEM:CMC	2.48	0.43
1:A:160:TYR:HB3	1:A:214:LEU:HB3	2.01	0.43
1:A:192:THR:CA	8:A:408:DAO:H112	2.36	0.43
1:A:176:SER:HB3	1:A:183:PHE:CD1	2.54	0.42
1:A:90:THR:OG1	1:A:102:THR:O	2.36	0.41
1:A:65[B]:GLU:CD	9:A:650:HOH:O	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/328 (100%)	320 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/273 (101%)	276 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	254	GLN
1	A	295	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	401	9,1,3	50,50,50	1.60	8 (16%)	67,82,82	1.54	11 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	410	1	14,14,15	0.77	1 (7%)	17,19,21	1.64	4 (23%)
8	DAO	A	408	-	13,13,13	0.65	0	13,13,13	0.66	0
5	NAG	A	405	1	14,14,15	0.77	0	17,19,21	1.51	4 (23%)
7	GOL	A	407	-	5,5,5	0.05	0	5,5,5	0.24	0
5	NAG	A	404	1	14,14,15	0.60	1 (7%)	17,19,21	0.90	1 (5%)
5	NAG	A	409	1	14,14,15	0.72	0	17,19,21	1.03	0
6	PO4	A	406	-	4,4,4	0.92	0	6,6,6	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	401	9,1,3	-	4/14/54/54	-
5	NAG	A	410	1	-	2/6/23/26	0/1/1/1
8	DAO	A	408	-	-	9/11/11/11	-
5	NAG	A	405	1	-	4/6/23/26	0/1/1/1
7	GOL	A	407	-	-	0/4/4/4	-
5	NAG	A	404	1	-	1/6/23/26	0/1/1/1
5	NAG	A	409	1	-	2/6/23/26	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	HEM	FE-NB	4.49	2.08	1.94
2	A	401	HEM	C1B-NB	-4.07	1.33	1.40
2	A	401	HEM	C4D-ND	-3.53	1.34	1.40
2	A	401	HEM	FE-NC	3.41	2.06	1.95
2	A	401	HEM	C1D-ND	-3.25	1.32	1.38
5	A	410	NAG	C1-C2	2.64	1.55	1.52
2	A	401	HEM	C4B-NB	-2.64	1.33	1.38
2	A	401	HEM	C4C-NC	-2.19	1.35	1.39
2	A	401	HEM	O2D-CGD	-2.06	1.24	1.30
5	A	404	NAG	C1-C2	2.02	1.55	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	CHC-C4B-NB	4.07	128.80	124.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	410	NAG	O5-C1-C2	3.29	116.38	111.29
2	A	401	HEM	CHD-C4C-NC	3.24	127.99	124.45
2	A	401	HEM	C1B-NB-C4B	3.20	109.00	105.21
5	A	405	NAG	O3-C3-C2	2.97	115.57	109.40
5	A	410	NAG	C2-N2-C7	-2.88	119.04	122.90
2	A	401	HEM	CMB-C2B-C1B	2.76	129.34	125.03
2	A	401	HEM	CHA-C4D-ND	2.65	127.65	124.37
5	A	410	NAG	C8-C7-N2	2.57	120.37	116.12
5	A	410	NAG	O7-C7-C8	-2.55	117.51	122.05
2	A	401	HEM	CHA-C4D-C3D	-2.50	120.61	125.23
5	A	405	NAG	O5-C5-C6	2.47	112.47	107.66
5	A	405	NAG	C6-C5-C4	-2.46	106.97	113.02
5	A	405	NAG	C1-C2-N2	2.46	114.31	110.43
2	A	401	HEM	CAA-CBA-CGA	-2.43	107.22	113.67
2	A	401	HEM	CHB-C1B-NB	2.40	127.34	124.37
2	A	401	HEM	CMD-C2D-C1D	2.28	128.59	125.03
2	A	401	HEM	O2A-CGA-CBA	2.18	120.87	114.00
5	A	404	NAG	C1-C2-N2	-2.10	107.12	110.43
2	A	401	HEM	C4C-NC-C1C	2.06	109.17	105.82

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	410	NAG	C8-C7-N2-C2
5	A	410	NAG	O7-C7-N2-C2
8	A	408	DAO	C7-C8-C9-C10
5	A	405	NAG	O5-C5-C6-O6
5	A	405	NAG	C4-C5-C6-O6
8	A	408	DAO	C2-C3-C4-C5
8	A	408	DAO	C5-C6-C7-C8
8	A	408	DAO	C9-C10-C11-C12
8	A	408	DAO	C1-C2-C3-C4
8	A	408	DAO	C4-C5-C6-C7
5	A	405	NAG	C1-C2-N2-C7
8	A	408	DAO	C11-C10-C9-C8
5	A	404	NAG	C8-C7-N2-C2
2	A	401	HEM	C2B-C3B-CAB-CBB
5	A	409	NAG	C4-C5-C6-O6
5	A	405	NAG	C3-C2-N2-C7
2	A	401	HEM	CAA-CBA-CGA-O2A
2	A	401	HEM	C2A-CAA-CBA-CGA

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	HEM	CAA-CBA-CGA-O1A
5	A	409	NAG	O5-C5-C6-O6
8	A	408	DAO	C3-C4-C5-C6
8	A	408	DAO	O2-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	HEM	1	0
8	A	408	DAO	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	325/328 (99%)	1.77	104 (32%) 1 1	6, 15, 27, 49	6 (1%)

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	317	ALA	6.5
1	A	327	ARG	5.9
1	A	295	ASN	5.7
1	A	298	VAL	5.6
1	A	318	GLY	5.2
1	A	278	CYS	4.7
1	A	316	ALA	4.6
1	A	107	PRO	4.6
1	A	119	GLY	4.5
1	A	109	PRO	4.4
1	A	299	GLN	4.2
1	A	319	CYS	4.2
1	A	324	PRO	4.1
1	A	4	LEU	4.0
1	A	191	PHE	4.0
1	A	315	VAL	4.0
1	A	322	VAL	3.9
1	A	193	ALA	3.9
1	A	323	PHE	3.8
1	A	297	THR	3.8
1	A	10	GLU	3.8
1	A	75	ILE	3.7
1	A	72	GLN	3.6
1	A	79	TYR	3.6
1	A	111	SER	3.6
1	A	279	LEU	3.6
1	A	320	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	230	ASP	3.5
1	A	326	GLY	3.5
1	A	280[A]	MET	3.5
1	A	302	LYS	3.3
1	A	296	PRO	3.2
1	A	20	GLU	3.2
1	A	287	ILE	3.2
1	A	3	GLY	3.1
1	A	16	LEU	3.1
1	A	209	ARG	3.0
1	A	245	GLU	3.0
1	A	194	TYR	3.0
1	A	306	THR	2.9
1	A	9	LEU	2.9
1	A	174[A]	GLN	2.9
1	A	104	PRO	2.9
1	A	105	ASP	2.9
1	A	28	ARG	2.9
1	A	156	GLY	2.9
1	A	259	VAL	2.8
1	A	76	PHE	2.8
1	A	300	LEU	2.8
1	A	303	ALA	2.8
1	A	48	LEU	2.8
1	A	188	PHE	2.7
1	A	312	PHE	2.7
1	A	19	ASP	2.7
1	A	261	LYS	2.7
1	A	217	ASP	2.7
1	A	29	PRO	2.7
1	A	260	GLY	2.7
1	A	204	PHE	2.7
1	A	110	ALA	2.6
1	A	146	GLU	2.6
1	A	262	ILE	2.6
1	A	301	ARG	2.5
1	A	121	PHE	2.5
1	A	101	LEU	2.5
1	A	290	LYS	2.5
1	A	92	LEU	2.5
1	A	13	SER	2.5
1	A	74	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	69	PHE	2.5
1	A	150	ASP	2.5
1	A	289	VAL	2.5
1	A	294	PRO	2.5
1	A	148	LEU	2.4
1	A	292	LEU	2.4
1	A	277	PRO	2.4
1	A	313	GLN	2.4
1	A	291	SER	2.4
1	A	249	GLN	2.4
1	A	30	GLY	2.4
1	A	227	ARG	2.3
1	A	106	PRO	2.3
1	A	7	GLY	2.3
1	A	155	PHE	2.3
1	A	286	ASN	2.3
1	A	108	PRO	2.3
1	A	175	ASP	2.3
1	A	127	MET	2.3
1	A	304	LEU	2.3
1	A	281	TYR	2.2
1	A	112	VAL	2.2
1	A	229	PRO	2.2
1	A	177	ILE	2.2
1	A	159	LYS	2.2
1	A	267	VAL	2.1
1	A	293	TYR	2.1
1	A	197	THR	2.1
1	A	84	VAL	2.1
1	A	26	PRO	2.1
1	A	314	GLY	2.1
1	A	73	ALA	2.1
1	A	89	ILE	2.0
1	A	274	PHE	2.0
1	A	285	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	404	14/15	0.44	0.30	42,47,58,61	4
8	DAO	A	408	14/14	0.53	0.46	41,51,53,54	0
5	NAG	A	410	14/15	0.58	0.24	32,34,43,46	0
7	GOL	A	407	6/6	0.75	0.20	24,31,32,35	0
5	NAG	A	405	14/15	0.77	0.16	17,19,21,23	0
6	PO4	A	406	5/5	0.80	0.16	30,33,35,36	0
5	NAG	A	409	14/15	0.86	0.11	12,13,15,17	0
2	HEM	A	401	43/43	0.92	0.12	10,10,12,15	0
3	MG	A	402	1/1	0.95	0.05	10,10,10,10	0
4	CL	A	403	1/1	0.96	0.10	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around DAO A 408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around HEM A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.