



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:55 PM UTC

PDB ID : 8PNM / pdb_00008pnm
Title : Structure of human KCTD15 BTB domain mutant G88D crystal form 2
Authors : Cruz Walma, D.A.; Cros, J.; Bradshaw, W.; Richardson, W.; Chen, Z.; Chalk, R.; Wilkie, A.; Bullock, A.N.
Deposited on : 2023-06-30
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	6% 78% 10% 10%
1	B	116	2% 81% 16%
1	C	116	3% 77% 5% 18%
1	D	116	5% 84% 5% 11%
1	E	116	6% 79% 7% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	116	
1	G	116	
1	H	116	
1	I	116	
1	J	116	
1	K	116	
1	L	116	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9618 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BTB/POZ domain-containing protein KCTD15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	95	Total	C	N	O	S	0	0	0
			776	501	125	148	2			
1	A	104	Total	C	N	O	S	0	0	0
			814	527	132	153	2			
1	G	105	Total	C	N	O	S	0	0	0
			816	528	134	152	2			
1	E	100	Total	C	N	O	S	0	0	0
			781	507	125	147	2			
1	J	108	Total	C	N	O	S	0	0	0
			811	522	135	153	1			
1	D	103	Total	C	N	O	S	0	0	0
			797	516	132	147	2			
1	B	98	Total	C	N	O	S	0	0	0
			775	502	124	147	2			
1	I	102	Total	C	N	O	S	0	0	0
			785	510	128	146	1			
1	H	102	Total	C	N	O	S	0	0	0
			779	505	127	145	2			
1	K	95	Total	C	N	O	S	0	0	0
			735	473	121	139	2			
1	F	95	Total	C	N	O	S	0	0	0
			743	480	121	140	2			
1	L	107	Total	C	N	O		0	0	0
			760	491	122	147				

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	SER	-	expression tag	UNP Q96SI1
C	2	MET	-	expression tag	UNP Q96SI1
C	39	ASP	GLY	engineered mutation	UNP Q96SI1
A	1	SER	-	expression tag	UNP Q96SI1
A	2	MET	-	expression tag	UNP Q96SI1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	GLY	engineered mutation	UNP Q96SI1
G	1	SER	-	expression tag	UNP Q96SI1
G	2	MET	-	expression tag	UNP Q96SI1
G	39	ASP	GLY	engineered mutation	UNP Q96SI1
E	1	SER	-	expression tag	UNP Q96SI1
E	2	MET	-	expression tag	UNP Q96SI1
E	39	ASP	GLY	engineered mutation	UNP Q96SI1
J	1	SER	-	expression tag	UNP Q96SI1
J	2	MET	-	expression tag	UNP Q96SI1
J	39	ASP	GLY	engineered mutation	UNP Q96SI1
D	1	SER	-	expression tag	UNP Q96SI1
D	2	MET	-	expression tag	UNP Q96SI1
D	39	ASP	GLY	engineered mutation	UNP Q96SI1
B	1	SER	-	expression tag	UNP Q96SI1
B	2	MET	-	expression tag	UNP Q96SI1
B	39	ASP	GLY	engineered mutation	UNP Q96SI1
I	1	SER	-	expression tag	UNP Q96SI1
I	2	MET	-	expression tag	UNP Q96SI1
I	39	ASP	GLY	engineered mutation	UNP Q96SI1
H	1	SER	-	expression tag	UNP Q96SI1
H	2	MET	-	expression tag	UNP Q96SI1
H	39	ASP	GLY	engineered mutation	UNP Q96SI1
K	1	SER	-	expression tag	UNP Q96SI1
K	2	MET	-	expression tag	UNP Q96SI1
K	39	ASP	GLY	engineered mutation	UNP Q96SI1
F	1	SER	-	expression tag	UNP Q96SI1
F	2	MET	-	expression tag	UNP Q96SI1
F	39	ASP	GLY	engineered mutation	UNP Q96SI1
L	1	SER	-	expression tag	UNP Q96SI1
L	2	MET	-	expression tag	UNP Q96SI1
L	39	ASP	GLY	engineered mutation	UNP Q96SI1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	34	Total O 34 34	0	0
2	A	29	Total O 29 29	0	0
2	G	11	Total O 11 11	0	0
2	E	27	Total O 27 27	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	J	15	Total	O	0	0
			15	15		
2	D	14	Total	O	0	0
			14	14		
2	B	28	Total	O	0	0
			28	28		
2	I	13	Total	O	0	0
			13	13		
2	H	17	Total	O	0	0
			17	17		
2	K	22	Total	O	0	0
			22	22		
2	F	22	Total	O	0	0
			22	22		
2	L	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15




- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15




- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15




- Molecule 1: BTB/POZ domain-containing protein KCTD15





- Molecule 1: BTB/POZ domain-containing protein KCTD15



- Molecule 1: BTB/POZ domain-containing protein KCTD15



- Molecule 1: BTB/POZ domain-containing protein KCTD15



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	71.26Å 72.09Å 94.41Å 89.63° 77.08° 60.50°	Depositor
Resolution (Å)	61.31 – 1.94 61.31 – 1.95	Depositor EDS
% Data completeness (in resolution range)	65.1 (61.31-1.94) 65.2 (61.31-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.94Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.236 , 0.273 0.237 , 0.271	Depositor DCC
R_{free} test set	3780 reflections (3.25%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.007 for -h+k,k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9618	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/836	0.42	0/1141
1	B	0.18	0/795	0.36	0/1082
1	C	0.31	0/796	0.56	1/1080 (0.1%)
1	D	0.22	0/818	0.49	2/1116 (0.2%)
1	E	0.20	0/802	0.38	0/1095
1	F	0.20	0/763	0.41	0/1040
1	G	0.16	0/837	0.36	0/1141
1	H	0.38	0/799	0.97	7/1091 (0.6%)
1	I	0.24	0/806	0.49	0/1100
1	J	0.26	0/832	0.74	6/1138 (0.5%)
1	K	0.19	0/753	0.41	0/1026
1	L	0.17	0/781	0.35	0/1076
All	All	0.23	0/9618	0.53	16/13126 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	I	0	1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	36	LEU	CB-CG-CD2	-17.43	58.40	110.70
1	H	36	LEU	CB-CG-CD1	13.56	151.37	110.70
1	J	94	LEU	CD1-CG-CD2	-9.07	90.84	110.80
1	J	94	LEU	CB-CG-CD2	-8.43	85.42	110.70
1	J	94	LEU	CB-CG-CD1	8.03	134.78	110.70
1	C	59	GLU	CA-CB-CG	7.01	128.12	114.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	36	LEU	CA-C-N	-6.90	109.22	122.61
1	H	36	LEU	C-N-CA	-6.90	109.22	122.61
1	J	98	VAL	CG1-CB-CG2	6.21	124.47	110.80
1	J	94	LEU	CA-CB-CG	6.11	137.69	116.30
1	H	36	LEU	CD1-CG-CD2	-5.93	97.75	110.80
1	H	35	ARG	CA-C-N	-5.41	113.03	120.28
1	H	35	ARG	C-N-CA	-5.41	113.03	120.28
1	J	29	PRO	N-CD-CG	-5.18	95.43	103.20
1	D	73	LEU	CA-C-N	-5.05	115.50	123.13
1	D	73	LEU	C-N-CA	-5.05	115.50	123.13

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	I	90	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	814	0	741	10	0
1	B	775	0	703	3	0
1	C	776	0	715	5	0
1	D	797	0	708	3	0
1	E	781	0	702	7	0
1	F	743	0	658	7	0
1	G	816	0	730	2	0
1	H	779	0	685	9	0
1	I	785	0	698	8	0
1	J	811	0	700	10	0
1	K	735	0	667	5	0
1	L	760	0	609	4	0
2	A	29	0	0	3	0
2	B	28	0	0	2	0
2	C	34	0	0	1	0
2	D	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	27	0	0	0	0
2	F	22	0	0	0	0
2	G	11	0	0	1	0
2	H	17	0	0	0	0
2	I	13	0	0	1	0
2	J	15	0	0	0	0
2	K	22	0	0	0	0
2	L	14	0	0	0	0
All	All	9618	0	8316	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:ALA:O	2:A:201:HOH:O	1.90	0.88
1:I:56:ARG:NH1	2:I:201:HOH:O	2.13	0.81
1:J:95:GLN:HA	1:J:98:VAL:HG22	1.62	0.80
1:J:44:VAL:HG12	1:J:50:GLN:HG3	1.65	0.78
1:A:93:GLN:N	2:A:201:HOH:O	2.18	0.76
1:E:11:ILE:HD11	1:E:25:LEU:HD11	1.74	0.69
1:A:100:GLU:OE1	1:A:103:ARG:NH2	2.27	0.68
1:A:5:SER:N	2:A:202:HOH:O	2.27	0.67
1:D:74:LEU:HD11	1:F:84:LEU:HD22	1.76	0.66
1:B:14:GLY:O	2:B:201:HOH:O	2.13	0.65
1:G:88:GLU:OE2	2:G:201:HOH:O	2.15	0.64
1:G:59:GLU:O	1:G:62:ARG:HG2	1.98	0.64
1:E:36:LEU:O	1:E:40:THR:HG22	1.99	0.62
1:K:3:THR:HG23	1:K:6:ASN:HB2	1.82	0.61
1:L:70:THR:O	1:L:72:LYS:N	2.37	0.57
1:K:60:ILE:HG21	1:K:88:GLU:HG3	1.88	0.56
1:F:36:LEU:O	1:F:40:THR:HG22	2.05	0.56
1:I:54:ILE:HG22	1:I:56:ARG:HB2	1.88	0.56
1:I:56:ARG:HG3	1:I:88:GLU:OE1	2.06	0.55
1:C:11:ILE:HD11	1:C:25:LEU:HD11	1.87	0.55
1:D:36:LEU:O	1:D:40:THR:HB	2.08	0.53
1:J:28:TYR:CE2	1:J:96:PRO:HG2	2.44	0.52
1:H:95:GLN:HA	1:H:98:VAL:HG13	1.92	0.52
1:B:40:THR:O	2:B:202:HOH:O	2.19	0.52
1:C:87:GLU:OE1	1:I:72:LYS:HE2	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:56:ARG:HD2	1:K:91:TYR:CE2	2.46	0.51
1:C:11:ILE:HD11	1:C:25:LEU:CD1	2.40	0.51
1:B:12:ASP:HB2	1:B:17:MET:HE1	1.92	0.51
1:L:70:THR:C	1:L:72:LYS:H	2.17	0.51
1:J:68:LEU:HD21	1:J:94:LEU:CD1	2.41	0.51
1:J:95:GLN:HA	1:J:98:VAL:CG2	2.35	0.50
1:H:94:LEU:O	1:H:98:VAL:HG12	2.13	0.49
1:H:41:GLU:O	1:H:41:GLU:HG2	2.13	0.49
1:F:95:GLN:O	1:F:98:VAL:HG22	2.14	0.48
1:C:12:ASP:OD2	2:C:201:HOH:O	2.20	0.48
1:A:42:PRO:HD3	1:A:52:TYR:CZ	2.49	0.47
1:C:25:LEU:HD13	1:C:37:PHE:CZ	2.50	0.46
1:J:68:LEU:HA	1:J:68:LEU:HD23	1.70	0.45
1:F:95:GLN:HA	1:F:98:VAL:HG22	1.99	0.45
1:I:84:LEU:HD22	1:I:84:LEU:H	1.82	0.45
1:A:94:LEU:O	1:A:98:VAL:HG23	2.16	0.45
1:E:25:LEU:HD13	1:E:37:PHE:CZ	2.52	0.44
1:K:82:PHE:C	1:K:82:PHE:CD1	2.94	0.44
1:A:43:ILE:HG22	1:A:44:VAL:HG13	1.99	0.44
1:I:36:LEU:O	1:I:40:THR:OG1	2.25	0.44
1:K:67:PHE:CD2	1:K:97:MET:HB2	2.52	0.44
1:L:28:TYR:CE1	1:L:96:PRO:HG2	2.53	0.44
1:D:35:ARG:HG2	1:D:39:ASP:OD2	2.18	0.43
1:I:42:PRO:HB3	1:I:52:TYR:CE1	2.53	0.43
1:H:59:GLU:O	1:H:62:ARG:NH1	2.52	0.43
1:E:11:ILE:HG12	1:E:37:PHE:CZ	2.53	0.43
1:J:94:LEU:HA	1:J:94:LEU:HD22	1.86	0.43
1:F:56:ARG:HD2	1:F:91:TYR:CE2	2.53	0.43
1:E:60:ILE:HD13	1:E:84:LEU:HD23	2.01	0.42
1:H:56:ARG:HA	1:H:56:ARG:HD2	1.93	0.42
1:A:100:GLU:OE1	1:A:103:ARG:CZ	2.67	0.42
1:H:40:THR:C	1:H:42:PRO:HD3	2.44	0.42
1:L:6:ASN:HA	1:L:21:SER:HB2	2.02	0.42
1:I:86:TYR:CE1	1:I:90:ARG:HD3	2.55	0.41
1:F:40:THR:OG1	1:F:41:GLU:N	2.51	0.41
1:E:25:LEU:HD13	1:E:37:PHE:CE1	2.55	0.41
1:A:42:PRO:HD3	1:A:52:TYR:OH	2.20	0.41
1:H:11:ILE:HD11	1:H:25:LEU:HD11	2.02	0.41
1:H:22:LEU:HD23	1:H:22:LEU:HA	1.86	0.41
1:H:84:LEU:H	1:H:84:LEU:HD22	1.85	0.41
1:J:95:GLN:CA	1:J:98:VAL:HG22	2.44	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:ILE:HG13	1:A:92:TYR:CD1	2.56	0.41
1:E:11:ILE:HG12	1:E:37:PHE:HZ	1.84	0.41
1:J:56:ARG:NH2	1:J:88:GLU:OE2	2.48	0.40
1:J:14:GLY:HA2	1:F:19:THR:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/116 (88%)	97 (95%)	4 (4%)	1 (1%)	12	5
1	B	94/116 (81%)	90 (96%)	4 (4%)	0	100	100
1	C	91/116 (78%)	88 (97%)	3 (3%)	0	100	100
1	D	99/116 (85%)	94 (95%)	5 (5%)	0	100	100
1	E	96/116 (83%)	92 (96%)	4 (4%)	0	100	100
1	F	91/116 (78%)	86 (94%)	4 (4%)	1 (1%)	11	4
1	G	101/116 (87%)	97 (96%)	4 (4%)	0	100	100
1	H	98/116 (84%)	90 (92%)	7 (7%)	1 (1%)	12	5
1	I	98/116 (84%)	94 (96%)	4 (4%)	0	100	100
1	J	104/116 (90%)	100 (96%)	4 (4%)	0	100	100
1	K	91/116 (78%)	87 (96%)	4 (4%)	0	100	100
1	L	103/116 (89%)	98 (95%)	4 (4%)	1 (1%)	12	5
All	All	1168/1392 (84%)	1113 (95%)	51 (4%)	4 (0%)	36	30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	71	SER
1	H	42	PRO
1	F	41	GLU
1	A	43	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/109 (76%)	83 (100%)	0	100	100
1	B	79/109 (72%)	79 (100%)	0	100	100
1	C	82/109 (75%)	82 (100%)	0	100	100
1	D	77/109 (71%)	77 (100%)	0	100	100
1	E	79/109 (72%)	78 (99%)	1 (1%)	61	54
1	F	74/109 (68%)	74 (100%)	0	100	100
1	G	80/109 (73%)	80 (100%)	0	100	100
1	H	74/109 (68%)	73 (99%)	1 (1%)	59	52
1	I	76/109 (70%)	75 (99%)	1 (1%)	61	54
1	J	76/109 (70%)	75 (99%)	1 (1%)	61	54
1	K	75/109 (69%)	75 (100%)	0	100	100
1	L	65/109 (60%)	65 (100%)	0	100	100
All	All	920/1308 (70%)	916 (100%)	4 (0%)	84	84

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	51	HIS
1	J	98	VAL
1	I	56	ARG
1	H	98	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	10	HIS
1	E	16	HIS
1	E	38	ASN
1	J	16	HIS
1	I	10	HIS
1	I	38	ASN
1	H	10	HIS
1	K	6	ASN
1	K	16	HIS
1	L	51	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/116 (89%)	0.40	7 (6%) 24 27	29, 44, 75, 104	0
1	B	98/116 (84%)	0.31	2 (2%) 65 72	34, 48, 70, 89	0
1	C	95/116 (81%)	0.17	3 (3%) 50 56	28, 38, 74, 91	0
1	D	103/116 (88%)	0.53	6 (5%) 29 32	35, 50, 81, 92	0
1	E	100/116 (86%)	0.35	7 (7%) 22 25	30, 43, 72, 88	0
1	F	95/116 (81%)	0.41	6 (6%) 26 29	33, 42, 67, 83	0
1	G	105/116 (90%)	0.68	8 (7%) 20 22	37, 54, 80, 85	0
1	H	102/116 (87%)	0.77	10 (9%) 13 14	36, 50, 80, 109	0
1	I	102/116 (87%)	0.58	4 (3%) 43 49	34, 49, 75, 91	0
1	J	108/116 (93%)	0.87	13 (12%) 9 10	38, 59, 79, 104	0
1	K	95/116 (81%)	0.47	9 (9%) 14 15	35, 44, 71, 85	0
1	L	107/116 (92%)	0.73	11 (10%) 12 13	37, 58, 78, 86	0
All	All	1214/1392 (87%)	0.53	86 (7%) 22 25	28, 48, 78, 109	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	40	THR	5.6
1	C	38	ASN	5.5
1	E	42	PRO	5.1
1	H	36	LEU	4.9
1	A	45	LEU	4.5
1	H	49	LYS	4.4
1	I	110	GLN	4.2
1	F	42	PRO	4.1
1	L	43	ILE	4.0
1	F	104	TRP	3.8
1	K	82	PHE	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	40	THR	3.7
1	J	94	LEU	3.7
1	H	44	VAL	3.7
1	J	44	VAL	3.6
1	D	43	ILE	3.6
1	D	44	VAL	3.5
1	C	110	GLN	3.5
1	J	104	TRP	3.5
1	J	77	ASP	3.5
1	F	82	PHE	3.5
1	A	48	LEU	3.4
1	G	113	ARG	3.3
1	J	33	ILE	3.3
1	L	3	THR	3.2
1	E	37	PHE	3.2
1	F	73	LEU	3.1
1	D	113	ARG	3.1
1	J	74	LEU	3.1
1	K	44	VAL	3.1
1	F	3	THR	3.1
1	J	95	GLN	3.0
1	F	101	LEU	3.0
1	K	42	PRO	3.0
1	H	106	GLN	2.9
1	C	51	HIS	2.9
1	B	104	TRP	2.9
1	H	50	GLN	2.8
1	J	98	VAL	2.8
1	E	40	THR	2.7
1	H	41	GLU	2.7
1	L	51	HIS	2.6
1	J	86	TYR	2.6
1	A	104	TRP	2.6
1	E	110	GLN	2.6
1	E	51	HIS	2.5
1	G	75	LEU	2.5
1	L	77	ASP	2.5
1	G	108	GLN	2.5
1	H	86	TYR	2.5
1	A	44	VAL	2.5
1	D	74	LEU	2.5
1	K	102	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	43	ILE	2.4
1	G	42	PRO	2.4
1	J	6	ASN	2.4
1	L	74	LEU	2.4
1	I	78	ASP	2.4
1	I	77	ASP	2.3
1	K	3	THR	2.3
1	H	43	ILE	2.3
1	E	41	GLU	2.3
1	K	73	LEU	2.3
1	A	47	SER	2.3
1	K	74	LEU	2.3
1	L	30	ASP	2.2
1	J	78	ASP	2.2
1	K	78	ASP	2.2
1	A	43	ILE	2.2
1	E	43	ILE	2.2
1	I	43	ILE	2.2
1	L	104	TRP	2.2
1	A	78	ASP	2.2
1	D	78	ASP	2.2
1	J	4	LYS	2.2
1	L	70	THR	2.1
1	G	76	PRO	2.1
1	L	42	PRO	2.1
1	G	104	TRP	2.1
1	L	98	VAL	2.1
1	D	42	PRO	2.1
1	L	6	ASN	2.1
1	K	50	GLN	2.1
1	J	92	TYR	2.1
1	H	77	ASP	2.1
1	G	48	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.