



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:42 AM UTC

PDB ID : 2POO / pdb_00002poo
Title : THERMOSTABLE PHYTASE IN FULLY CALCIUM LOADED STATE
Authors : Ha, N.-C.; Oh, B.-H.
Deposited on : 1999-04-16
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	355	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (PHYTASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total	C	N	O	S	0	0	0
			2736	1711	461	558	6			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	Ca	0	0
			6	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total	O	0	0
			53	53		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTEIN (PHYTASE)

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.98Å 65.15Å 105.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.05	Depositor
% Data completeness (in resolution range)	92.9 (8.00-2.05)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	5.40	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.228 , 0.262	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2795	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2796	1.01	15/3785 (0.4%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	ASP	N-CA-C	7.49	121.72	111.54
1	A	116	ILE	N-CA-C	7.14	118.76	108.48
1	A	75	ASN	N-CA-C	-7.00	97.00	108.41
1	A	373	ASP	CA-C-N	6.30	126.50	119.32
1	A	373	ASP	C-N-CA	6.30	126.50	119.32
1	A	256	THR	CA-C-N	5.93	125.94	119.90
1	A	256	THR	C-N-CA	5.93	125.94	119.90
1	A	227	GLU	N-CA-C	5.91	123.40	110.80
1	A	123	SER	N-CA-C	5.52	118.23	110.23
1	A	207	ASN	N-CA-C	5.43	117.91	111.33
1	A	260	GLU	N-CA-C	5.42	113.22	108.78
1	A	299	PHE	N-CA-C	5.33	116.90	108.96
1	A	235	SER	N-CA-C	-5.13	102.16	110.32
1	A	55	ASP	N-CA-C	5.04	121.54	110.80
1	A	372	VAL	N-CA-C	5.02	115.97	108.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2736	0	2594	29	0
2	A	6	0	0	0	0
3	A	53	0	0	1	0
All	All	2795	0	2594	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:PRO:HD3	1:A:253:ARG:HH22	1.40	0.84
1:A:44:THR:HG22	1:A:349:ASN:HB2	1.69	0.73
1:A:160:GLY:H	1:A:177:THR:HG22	1.57	0.69
1:A:47:VAL:HG12	1:A:49:THR:H	1.58	0.67
1:A:133:ALA:HB3	1:A:142:GLN:HG2	1.82	0.60
1:A:129:ILE:O	1:A:129:ILE:HG13	2.01	0.59
1:A:307:THR:HG22	1:A:308:ASP:O	2.02	0.59
1:A:364:PHE:CE2	1:A:366:PRO:HG3	2.40	0.56
1:A:80:LEU:HD23	1:A:141:LEU:HD13	1.88	0.56
1:A:202:ARG:HH11	1:A:204:PHE:HE2	1.53	0.56
1:A:349:ASN:HD21	1:A:351:LYS:NZ	2.05	0.55
1:A:160:GLY:N	1:A:177:THR:HG22	2.23	0.54
1:A:58:ALA:HB1	1:A:103:ILE:HG23	1.91	0.52
1:A:103:ILE:HD11	1:A:105:TYR:CZ	2.44	0.51
1:A:216:ASP:OD2	1:A:289:ARG:HD3	2.11	0.51
1:A:272:GLY:C	1:A:289:ARG:HB2	2.36	0.50
1:A:290:GLN:H	1:A:290:GLN:CD	2.19	0.49
1:A:126:LYS:HD2	1:A:128:THR:HG23	1.97	0.47
1:A:349:ASN:HD21	1:A:351:LYS:HZ2	1.66	0.43
1:A:250:ALA:HB1	1:A:257:PRO:HG3	2.01	0.43
1:A:233:LYS:NZ	1:A:293:ASN:HD21	2.17	0.43
1:A:375:ARG:HD3	3:A:527:HOH:O	2.18	0.42
1:A:86:GLU:H	1:A:86:GLU:HG3	1.64	0.42
1:A:321:PHE:O	1:A:323:LEU:HD22	2.20	0.42
1:A:177:THR:HG23	1:A:178:GLY:N	2.35	0.41
1:A:210:THR:HG22	1:A:226:GLU:HA	2.02	0.41
1:A:32:ASP:HA	1:A:33:PRO:HA	1.86	0.41
1:A:67:GLN:H	1:A:67:GLN:CD	2.29	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ARG:NH1	1:A:204:PHE:HE2	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	351/355 (99%)	327 (93%)	22 (6%)	2 (1%)	21 13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	200	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	287/288 (100%)	256 (89%)	31 (11%)	6 2

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	52	ASP
1	A	65	ASN
1	A	67	GLN
1	A	77	LYS
1	A	80	LEU
1	A	86	GLU
1	A	89	MET
1	A	103	ILE
1	A	122	ARG
1	A	126	LYS
1	A	129	ILE
1	A	163	LEU
1	A	175	MET
1	A	176	VAL
1	A	177	THR
1	A	189	ASN
1	A	200	LYS
1	A	262	LEU
1	A	289	ARG
1	A	290	GLN
1	A	306	GLU
1	A	318	VAL
1	A	323	LEU
1	A	326	GLU
1	A	331	LEU
1	A	345	LYS
1	A	349	ASN
1	A	357	ARG
1	A	375	ARG
1	A	378	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	ASN
1	A	67	GLN
1	A	75	ASN
1	A	91	HIS
1	A	167	GLN
1	A	185	GLN
1	A	209	GLN
1	A	290	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	ASN
1	A	349	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.