



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:56 AM UTC

PDB ID : 3POS / pdb_00003pos
Title : Crystal structure of the globular domain of human calreticulin
Authors : Chouquet, A.; Paidassi, H.; Ling, W.-L.; Frachet, P.; Houen, G.; Arlaud, G.J.; Gaboriaud, C.
Deposited on : 2010-11-23
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	2% 85% 8% 6%
1	B	265	2% 89% 6% 5%
1	C	265	2% 84% 9% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calreticulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	3	0
			2025	1287	325	406	7			
1	B	253	Total	C	N	O	S	0	1	0
			2043	1298	331	407	7			
1	C	251	Total	C	N	O	S	0	3	0
			2038	1297	330	404	7			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	ASN	-	expression tag	UNP P27797
A	11	LYS	-	expression tag	UNP P27797
A	12	GLY	-	expression tag	UNP P27797
A	13	SER	-	expression tag	UNP P27797
A	14	ILE	-	expression tag	UNP P27797
A	15	GLU	-	expression tag	UNP P27797
A	16	GLY	-	expression tag	UNP P27797
A	17	ARG	-	expression tag	UNP P27797
A	299	GLY	-	linker	UNP P27797
A	300	SER	-	linker	UNP P27797
A	301	GLY	-	linker	UNP P27797
B	10	ASN	-	expression tag	UNP P27797
B	11	LYS	-	expression tag	UNP P27797
B	12	GLY	-	expression tag	UNP P27797
B	13	SER	-	expression tag	UNP P27797
B	14	ILE	-	expression tag	UNP P27797
B	15	GLU	-	expression tag	UNP P27797
B	16	GLY	-	expression tag	UNP P27797
B	17	ARG	-	expression tag	UNP P27797
B	299	GLY	-	linker	UNP P27797
B	300	SER	-	linker	UNP P27797
B	301	GLY	-	linker	UNP P27797
C	10	ASN	-	expression tag	UNP P27797

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	11	LYS	-	expression tag	UNP P27797
C	12	GLY	-	expression tag	UNP P27797
C	13	SER	-	expression tag	UNP P27797
C	14	ILE	-	expression tag	UNP P27797
C	15	GLU	-	expression tag	UNP P27797
C	16	GLY	-	expression tag	UNP P27797
C	17	ARG	-	expression tag	UNP P27797
C	299	GLY	-	linker	UNP P27797
C	300	SER	-	linker	UNP P27797
C	301	GLY	-	linker	UNP P27797

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	253	Total O 253 253	0	0
3	B	227	Total O 227 227	0	0
3	C	259	Total O 259 259	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Calreticulin



• Molecule 1: Calreticulin



• Molecule 1: Calreticulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.23Å 91.10Å 194.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 1.65 19.94 – 1.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.94-1.65) 96.8 (19.94-1.65)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 1.65Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.170 , 0.205 0.177 , 0.213	Depositor DCC
R_{free} test set	4379 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtrriage
Anisotropy	0.248	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6848	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7374e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	0/2073	1.04	2/2794 (0.1%)
1	B	0.78	0/2092	1.07	5/2820 (0.2%)
1	C	0.78	0/2086	1.04	7/2812 (0.2%)
All	All	0.78	0/6251	1.05	14/8426 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	310	ASN	CA-CB-CG	6.51	119.11	112.60
1	A	104	ASP	CA-CB-CG	6.26	118.86	112.60
1	C	328	ASP	CA-CB-CG	6.00	118.60	112.60
1	B	335	ASP	CA-CB-CG	5.98	118.58	112.60
1	C	71	ASP	CA-CB-CG	5.88	118.48	112.60
1	C	104	ASP	CA-CB-CG	5.83	118.43	112.60
1	C	310	ASN	CA-CB-CG	5.71	118.31	112.60
1	B	335	ASP	CA-C-N	5.66	127.86	120.28
1	B	335	ASP	C-N-CA	5.66	127.86	120.28
1	A	199	ASP	N-CA-C	5.30	119.73	113.16
1	B	104	ASP	CA-CB-CG	5.21	117.81	112.60
1	C	335	ASP	CA-CB-CG	5.16	117.76	112.60
1	C	307	ALA	N-CA-C	5.13	117.98	109.46
1	C	310	ASN	N-CA-C	5.06	116.75	108.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2025	0	1892	10	0
1	B	2043	0	1918	6	0
1	C	2038	0	1917	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	253	0	0	0	0
3	B	227	0	0	0	0
3	C	259	0	0	1	0
All	All	6848	0	5727	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:105:CYS:HG	1:C:137:CYS:HG	0.77	0.74
1:A:48:LYS:H	1:A:70:GLN:HE21	1.37	0.70
1:B:48:LYS:H	1:B:70:GLN:HE21	1.39	0.70
1:C:89:GLN:HE22	1:C:334:ASN:HD22	1.54	0.54
1:B:26:GLN:HG2	1:B:329:ASN:HD22	1.74	0.52
1:B:48:LYS:H	1:B:70:GLN:NE2	2.08	0.49
1:A:48:LYS:H	1:A:70:GLN:NE2	2.06	0.49
1:C:26:GLN:HG2	1:C:329:ASN:HD22	1.80	0.47
1:A:26:GLN:HG2	1:A:329:ASN:HD22	1.79	0.46
1:C:89:GLN:NE2	1:C:334:ASN:HD22	2.14	0.45
1:A:93[B]:VAL:HG22	1:A:112:LEU:HD11	1.97	0.45
1:C:48:LYS:H	1:C:70:GLN:HE21	1.62	0.45
1:B:67:GLN:HA	1:B:325:THR:O	2.18	0.44
1:C:27:PHE:CZ	1:C:66:LEU:HG	2.53	0.44
1:B:131[B]:MET:HB3	1:B:147:ILE:HB	2.00	0.44
1:C:43:LYS:HD3	3:C:430:HOH:O	2.17	0.44
1:C:143:LYS:HE2	1:C:145:HIS:CE1	2.53	0.43
1:C:159:LYS:HD3	1:C:192:GLU:CD	2.44	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:LYS:O	1:A:364:GLU:HG2	2.20	0.42
1:A:159:LYS:HD3	1:A:192:GLU:CD	2.45	0.42
1:B:159:LYS:HD3	1:B:192:GLU:CD	2.46	0.41
1:C:67:GLN:HA	1:C:325:THR:O	2.20	0.41
1:A:67:GLN:HB2	1:A:326:ILE:HG12	2.02	0.41
1:C:180:ASN:HD22	1:C:306:TYR:HB3	1.85	0.41
1:A:56:PHE:O	1:A:98:LYS:NZ	2.53	0.40
1:A:67:GLN:HA	1:A:325:THR:O	2.19	0.40
1:A:173:THR:HB	1:A:185:LYS:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/265 (93%)	235 (96%)	11 (4%)	0	100	100
1	B	250/265 (94%)	237 (95%)	13 (5%)	0	100	100
1	C	248/265 (94%)	236 (95%)	12 (5%)	0	100	100
All	All	744/795 (94%)	708 (95%)	36 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/231 (96%)	216 (98%)	5 (2%)	44	21
1	B	222/231 (96%)	218 (98%)	4 (2%)	51	30
1	C	222/231 (96%)	216 (97%)	6 (3%)	39	16
All	All	665/693 (96%)	650 (98%)	15 (2%)	44	21

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ASP
1	A	35	SER
1	A	48	LYS
1	A	142	LYS
1	A	143	LYS
1	B	31	ASP
1	B	143	LYS
1	B	180	ASN
1	B	310	ASN
1	C	89	GLN
1	C	142	LYS
1	C	180	ASN
1	C	203	LEU
1	C	310	ASN
1	C	318	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	89	GLN
1	A	149	ASN
1	A	154	ASN
1	A	180	ASN
1	A	329	ASN
1	B	70	GLN
1	B	89	GLN
1	B	180	ASN
1	B	329	ASN
1	B	356	GLN
1	C	26	GLN
1	C	70	GLN
1	C	89	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	149	ASN
1	C	154	ASN
1	C	180	ASN
1	C	329	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/265 (93%)	-0.32	3 (1%) 76 81	4, 14, 33, 66	3 (1%)
1	B	253/265 (95%)	-0.19	4 (1%) 70 75	5, 16, 37, 58	1 (0%)
1	C	251/265 (94%)	-0.37	4 (1%) 70 75	4, 14, 34, 65	3 (1%)
All	All	753/795 (94%)	-0.29	11 (1%) 72 77	4, 15, 36, 66	7 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	17	ARG	3.6
1	B	303	PRO	3.0
1	C	20	ALA	2.9
1	B	17	ARG	2.9
1	C	203	LEU	2.7
1	B	202	PHE	2.7
1	A	364	GLU	2.6
1	A	18	GLU	2.5
1	A	12	GLY	2.4
1	C	31	ASP	2.2
1	B	115	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	1	1/1	1.00	0.02	10,10,10,10	0
2	CA	B	2	1/1	1.00	0.01	10,10,10,10	0
2	CA	C	3	1/1	1.00	0.01	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.