



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:29 PM UTC

PDB ID : 6POL / pdb_00006pol
Title : Crystal structure of the human NELL1 EGF1-3-Robo3 FN1 complex
Authors : Wang, J.; Pak, J.S.; Ozkan, E.
Deposited on : 2019-07-04
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	114	6% 86% 11%
1	C	114	4% 82% 7% 11%
1	E	114	4% 82% 5% 12%
2	B	139	7% 91% 7%
2	D	139	8% 86% 7% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	139	9% 89% • • 7%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Roundabout homolog 3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	102	Total	C	N	O	0	0	0
			757	480	127	150			
1	C	101	Total	C	N	O	0	0	0
			751	476	126	149			
1	E	100	Total	C	N	O	0	1	0
			750	475	126	149			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	547	ALA	-	expression tag	UNP Q96MS0
A	548	ASP	-	expression tag	UNP Q96MS0
A	653	GLU	-	expression tag	UNP Q96MS0
A	654	PHE	-	expression tag	UNP Q96MS0
A	655	HIS	-	expression tag	UNP Q96MS0
A	656	HIS	-	expression tag	UNP Q96MS0
A	657	HIS	-	expression tag	UNP Q96MS0
A	658	HIS	-	expression tag	UNP Q96MS0
A	659	HIS	-	expression tag	UNP Q96MS0
A	660	HIS	-	expression tag	UNP Q96MS0
C	547	ALA	-	expression tag	UNP Q96MS0
C	548	ASP	-	expression tag	UNP Q96MS0
C	653	GLU	-	expression tag	UNP Q96MS0
C	654	PHE	-	expression tag	UNP Q96MS0
C	655	HIS	-	expression tag	UNP Q96MS0
C	656	HIS	-	expression tag	UNP Q96MS0
C	657	HIS	-	expression tag	UNP Q96MS0
C	658	HIS	-	expression tag	UNP Q96MS0
C	659	HIS	-	expression tag	UNP Q96MS0
C	660	HIS	-	expression tag	UNP Q96MS0
E	547	ALA	-	expression tag	UNP Q96MS0
E	548	ASP	-	expression tag	UNP Q96MS0
E	653	GLU	-	expression tag	UNP Q96MS0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	654	PHE	-	expression tag	UNP Q96MS0
E	655	HIS	-	expression tag	UNP Q96MS0
E	656	HIS	-	expression tag	UNP Q96MS0
E	657	HIS	-	expression tag	UNP Q96MS0
E	658	HIS	-	expression tag	UNP Q96MS0
E	659	HIS	-	expression tag	UNP Q96MS0
E	660	HIS	-	expression tag	UNP Q96MS0

- Molecule 2 is a protein called Protein kinase C-binding protein NELL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	129	Total	C	N	O	S	0	2	0
			967	577	170	201	19			
2	D	129	Total	C	N	O	S	0	2	0
			958	573	166	200	19			
2	F	129	Total	C	N	O	S	0	2	0
			958	573	166	200	19			

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	387	ALA	-	expression tag	UNP Q92832
B	388	ASP	-	expression tag	UNP Q92832
B	389	PRO	-	expression tag	UNP Q92832
B	518	GLU	-	expression tag	UNP Q92832
B	519	PHE	-	expression tag	UNP Q92832
B	520	HIS	-	expression tag	UNP Q92832
B	521	HIS	-	expression tag	UNP Q92832
B	522	HIS	-	expression tag	UNP Q92832
B	523	HIS	-	expression tag	UNP Q92832
B	524	HIS	-	expression tag	UNP Q92832
B	525	HIS	-	expression tag	UNP Q92832
D	387	ALA	-	expression tag	UNP Q92832
D	388	ASP	-	expression tag	UNP Q92832
D	389	PRO	-	expression tag	UNP Q92832
D	518	GLU	-	expression tag	UNP Q92832
D	519	PHE	-	expression tag	UNP Q92832
D	520	HIS	-	expression tag	UNP Q92832
D	521	HIS	-	expression tag	UNP Q92832
D	522	HIS	-	expression tag	UNP Q92832
D	523	HIS	-	expression tag	UNP Q92832
D	524	HIS	-	expression tag	UNP Q92832

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	525	HIS	-	expression tag	UNP Q92832
F	387	ALA	-	expression tag	UNP Q92832
F	388	ASP	-	expression tag	UNP Q92832
F	389	PRO	-	expression tag	UNP Q92832
F	518	GLU	-	expression tag	UNP Q92832
F	519	PHE	-	expression tag	UNP Q92832
F	520	HIS	-	expression tag	UNP Q92832
F	521	HIS	-	expression tag	UNP Q92832
F	522	HIS	-	expression tag	UNP Q92832
F	523	HIS	-	expression tag	UNP Q92832
F	524	HIS	-	expression tag	UNP Q92832
F	525	HIS	-	expression tag	UNP Q92832

- Molecule 3 is NITRATE ION (CCD ID: NO3) (formula: NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			4	1	3		
3	A	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	N	O	0	0
			4	1	3		
3	C	1	Total	N	O	0	0
			4	1	3		
3	D	1	Total	N	O	0	0
			4	1	3		
3	E	1	Total	N	O	0	0
			4	1	3		
3	E	1	Total	N	O	0	0
			4	1	3		
3	E	1	Total	N	O	0	0
			4	1	3		
3	F	1	Total	N	O	0	0
			4	1	3		
3	F	1	Total	N	O	0	0
			4	1	3		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	D	1	Total	Ca	0	0
			1	1		
5	F	1	Total	Ca	0	0
			1	1		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	70	Total	O	0	0
			70	70		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	140	Total 140	O 140	0	0
7	C	82	Total 82	O 82	0	0
7	D	129	Total 129	O 129	0	0
7	E	83	Total 83	O 83	0	0
7	F	138	Total 138	O 138	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Roundabout homolog 3




- Molecule 1: Roundabout homolog 3





- Molecule 1: Roundabout homolog 3





- Molecule 2: Protein kinase C-binding protein NELL1



- Molecule 2: Protein kinase C-binding protein NELL1



- Molecule 2: Protein kinase C-binding protein NELL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.00Å 87.00Å 211.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	51.47 – 1.80 51.47 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (51.47-1.80) 99.6 (51.47-1.80)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.81Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.230 , 0.267 0.234 , 0.269	Depositor DCC
R_{free} test set	1998 reflections (2.32%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.652	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.57$, $\langle L^2 \rangle = 0.42$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5896	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0219e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NO3, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/779	0.50	0/1076
1	C	0.33	0/773	0.50	0/1068
1	E	0.29	0/774	0.45	0/1069
2	B	0.30	0/989	0.50	0/1339
2	D	0.31	0/980	0.48	0/1328
2	F	0.29	0/980	0.49	1/1328 (0.1%)
All	All	0.30	0/5275	0.49	1/7208 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	457	LEU	CA-CB-CG	5.16	134.36	116.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	757	0	742	3	0
1	C	751	0	738	6	0
1	E	750	0	738	5	0
2	B	967	0	843	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	958	0	828	7	0
2	F	958	0	828	5	0
3	A	8	0	0	0	0
3	B	12	0	0	0	0
3	C	8	0	0	0	0
3	D	4	0	0	1	0
3	E	16	0	0	0	0
3	F	8	0	0	0	0
4	B	14	0	13	0	0
4	D	14	0	13	0	0
4	F	14	0	13	0	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
5	F	1	0	0	0	0
6	B	6	0	8	0	0
6	D	6	0	8	0	0
7	A	70	0	0	1	0
7	B	140	0	0	0	0
7	C	82	0	0	0	0
7	D	129	0	0	2	0
7	E	83	0	0	0	0
7	F	138	0	0	1	0
All	All	5896	0	4772	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:489:GLU:OE1	7:F:1001:HOH:O	2.11	0.68
1:A:575:THR:OG1	7:A:1101:HOH:O	2.13	0.65
2:D:494[B]:THR:OG1	2:D:501:SER:OG	2.18	0.61
2:D:477:HIS:O	7:D:1001:HOH:O	2.16	0.59
2:D:429:SER:OG	7:D:1002:HOH:O	2.17	0.58
2:F:494[B]:THR:OG1	2:F:501:SER:OG	2.24	0.55
1:A:555:PRO:HG3	2:B:443:HIS:HB2	1.88	0.53
1:C:555:PRO:HG3	2:D:443:HIS:HB2	1.91	0.53
1:E:555:PRO:HG3	2:F:443:HIS:HB2	1.91	0.52
2:B:494[B]:THR:OG1	2:B:501:SER:OG	2.29	0.50
1:E:622:ASN:OD1	1:E:648:ARG:HD3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:632:VAL:HG22	1:A:637:LEU:HD22	1.97	0.46
1:E:594:PHE:CZ	2:F:494[B]:THR:HG22	2.53	0.43
1:E:594:PHE:CE1	2:F:494[B]:THR:HG22	2.54	0.43
1:C:632:VAL:HG22	1:C:637:LEU:HD22	2.00	0.42
1:C:567:ILE:HG12	1:C:572:ILE:CD1	2.49	0.42
1:C:567:ILE:HG12	1:C:572:ILE:HD13	2.01	0.42
1:E:632:VAL:HG22	1:E:637:LEU:CD2	2.50	0.41
2:D:505:LYS:HE2	2:D:508:TYR:CE2	2.56	0.41
2:D:412:THR:OG1	3:D:903:NO3:O2	2.36	0.41
1:C:622:ASN:OD1	1:C:648:ARG:HD3	2.21	0.41
1:C:602:TRP:HB2	2:D:470:ASP:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/114 (88%)	98 (98%)	2 (2%)	0	100	100
1	C	99/114 (87%)	96 (97%)	3 (3%)	0	100	100
1	E	99/114 (87%)	98 (99%)	1 (1%)	0	100	100
2	B	129/139 (93%)	123 (95%)	6 (5%)	0	100	100
2	D	129/139 (93%)	123 (95%)	6 (5%)	0	100	100
2	F	129/139 (93%)	123 (95%)	6 (5%)	0	100	100
All	All	685/759 (90%)	661 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/97 (88%)	85 (100%)	0	100	100
1	C	85/97 (88%)	85 (100%)	0	100	100
1	E	85/97 (88%)	85 (100%)	0	100	100
2	B	108/117 (92%)	108 (100%)	0	100	100
2	D	106/117 (91%)	105 (99%)	1 (1%)	70	67
2	F	106/117 (91%)	105 (99%)	1 (1%)	70	67
All	All	575/642 (90%)	573 (100%)	2 (0%)	84	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	467	ILE
2	F	457	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	610	GLN
2	D	443	HIS
1	E	610	GLN
2	F	443	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 3 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	D	904	-	5,5,5	1.06	0	5,5,5	1.09	0
4	NAG	B	901	2	14,14,15	0.33	0	17,19,21	0.43	0
4	NAG	D	901	2	14,14,15	0.50	0	17,19,21	0.60	0
3	NO3	B	903	-	1,3,3	0.50	0	0,3,3	-	-
3	NO3	A	1002	-	1,3,3	0.60	0	0,3,3	-	-
3	NO3	E	1001	-	1,3,3	0.60	0	0,3,3	-	-
4	NAG	F	901	2	14,14,15	0.43	0	17,19,21	0.52	0
3	NO3	A	1001	-	1,3,3	0.50	0	0,3,3	-	-
3	NO3	E	1004	-	1,3,3	0.83	0	0,3,3	-	-
3	NO3	F	904	-	1,3,3	0.63	0	0,3,3	-	-
3	NO3	B	906	-	1,3,3	0.60	0	0,3,3	-	-
3	NO3	F	903	-	1,3,3	0.33	0	0,3,3	-	-
3	NO3	B	905	-	1,3,3	0.63	0	0,3,3	-	-
3	NO3	E	1002	-	1,3,3	0.61	0	0,3,3	-	-
3	NO3	C	1001	-	1,3,3	0.74	0	0,3,3	-	-
3	NO3	E	1003	-	1,3,3	0.88	0	0,3,3	-	-
3	NO3	D	903	-	1,3,3	0.46	0	0,3,3	-	-
3	NO3	C	1002	-	1,3,3	0.70	0	0,3,3	-	-
6	GOL	B	904	-	5,5,5	0.99	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	901	2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	F	901	2	-	2/6/23/26	0/1/1/1
6	GOL	D	904	-	-	4/4/4/4	-
4	NAG	D	901	2	-	1/6/23/26	0/1/1/1
6	GOL	B	904	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	D	904	GOL	O1-C1-C2-C3
6	D	904	GOL	C1-C2-C3-O3
4	B	901	NAG	O5-C5-C6-O6
6	D	904	GOL	O1-C1-C2-O2
6	D	904	GOL	O2-C2-C3-O3
4	F	901	NAG	O5-C5-C6-O6
4	B	901	NAG	C4-C5-C6-O6
4	D	901	NAG	C1-C2-N2-C7
4	F	901	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	903	NO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	102/114 (89%)	0.55	7 (6%)	23	21	29, 41, 72, 103	0
1	C	101/114 (88%)	0.54	4 (3%)	42	41	25, 38, 64, 112	0
1	E	100/114 (87%)	0.47	5 (5%)	34	33	29, 41, 64, 96	1 (1%)
2	B	129/139 (92%)	0.56	10 (7%)	19	17	20, 42, 76, 98	2 (1%)
2	D	129/139 (92%)	0.57	11 (8%)	16	14	18, 40, 71, 101	2 (1%)
2	F	129/139 (92%)	0.60	12 (9%)	14	12	19, 41, 72, 97	2 (1%)
All	All	690/759 (90%)	0.55	49 (7%)	22	20	18, 41, 73, 112	7 (1%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	552	PRO	6.2
1	A	552	PRO	4.7
1	A	582	THR	4.4
2	B	514	ILE	4.1
1	C	582	THR	3.8
2	F	514	ILE	3.7
2	D	390	ARG	3.7
2	F	390	ARG	3.5
1	E	652	SER	3.4
2	D	514	ILE	3.4
1	A	653	GLU	3.3
2	F	484	GLN	3.1
2	B	390	ARG	3.0
2	B	484	GLN	3.0
2	F	517	ALA	3.0
2	B	516	ARG	2.8
1	E	554	SER	2.8
2	D	399	PRO	2.7
1	E	582	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	486	ASN	2.6
1	A	554	SER	2.6
2	D	518	GLU	2.6
2	F	518	GLU	2.5
1	A	652	SER	2.5
2	F	399	PRO	2.5
1	A	651	ASP	2.5
2	D	509	VAL	2.5
2	B	489	GLU	2.4
1	A	553	SER	2.4
2	D	441	LYS	2.4
2	D	484	GLN	2.4
2	F	441	LYS	2.4
2	D	391	GLY	2.4
2	B	517	ALA	2.3
2	F	482	SER	2.3
2	B	518	GLU	2.3
2	D	517	ALA	2.3
1	C	652	SER	2.3
2	B	486	ASN	2.2
2	F	516	ARG	2.2
1	E	553	SER	2.2
2	B	483	GLY	2.2
2	F	510	GLY	2.1
1	C	569	LYS	2.1
2	F	440	ALA	2.1
2	F	486	ASN	2.1
1	E	570	ASN	2.1
2	B	399	PRO	2.1
2	D	457	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	F	901	14/15	0.53	0.16	65,78,82,82	0
3	NO3	B	905	4/4	0.64	0.15	87,87,87,88	0
4	NAG	D	901	14/15	0.65	0.15	64,79,81,81	0
4	NAG	B	901	14/15	0.66	0.16	60,75,80,80	0
6	GOL	D	904	6/6	0.73	0.19	55,57,60,61	0
3	NO3	B	906	4/4	0.76	0.17	77,78,78,78	0
3	NO3	F	904	4/4	0.77	0.13	78,79,79,79	0
3	NO3	E	1004	4/4	0.78	0.17	44,49,50,53	0
3	NO3	A	1002	4/4	0.78	0.22	77,78,78,78	0
3	NO3	C	1002	4/4	0.80	0.16	57,58,58,59	0
3	NO3	E	1002	4/4	0.83	0.14	60,61,61,61	0
6	GOL	B	904	6/6	0.84	0.14	48,50,51,55	0
3	NO3	E	1003	4/4	0.86	0.17	42,44,46,48	0
3	NO3	E	1001	4/4	0.87	0.11	48,49,50,51	0
3	NO3	F	903	4/4	0.88	0.13	47,47,49,49	0
3	NO3	B	903	4/4	0.90	0.15	45,46,47,48	0
3	NO3	D	903	4/4	0.92	0.11	45,45,46,48	0
3	NO3	A	1001	4/4	0.92	0.09	50,51,52,52	0
3	NO3	C	1001	4/4	0.93	0.11	41,45,45,45	0
5	CA	D	902	1/1	0.99	0.02	30,30,30,30	0
5	CA	F	902	1/1	0.99	0.02	29,29,29,29	0
5	CA	B	902	1/1	1.00	0.02	29,29,29,29	0

6.5 Other polymers

There are no such residues in this entry.