



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:12 AM UTC

PDB ID : 8PO0 / pdb_00008po0
Title : Discovery and Optimisation of Potent, Efficacious and Selective Inhibitors Targeting EGFR Exon20 Insertion Mutations. Compound 12 bound to EGFRin-sNPG
Authors : Hargreaves, D.
Deposited on : 2023-07-03
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	17% 63% 27% • 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2503 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2466	1580	413	455	18			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	694	GLY	-	expression tag	UNP P00533
A	773	GLY	-	insertion	UNP P00533
A	774	ASN	-	insertion	UNP P00533
A	775	PRO	-	insertion	UNP P00533

- Molecule 2 is 1-[3-[7-methoxy-4-[3-(3-methoxyphenyl)-1 {H}-pyrazol-4-yl]quinazolin-6-yl]oxyxyazetidin-1-yl]propan-1-one (CCD ID: 2I6) (formula: C₂₅H₂₅N₅O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			34	25	5	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	150.77Å 150.77Å 150.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.68 – 2.52 47.68 – 2.52	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.68-2.52) 99.6 (47.68-2.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.8 (24-FEB-2021)	Depositor
R, R_{free}	0.315 , 0.376 0.308 , 0.381	Depositor DCC
R_{free} test set	989 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	87.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 67.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2503	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2I6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/2518	1.21	13/3407 (0.4%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	858	ASP	CA-CB-CG	7.02	119.62	112.60
1	A	797	PRO	N-CA-C	6.66	122.50	113.84
1	A	792	ILE	N-CA-C	5.52	116.06	107.28
1	A	740	ILE	N-CA-C	5.29	113.76	107.73
1	A	761	ASP	CA-C-N	5.26	127.28	120.44
1	A	761	ASP	C-N-CA	5.26	127.28	120.44
1	A	744	ILE	N-CA-C	5.26	115.64	107.28
1	A	837	VAL	CA-C-N	5.13	127.16	120.28
1	A	837	VAL	C-N-CA	5.13	127.16	120.28
1	A	976	ARG	CA-C-N	5.10	126.76	122.28
1	A	976	ARG	C-N-CA	5.10	126.76	122.28
1	A	983	VAL	N-CA-C	5.09	111.66	106.21
1	A	942	CYS	N-CA-C	5.03	116.51	108.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2466	0	2472	42	0
2	A	34	0	0	2	0
3	A	3	0	0	0	0
All	All	2503	0	2472	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:798:PHE:HB2	1:A:848:VAL:HB	1.50	0.94
1:A:958:ALA:HA	1:A:961:ARG:NE	1.82	0.94
1:A:958:ALA:HA	1:A:961:ARG:HE	1.38	0.87
1:A:774:ASN:HD22	1:A:776:HIS:H	1.32	0.77
1:A:929:ILE:HA	1:A:932:LYS:HE2	1.67	0.77
1:A:907:VAL:HG12	1:A:950:MET:HE2	1.76	0.66
1:A:836:LEU:HD11	1:A:862:ALA:HB1	1.80	0.64
1:A:774:ASN:ND2	1:A:776:HIS:H	1.99	0.61
1:A:964:PHE:O	1:A:968:ILE:HG13	2.00	0.61
1:A:809:LYS:HG3	1:A:913:PHE:HB3	1.83	0.60
1:A:988:GLU:H	1:A:988:GLU:CD	2.07	0.60
1:A:798:PHE:HB2	1:A:848:VAL:CB	2.30	0.56
1:A:776:HIS:HA	1:A:855:LYS:HG2	1.86	0.55
1:A:821:CYS:O	1:A:903:TYR:OH	2.24	0.55
1:A:935:ARG:NH1	1:A:954:TRP:O	2.41	0.53
1:A:732:ILE:HG12	1:A:739:LYS:HG2	1.91	0.53
1:A:912:THR:HB	1:A:915:SER:HB2	1.91	0.52
1:A:801:LEU:HD12	1:A:804:TYR:HB3	1.92	0.51
1:A:847:LEU:HG	1:A:857:THR:HG21	1.94	0.50
1:A:771:ASN:HB2	1:A:772:PRO:HD3	1.93	0.50
1:A:1015:ASP:OD1	1:A:1017:ASP:OD1	2.30	0.49
1:A:828:MET:HE2	1:A:841:LEU:HD22	1.97	0.47
1:A:907:VAL:HA	1:A:910:LEU:HD12	1.96	0.46
1:A:911:MET:HG3	1:A:950:MET:HE1	1.97	0.46
1:A:802:LEU:HD13	1:A:843:ALA:HB3	1.98	0.46
1:A:929:ILE:HG22	1:A:930:LEU:HD23	1.97	0.46
1:A:784:CYS:HB3	1:A:790:GLN:HB2	1.96	0.46
1:A:785:LEU:HA	1:A:789:VAL:HG22	1.98	0.46
1:A:881:ILE:HD11	1:A:889:ILE:HG23	1.97	0.45
1:A:744:ILE:HG12	1:A:792:ILE:HD12	1.98	0.45
1:A:912:THR:CB	1:A:915:SER:HB2	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:800:CYS:SG	1:A:803:ASP:OD2	2.75	0.44
1:A:763:ALA:HB2	1:A:791:LEU:HD21	2.00	0.43
2:A:1101:2I6:C23	2:A:1101:2I6:C19	2.97	0.42
1:A:836:LEU:HD21	1:A:862:ALA:HB1	2.01	0.42
2:A:1101:2I6:C23	2:A:1101:2I6:C8	2.99	0.41
1:A:769:VAL:O	1:A:830:TYR:OH	2.28	0.41
1:A:885:ALA:HB1	1:A:887:GLU:OE2	2.21	0.41
1:A:771:ASN:CB	1:A:772:PRO:HD3	2.51	0.41
1:A:832:GLU:HA	1:A:896:HIS:CE1	2.56	0.41
1:A:882:LYS:HG3	1:A:918:TYR:HD2	1.86	0.41
1:A:926:ILE:O	1:A:930:LEU:HG	2.21	0.41
1:A:702:ALA:HB2	1:A:768:SER:HA	2.03	0.40
1:A:921:ILE:HA	1:A:922:PRO:HD3	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	304/332 (92%)	267 (88%)	33 (11%)	4 (1%)	9 17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	890	LEU
1	A	723	PHE
1	A	766	MET
1	A	771	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	269/290 (93%)	237 (88%)	32 (12%)	5 9

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	707	LEU
1	A	709	GLU
1	A	718	LEU
1	A	730	LEU
1	A	734	GLU
1	A	746	GLU
1	A	757	LYS
1	A	759	ILE
1	A	760	LEU
1	A	766	MET
1	A	779	ARG
1	A	783	ILE
1	A	785	LEU
1	A	815	GLN
1	A	817	LEU
1	A	819	ASN
1	A	821	CYS
1	A	861	LEU
1	A	863	LYS
1	A	868	GLU
1	A	879	VAL
1	A	881	ILE
1	A	887	GLU
1	A	891	HIS
1	A	929	ILE
1	A	942	CYS
1	A	951	VAL
1	A	965	ARG
1	A	967	LEU
1	A	988	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1006	ASP
1	A	1008	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	700	ASN
1	A	774	ASN
1	A	790	GLN
1	A	819	ASN
1	A	896	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2I6	A	1101	1	37,38,38	0.71	2 (5%)	45,54,54	1.44	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2I6	A	1101	1	-	9/20/30/30	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	2I6	C18-C17	2.51	1.58	1.53
2	A	1101	2I6	C19-C17	2.06	1.57	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	2I6	C8-N1-N	-5.42	104.30	112.59
2	A	1101	2I6	C17-C18-N4	-4.21	83.79	87.96
2	A	1101	2I6	C17-C19-N4	-3.84	84.15	87.96
2	A	1101	2I6	C18-N4-C19	-3.32	92.37	94.69
2	A	1101	2I6	C7-N-N1	2.41	112.61	106.66
2	A	1101	2I6	C9-C8-N1	2.39	112.22	108.17

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	2I6	N4-C20-C21-C22
2	A	1101	2I6	O3-C20-N4-C18
2	A	1101	2I6	C6-C1-O-C
2	A	1101	2I6	C2-C1-O-C
2	A	1101	2I6	O3-C20-C21-C22
2	A	1101	2I6	C23-C16-O2-C17
2	A	1101	2I6	C19-C17-O2-C16
2	A	1101	2I6	C18-C17-O2-C16
2	A	1101	2I6	O3-C20-N4-C19

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	2I6	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/332 (93%)	1.37	57 (18%) 3 3	70, 95, 126, 135	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	918	TYR	6.5
1	A	1020	LEU	5.2
1	A	812	ILE	4.7
1	A	1004	LEU	4.6
1	A	929	ILE	4.1
1	A	760	LEU	4.1
1	A	862	ALA	3.8
1	A	759	ILE	3.5
1	A	1013	VAL	3.3
1	A	926	ILE	3.2
1	A	879	VAL	3.2
1	A	1010	MET	3.1
1	A	983	VAL	3.1
1	A	785	LEU	3.0
1	A	763	ALA	2.9
1	A	754	LYS	2.9
1	A	976	ARG	2.8
1	A	757	LYS	2.8
1	A	881	ILE	2.8
1	A	755	ALA	2.8
1	A	744	ILE	2.7
1	A	851	PRO	2.6
1	A	765	VAL	2.6
1	A	743	ALA	2.6
1	A	761	ASP	2.6
1	A	703	LEU	2.6
1	A	773	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	730	LEU	2.5
1	A	778	CYS	2.4
1	A	869	GLU	2.4
1	A	864	LEU	2.4
1	A	767	ALA	2.4
1	A	910	LEU	2.3
1	A	792	ILE	2.3
1	A	901	TRP	2.3
1	A	801	LEU	2.3
1	A	712	PHE	2.3
1	A	923	ALA	2.3
1	A	941	ILE	2.3
1	A	769	VAL	2.3
1	A	775	PRO	2.2
1	A	805	VAL	2.2
1	A	771	ASN	2.2
1	A	865	LEU	2.2
1	A	706	ILE	2.2
1	A	984	ILE	2.2
1	A	1021	ILE	2.2
1	A	774	ASN	2.1
1	A	992	LEU	2.1
1	A	946	VAL	2.1
1	A	989	ARG	2.1
1	A	909	GLU	2.1
1	A	726	VAL	2.0
1	A	762	GLU	2.0
1	A	780	LEU	2.0
1	A	883	TRP	2.0
1	A	939	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	2I6	A	1101	34/34	0.82	0.17	91,92,97,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.