



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:29 AM UTC

PDB ID : 8PPE / pdb_00008ppe
Title : Human inositol 1,4,5-trisphosphate 3-kinase A (IP3K) catalytic domain in complex with DL-6-deoxy-6-phosphoryloxymethyl-scylo-inositol 1,2,4-trisphosphate/ADP/Mn
Authors : Marquez-Monino, M.A.; Gonzalez, B.
Deposited on : 2023-07-07
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	16% 89% 8% ••
1	B	279	24% 85% 9% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Inositol-trisphosphate 3-kinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	2	0
			2203	1376	400	413	14			
1	B	263	Total	C	N	O	S	0	3	0
			2132	1333	386	399	14			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	183	GLY	-	expression tag	UNP P23677
A	184	SER	-	expression tag	UNP P23677
A	185	HIS	-	expression tag	UNP P23677
A	186	MET	-	expression tag	UNP P23677
A	187	SER	-	expression tag	UNP P23677
B	183	GLY	-	expression tag	UNP P23677
B	184	SER	-	expression tag	UNP P23677
B	185	HIS	-	expression tag	UNP P23677
B	186	MET	-	expression tag	UNP P23677
B	187	SER	-	expression tag	UNP P23677

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 3 is DL-6-deoxy-6-phosphoryloxymethyl-scylo-inositol 1,2,4-trisphosphate (CCD ID: 7I8) (formula: $C_7H_{18}O_{18}P_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	1
			58	14	36	8		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	P	0	0
			29	7	18	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mn	0	0
			1	1		
5	B	1	Total	Mn	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	198	Total	O	0	0
			198	198		
6	B	190	Total	O	0	0
			190	190		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Inositol-trisphosphate 3-kinase A



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.55Å 97.55Å 190.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.28 – 1.59 58.28 – 1.59	Depositor EDS
% Data completeness (in resolution range)	88.2 (58.28-1.59) 88.3 (58.28-1.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.59Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.200 , 0.240 0.209 , 0.250	Depositor DCC
R_{free} test set	4019 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4872	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 7I8, ADP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/2245	1.26	1/3019 (0.0%)
1	B	1.01	0/2171	1.28	1/2917 (0.0%)
All	All	1.00	0/4416	1.27	2/5936 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	430	ASP	CA-CB-CG	5.21	117.81	112.60
1	A	430	ASP	CA-CB-CG	5.16	117.75	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2203	0	2168	16	0
1	B	2132	0	2108	15	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	58	0	0	0	0
3	B	29	0	0	0	0
4	A	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	198	0	0	1	0
6	B	190	0	0	1	0
All	All	4872	0	4308	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:316[B]:MET:HA	1:B:316[B]:MET:HE2	1.64	0.80
1:B:316[B]:MET:HA	1:B:316[B]:MET:CE	2.15	0.77
1:A:285:ARG:HG3	1:A:288:MET:HE3	1.85	0.58
1:A:286:LYS:HG3	1:A:308:ARG:O	2.04	0.58
1:A:288:MET:HE1	1:A:312:LYS:HG3	1.87	0.56
1:A:183:GLY:O	1:A:253:ASP:HA	2.08	0.53
1:B:291:LYS:O	1:B:294:ALA:HB3	2.09	0.53
1:A:293:LEU:HD12	1:A:297:PRO:HA	1.93	0.50
1:A:338:ALA:HB1	1:B:294:ALA:HB1	1.95	0.48
1:A:285:ARG:CB	1:A:288:MET:HG3	2.43	0.48
1:A:255:PHE:CZ	4:A:503:GOL:H2	2.51	0.46
1:B:299:ALA:HB3	1:B:300:PRO:HD3	1.97	0.45
1:B:292:MET:HG3	1:B:300:PRO:HG3	1.97	0.45
1:A:285:ARG:CZ	1:A:288:MET:HE2	2.47	0.44
1:A:184:SER:HA	1:A:250:ASP:OD2	2.18	0.44
1:A:285:ARG:HB2	1:A:288:MET:HG3	1.99	0.44
1:A:299:ALA:HB3	1:A:300:PRO:HD3	2.00	0.44
1:A:213:GLU:N	1:A:214:PRO:CD	2.81	0.43
1:B:291:LYS:HB3	6:B:701:HOH:O	2.18	0.43
1:B:213:GLU:N	1:B:214:PRO:CD	2.82	0.43
1:B:292:MET:HE2	1:B:292:MET:HB2	1.93	0.42
1:B:261:LEU:C	1:B:261:LEU:HD23	2.43	0.42
1:B:289:TYR:HD1	1:B:310:VAL:HG13	1.84	0.42
1:A:264:LYS:HD2	1:A:397:GLY:HA2	2.02	0.42
1:A:341:SER:HB2	1:B:345:ASP:HA	2.02	0.41
1:A:319:ARG:HD2	6:A:660:HOH:O	2.20	0.41
1:B:312:LYS:HB3	1:B:313:PRO:HD3	2.02	0.41
1:B:423:LEU:HD21	1:B:429:LEU:HG	2.02	0.40
1:B:436:GLU:O	1:B:437:GLU:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/279 (97%)	268 (99%)	2 (1%)	0	100	100
1	B	262/279 (94%)	257 (98%)	5 (2%)	0	100	100
All	All	532/558 (95%)	525 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/239 (99%)	232 (98%)	5 (2%)	47	23
1	B	229/239 (96%)	221 (96%)	8 (4%)	32	11
All	All	466/478 (98%)	453 (97%)	13 (3%)	38	15

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	213	GLU
1	A	240	ARG
1	A	264	LYS
1	A	288	MET
1	A	446	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	264	LYS
1	B	285	ARG
1	B	288	MET
1	B	291	LYS
1	B	295	VAL
1	B	305	HIS
1	B	342	CYS
1	B	392	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	247	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	503	-	5,5,5	0.17	0	5,5,5	0.51	0
3	7I8	B	502	-	29,29,29	0.79	0	44,47,47	1.54	3 (6%)
2	ADP	A	501	5	28,29,29	0.51	0	43,45,45	0.43	0
2	ADP	B	501	5	28,29,29	0.56	1 (3%)	43,45,45	0.43	0
3	7I8	A	502[B]	5	29,29,29	0.81	1 (3%)	44,47,47	1.51	3 (6%)
3	7I8	A	502[A]	5	29,29,29	0.80	1 (3%)	44,47,47	1.50	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	503	-	-	4/4/4/4	-
3	7I8	B	502	-	-	10/21/45/45	0/1/1/1
2	ADP	A	501	5	-	1/16/32/32	0/3/3/3
2	ADP	B	501	5	-	0/16/32/32	0/3/3/3
3	7I8	A	502[B]	5	-	5/21/45/45	0/1/1/1
3	7I8	A	502[A]	5	-	9/21/45/45	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502[B]	7I8	P2-O2	2.16	1.63	1.59
2	B	501	ADP	PA-O3A	2.04	1.61	1.59
3	A	502[A]	7I8	P2-O2	2.03	1.63	1.59

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502[B]	7I8	P4-O4-C4	7.94	144.65	123.43
3	A	502[A]	7I8	P4-O4-C4	7.84	144.37	123.43
3	B	502	7I8	P4-O4-C4	7.72	144.06	123.43
3	B	502	7I8	P1-O1-C1	4.41	135.21	123.43
3	A	502[B]	7I8	P1-O1-C1	3.65	133.19	123.43
3	A	502[A]	7I8	P1-O1-C1	3.65	133.19	123.43
3	B	502	7I8	P2-O2-C2	3.20	131.99	123.43
3	A	502[B]	7I8	P2-O2-C2	2.75	130.77	123.43
3	A	502[A]	7I8	P2-O2-C2	2.52	130.15	123.43

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502[A]	7I8	C06-O36-P6-O46
3	A	502[A]	7I8	C06-O36-P6-O66
3	A	502[B]	7I8	C06-O36-P6-O66
3	B	502	7I8	C06-O36-P6-O56
3	B	502	7I8	C06-O36-P6-O46
3	B	502	7I8	C06-O36-P6-O66
3	B	502	7I8	O36-C06-C6-C1
3	B	502	7I8	O36-C06-C6-C5
3	B	502	7I8	C3-C4-O4-P4
3	B	502	7I8	C5-C4-O4-P4
4	A	503	GOL	O1-C1-C2-C3
4	A	503	GOL	C1-C2-C3-O3
4	A	503	GOL	O2-C2-C3-O3
3	A	502[A]	7I8	C3-C4-O4-P4
4	A	503	GOL	O1-C1-C2-O2
3	A	502[A]	7I8	C06-O36-P6-O56
3	A	502[A]	7I8	C5-C4-O4-P4
3	A	502[A]	7I8	O36-C06-C6-C1
3	A	502[B]	7I8	C06-O36-P6-O56
3	B	502	7I8	C1-O1-P1-O21
3	A	502[A]	7I8	C1-O1-P1-O31
3	A	502[B]	7I8	C1-O1-P1-O31
3	A	502[B]	7I8	C06-O36-P6-O46
3	A	502[A]	7I8	C6-C06-O36-P6
3	B	502	7I8	C6-C06-O36-P6
3	A	502[B]	7I8	C1-O1-P1-O11
3	B	502	7I8	C1-O1-P1-O31
3	A	502[A]	7I8	O36-C06-C6-C5
2	A	501	ADP	PB-O3A-PA-O2A


There are no ring outliers.



1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is


within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand 7I8 A 502 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/279 (97%)	1.03	46 (16%) 4 3	12, 33, 66, 86	2 (0%)
1	B	263/279 (94%)	1.27	66 (25%) 1 1	15, 32, 95, 114	3 (1%)
All	All	535/558 (95%)	1.15	112 (20%) 2 2	12, 33, 86, 114	5 (0%)

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	293	LEU	9.2
1	B	284	LEU	8.0
1	B	295	VAL	7.3
1	B	278	ALA	6.6
1	A	188	TRP	6.4
1	B	338	ALA	6.2
1	A	189	VAL	6.2
1	B	294	ALA	6.0
1	B	282	PRO	5.9
1	A	338	ALA	5.7
1	B	297	PRO	5.5
1	A	295	VAL	5.3
1	B	198	PHE	4.9
1	B	301	THR	4.7
1	A	294	ALA	4.7
1	A	293	LEU	4.7
1	B	289	TYR	4.4
1	B	296	ASP	4.4
1	A	183	GLY	4.4
1	B	306	ALA	4.3
1	B	310	VAL	4.1
1	B	309	ALA	4.1
1	B	291	LYS	4.0
1	B	292	MET	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	305	HIS	3.8
1	B	242	GLY	3.7
1	B	200	ALA	3.7
1	B	299	ALA	3.7
1	B	300	PRO	3.6
1	B	288	MET	3.6
1	A	289	TYR	3.5
1	B	276	THR	3.5
1	B	196	GLY	3.4
1	A	290	LYS	3.3
1	A	341	SER	3.3
1	B	318	TRP	3.3
1	B	290	LYS	3.2
1	B	302	GLU	3.2
1	A	185	HIS	3.2
1	B	204	SER	3.2
1	B	340	GLY	3.2
1	A	424	PRO	3.1
1	B	237	VAL	3.1
1	A	243	GLU	3.1
1	B	342	CYS	3.0
1	A	242	GLY	2.9
1	B	426	GLY	2.9
1	A	335	ILE	2.9
1	B	339	ASP	2.9
1	A	284	LEU	2.9
1	A	283	LYS	2.9
1	A	186	MET	2.9
1	A	187	SER	2.9
1	B	240	ARG	2.8
1	A	286	LYS	2.8
1	B	311	THR	2.8
1	A	184	SER	2.7
1	B	246	LEU	2.7
1	B	238	VAL	2.7
1	B	277	LYS	2.7
1	A	211	CYS	2.6
1	A	201	ALA	2.6
1	A	203	THR	2.6
1	B	285	ARG	2.6
1	A	342	CYS	2.5
1	B	298	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	308	ARG	2.5
1	A	198	PHE	2.5
1	B	210	ARG	2.5
1	A	309	ALA	2.5
1	A	202	GLY	2.5
1	B	239	GLU	2.4
1	A	299	ALA	2.4
1	A	306	ALA	2.4
1	B	202	GLY	2.4
1	B	197	SER	2.4
1	B	307	GLN	2.4
1	B	211	CYS	2.4
1	B	461	ARG	2.4
1	A	241	ASP	2.3
1	A	339	ASP	2.3
1	B	425	ASP	2.3
1	B	424	PRO	2.3
1	B	283	LYS	2.3
1	B	275	LEU	2.3
1	A	426	GLY	2.3
1	A	277	LYS	2.3
1	B	199	LYS	2.3
1	A	288	MET	2.3
1	A	336	LYS	2.3
1	B	459	ALA	2.3
1	B	243	GLU	2.3
1	B	313	PRO	2.3
1	A	278	ALA	2.2
1	B	303	GLU	2.2
1	A	425	ASP	2.2
1	B	214	PRO	2.2
1	B	357	ARG	2.2
1	B	201	ALA	2.2
1	A	281	ARG	2.2
1	A	213	GLU	2.2
1	B	208	LEU	2.1
1	B	207	ILE	2.1
1	A	197	SER	2.1
1	B	241	ASP	2.1
1	A	292	MET	2.1
1	A	210	ARG	2.1
1	A	346	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	214	PRO	2.1
1	B	287	ASP	2.1
1	A	357	ARG	2.0
1	B	392	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	7I8	A	502[A]	29/29	0.74	0.13	62,75,85,86	29
3	7I8	A	502[B]	29/29	0.74	0.13	43,56,66,68	29
3	7I8	B	502	29/29	0.75	0.12	87,96,107,109	0
2	ADP	B	501	27/27	0.80	0.14	32,43,61,65	27
4	GOL	A	503	6/6	0.81	0.17	46,48,48,52	0
2	ADP	A	501	27/27	0.83	0.14	30,40,54,58	27
5	MN	B	503	1/1	0.88	0.14	51,51,51,51	1
5	MN	A	504	1/1	0.97	0.12	35,35,35,35	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 7I8 A 502 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 7I8 A 502 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 7I8 B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP A 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.