



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 02:53 PM UTC

PDB ID : 2PQF / pdb_00002pqf
Title : Human Poly(ADP-Ribose) Polymerase 12, Catalytic fragment in complex with an inhibitor 3-Aminobenzoic acid
Authors : Karlberg, T.; Lehtio, L.; Arrowsmith, C.H.; Berglund, H.; Busam, R.D.; Collins, R.; Dahlgren, L.G.; Edwards, A.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hogbom, M.; Johansson, I.; Kallas, A.; Kotenyova, T.; Moche, M.; Nordlund, P.; Nyman, T.; Persson, C.; Sagemark, J.; Sundstrom, M.; Thorsell, A.G.; Van Den Berg, S.; Weigelt, J.; Holmberg-Schiavone, L.; Structural Genomics Consortium (SGC)
Deposited on : 2007-05-02
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	
1	B	198	
1	C	198	

Continued on next page...

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	198	 5% 80% 12% • 8%
1	E	198	 5% 81% 11% • 7%
1	F	198	 2% 83% 8% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CIT	B	801	-	X	-	-
3	CIT	E	801	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 12.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	Se	0	1	0
			1519	976	260	278	3	2			
1	B	181	Total	C	N	O	S	Se	0	0	0
			1479	950	252	272	3	2			
1	C	185	Total	C	N	O	S	Se	0	0	0
			1513	972	259	277	3	2			
1	D	182	Total	C	N	O	S	Se	0	1	0
			1498	967	254	272	3	2			
1	E	185	Total	C	N	O	S	Se	0	0	0
			1513	972	259	277	3	2			
1	F	183	Total	C	N	O	S	Se	0	0	0
			1493	961	254	273	3	2			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	487	SER	-	cloning artifact	UNP Q9H0J9
A	488	MSE	-	cloning artifact	UNP Q9H0J9
A	548	MSE	MET	modified residue	UNP Q9H0J9
A	617	MSE	MET	modified residue	UNP Q9H0J9
B	487	SER	-	cloning artifact	UNP Q9H0J9
B	488	MSE	-	cloning artifact	UNP Q9H0J9
B	548	MSE	MET	modified residue	UNP Q9H0J9
B	617	MSE	MET	modified residue	UNP Q9H0J9
C	487	SER	-	cloning artifact	UNP Q9H0J9
C	488	MSE	-	cloning artifact	UNP Q9H0J9
C	548	MSE	MET	modified residue	UNP Q9H0J9
C	617	MSE	MET	modified residue	UNP Q9H0J9
D	487	SER	-	cloning artifact	UNP Q9H0J9
D	488	MSE	-	cloning artifact	UNP Q9H0J9
D	548	MSE	MET	modified residue	UNP Q9H0J9
D	617	MSE	MET	modified residue	UNP Q9H0J9
E	487	SER	-	cloning artifact	UNP Q9H0J9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	488	MSE	-	cloning artifact	UNP Q9H0J9
E	548	MSE	MET	modified residue	UNP Q9H0J9
E	617	MSE	MET	modified residue	UNP Q9H0J9
F	487	SER	-	cloning artifact	UNP Q9H0J9
F	488	MSE	-	cloning artifact	UNP Q9H0J9
F	548	MSE	MET	modified residue	UNP Q9H0J9
F	617	MSE	MET	modified residue	UNP Q9H0J9

- Molecule 2 is 3-AMINOBENZOIC ACID (CCD ID: GAB) (formula: $C_7H_7NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	7	1	2		
2	B	1	Total	C	N	O	0	0
			10	7	1	2		
2	C	1	Total	C	N	O	0	0
			10	7	1	2		
2	D	1	Total	C	N	O	0	0
			10	7	1	2		
2	E	1	Total	C	N	O	0	0
			10	7	1	2		
2	F	1	Total	C	N	O	0	0
			10	7	1	2		

- Molecule 3 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		
3	D	1	Total	C	O	0	0
			13	6	7		
3	E	1	Total	C	O	0	0
			13	6	7		
3	F	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	66	Total	O	0	0
			66	66		
4	B	51	Total	O	0	0
			51	51		
4	C	62	Total	O	0	0
			62	62		
4	D	42	Total	O	0	0
			42	42		
4	E	56	Total	O	0	0
			56	56		
4	F	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poly [ADP-ribose] polymerase 12




- Molecule 1: Poly [ADP-ribose] polymerase 12




- Molecule 1: Poly [ADP-ribose] polymerase 12




- Molecule 1: Poly [ADP-ribose] polymerase 12







- Molecule 1: Poly [ADP-ribose] polymerase 12



- Molecule 1: Poly [ADP-ribose] polymerase 12



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	206.59Å 206.59Å 84.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.60 – 2.20 19.60 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.60-2.20) 99.8 (19.60-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.3.0032	Depositor
R, R_{free}	0.198 , 0.244 0.199 , 0.245	Depositor DCC
R_{free} test set	1810 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.044 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9471	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, GAB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	0/1562	0.93	0/2112
1	B	0.99	1/1518 (0.1%)	0.98	3/2050 (0.1%)
1	C	0.97	2/1557 (0.1%)	0.96	4/2106 (0.2%)
1	D	1.06	2/1540 (0.1%)	0.96	0/2080
1	E	1.06	1/1557 (0.1%)	0.95	0/2106
1	F	0.82	1/1536 (0.1%)	0.90	1/2078 (0.0%)
All	All	0.99	7/9270 (0.1%)	0.95	8/12532 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	1	1
All	All	1	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	637	PRO	CA-C	6.41	1.56	1.52
1	D	622	VAL	CB-CG2	5.46	1.70	1.52
1	B	557	VAL	CA-CB	5.39	1.60	1.54
1	C	622	VAL	CA-CB	5.31	1.61	1.54
1	E	545	LYS	C-O	-5.28	1.18	1.24

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	636	ARG	CA-C-N	-6.54	114.95	119.66
1	C	636	ARG	C-N-CA	-6.54	114.95	119.66
1	C	672	GLU	N-CA-C	6.28	118.21	111.36
1	C	554	GLY	N-CA-C	-5.67	107.73	114.48
1	F	672	GLU	N-CA-C	5.17	118.06	111.69

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	C	645	ASN	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	680	SER	Peptide
1	C	645	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1519	0	1434	15	0
1	B	1479	0	1402	8	0
1	C	1513	0	1433	10	0
1	D	1498	0	1419	13	0
1	E	1513	0	1433	14	0
1	F	1493	0	1400	9	0
2	A	10	0	6	2	0
2	B	10	0	6	1	0
2	C	10	0	6	1	0
2	D	10	0	6	3	0
2	E	10	0	6	2	0
2	F	10	0	6	0	0
3	A	13	0	5	1	0
3	B	13	0	5	1	0
3	C	13	0	5	1	0
3	D	13	0	5	2	0
3	E	13	0	5	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	13	0	5	0	0
4	A	66	0	0	0	0
4	B	51	0	0	0	0
4	C	62	0	0	0	0
4	D	42	0	0	0	0
4	E	56	0	0	0	0
4	F	41	0	0	0	0
All	All	9471	0	8587	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:644:SER:O	1:C:645:ASN:HB3	1.47	1.13
1:B:520:THR:HG21	1:B:600:ASP:HB3	1.65	0.77
1:B:598:ALA:HB2	2:B:701:GAB:HN31	1.52	0.75
1:D:517:PHE:HD1	1:D:617:MSE:HE1	1.51	0.74
1:D:598:ALA:HB2	2:D:701:GAB:HN31	1.53	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/198 (92%)	179 (98%)	3 (2%)	0	100	100
1	B	175/198 (88%)	173 (99%)	2 (1%)	0	100	100
1	C	183/198 (92%)	178 (97%)	5 (3%)	0	100	100
1	D	177/198 (89%)	171 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	183/198 (92%)	180 (98%)	3 (2%)	0	100	100
1	F	179/198 (90%)	177 (99%)	2 (1%)	0	100	100
All	All	1079/1188 (91%)	1058 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/171 (95%)	151 (93%)	11 (7%)	14	17
1	B	159/171 (93%)	153 (96%)	6 (4%)	29	40
1	C	162/171 (95%)	153 (94%)	9 (6%)	19	24
1	D	160/171 (94%)	154 (96%)	6 (4%)	29	40
1	E	162/171 (95%)	155 (96%)	7 (4%)	26	35
1	F	158/171 (92%)	149 (94%)	9 (6%)	18	23
All	All	963/1026 (94%)	915 (95%)	48 (5%)	22	28

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	586	VAL
1	E	555	LYS
1	D	643	TRP
1	E	506	SER
1	E	641	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	547	GLN
1	E	499	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	631	ASN
1	D	676	GLN
1	E	515	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CIT	D	801	-	12,12,12	1.50	1 (8%)	17,17,17	2.40	9 (52%)
3	CIT	E	801	-	12,12,12	2.37	6 (50%)	17,17,17	5.72	12 (70%)
2	GAB	E	701	-	10,10,10	1.01	1 (10%)	13,13,13	1.80	3 (23%)
3	CIT	B	801	-	12,12,12	1.98	3 (25%)	17,17,17	2.08	8 (47%)
3	CIT	A	801	-	12,12,12	2.07	2 (16%)	17,17,17	1.86	5 (29%)
3	CIT	F	801	-	12,12,12	1.24	0	17,17,17	2.23	7 (41%)
2	GAB	A	701	-	10,10,10	1.23	0	13,13,13	1.90	4 (30%)
2	GAB	B	701	-	10,10,10	1.10	2 (20%)	13,13,13	1.78	3 (23%)
3	CIT	C	801	-	12,12,12	1.84	3 (25%)	17,17,17	1.57	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GAB	F	701	-	10,10,10	1.12	0	13,13,13	1.70	2 (15%)
2	GAB	D	701	-	10,10,10	1.02	1 (10%)	13,13,13	1.57	3 (23%)
2	GAB	C	701	-	10,10,10	1.17	0	13,13,13	1.86	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	D	801	-	-	6/16/16/16	-
3	CIT	E	801	-	-	6/16/16/16	-
2	GAB	E	701	-	-	4/4/4/4	0/1/1/1
3	CIT	B	801	-	-	7/16/16/16	-
3	CIT	A	801	-	-	5/16/16/16	-
3	CIT	F	801	-	-	1/16/16/16	-
2	GAB	A	701	-	-	2/4/4/4	0/1/1/1
2	GAB	B	701	-	-	4/4/4/4	0/1/1/1
3	CIT	C	801	-	-	3/16/16/16	-
2	GAB	F	701	-	-	4/4/4/4	0/1/1/1
2	GAB	D	701	-	-	4/4/4/4	0/1/1/1
2	GAB	C	701	-	-	0/4/4/4	0/1/1/1

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	CIT	C3-C6	-5.56	1.47	1.53
3	B	801	CIT	C3-C6	-4.57	1.48	1.53
3	E	801	CIT	C3-C6	-4.37	1.48	1.53
3	C	801	CIT	C3-C6	-4.15	1.49	1.53
3	A	801	CIT	O7-C3	3.48	1.49	1.43

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	801	CIT	C4-C3-C2	13.67	144.38	109.31
3	E	801	CIT	O7-C3-C4	-11.00	84.28	109.38
3	E	801	CIT	O7-C3-C6	9.48	122.41	108.96
3	E	801	CIT	O2-C1-O1	-5.81	108.38	123.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	801	CIT	O7-C3-C2	-5.01	97.95	109.38

There are no chirality outliers.

5 of 46 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	801	CIT	C2-C3-C4-C5
3	D	801	CIT	O7-C3-C4-C5
3	D	801	CIT	C6-C3-C4-C5
3	E	801	CIT	C1-C2-C3-O7
3	A	801	CIT	O7-C3-C4-C5

There are no ring outliers.

10 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	801	CIT	2	0
3	E	801	CIT	1	0
2	E	701	GAB	2	0
3	B	801	CIT	1	0
3	A	801	CIT	1	0
2	A	701	GAB	2	0
2	B	701	GAB	1	0
3	C	801	CIT	1	0
2	D	701	GAB	3	0
2	C	701	GAB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/198 (92%)	0.39	15 (8%) 17 15	26, 55, 68, 81	1 (0%)
1	B	179/198 (90%)	0.57	9 (5%) 34 31	45, 55, 64, 67	0
1	C	183/198 (92%)	0.32	4 (2%) 62 59	47, 56, 65, 72	0
1	D	180/198 (90%)	0.51	9 (5%) 34 31	29, 55, 64, 69	1 (0%)
1	E	183/198 (92%)	0.33	9 (4%) 35 31	44, 55, 65, 69	0
1	F	181/198 (91%)	0.40	4 (2%) 62 59	47, 56, 65, 71	0
All	All	1089/1188 (91%)	0.42	50 (4%) 37 34	26, 55, 65, 81	2 (0%)

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	643	TRP	4.9
1	B	586	VAL	4.5
1	E	496	PRO	4.0
1	D	681	SER	3.9
1	D	525[A]	PHE	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CIT	F	801	13/13	0.77	0.20	86,92,96,96	0
3	CIT	E	801	13/13	0.80	0.15	61,66,68,69	0
3	CIT	D	801	13/13	0.85	0.15	70,76,79,80	0
2	GAB	D	701	10/10	0.85	0.15	54,57,59,60	0
3	CIT	B	801	13/13	0.85	0.14	60,67,70,71	0
2	GAB	B	701	10/10	0.86	0.15	51,57,57,59	0
3	CIT	C	801	13/13	0.88	0.12	60,68,73,74	0
2	GAB	E	701	10/10	0.89	0.14	47,54,55,59	0
3	CIT	A	801	13/13	0.90	0.12	59,67,71,72	0
2	GAB	C	701	10/10	0.91	0.13	46,47,49,53	0
2	GAB	F	701	10/10	0.91	0.11	52,54,57,60	0
2	GAB	A	701	10/10	0.92	0.11	48,54,56,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.