



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:23 PM UTC

PDB ID : 7PQ0 / pdb_00007pq0
Title : Crystal structure of the Burkholderia Lethal Factor 1 (BLF1) C94S inactive mutant in complex with human eIF4A - Crystal form B
Authors : Mobbs, G.W.; Aziz, A.A.; Dix, S.R.; Blackburn, G.M.; Sedelnikova, S.E.; Minshull, T.C.; Dickman, M.J.; Baker, P.J.; Nathan, S.; Firdaus-Raih, M.; Rice, D.W.
Deposited on : 2021-09-15
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	% 78% 20% .
2	B	394	2% 76% 19% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4716 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Burkholderia Lethal Factor 1 (BLF1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1638	1031	283	319	5			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	SER	CYS	engineered mutation	UNP Q63UP7

- Molecule 2 is a protein called Eukaryotic initiation factor 4A-I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	385	Total	C	N	O	S	0	0	0
			3078	1941	534	584	19			


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	13	MET	-	initiating methionine	UNP P60842
B	14	HIS	-	expression tag	UNP P60842
B	15	HIS	-	expression tag	UNP P60842
B	16	HIS	-	expression tag	UNP P60842
B	17	HIS	-	expression tag	UNP P60842
B	18	HIS	-	expression tag	UNP P60842
B	19	HIS	-	expression tag	UNP P60842


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Burkholderia Lethal Factor 1 (BLF1)



• Molecule 2: Eukaryotic initiation factor 4A-I



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	199.53Å 199.53Å 51.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	65.31 – 3.00 65.31 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (65.31-3.00) 99.7 (65.31-3.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.76 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.186 , 0.230 0.188 , 0.230	Depositor DCC
R_{free} test set	1208 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	67.4	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 70.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4716	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1680	1.13	0/2284
2	B	0.92	3/3124 (0.1%)	1.15	6/4216 (0.1%)
All	All	0.90	3/4804 (0.1%)	1.14	6/6500 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	332	LEU	N-CA	6.47	1.54	1.46
2	B	232	ILE	C-O	-5.79	1.17	1.24
2	B	23	ILE	C-O	-5.40	1.17	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	330	ASP	CA-CB-CG	6.39	118.99	112.60
2	B	240	GLU	CB-CG-CD	5.99	122.78	112.60
2	B	26	ASN	N-CA-C	-5.50	104.82	112.30
2	B	330	ASP	CB-CA-C	-5.42	101.71	109.84
2	B	71	ASP	CA-CB-CG	5.13	117.73	112.60
2	B	55	PRO	CB-CA-C	-5.11	104.69	111.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1638	0	1564	34	0
2	B	3078	0	3112	46	0
All	All	4716	0	4676	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:187:MET:HE3	2:B:196:ILE:HD11	1.40	1.01
2:B:187:MET:HE3	2:B:196:ILE:CD1	2.19	0.72
1:A:73:THR:HG1	1:A:105:TRP:CD1	2.08	0.72
1:A:94:SER:OG	2:B:339:GLN:OE1	2.04	0.72
1:A:128:ASP:O	1:A:135:ARG:NH2	2.25	0.68
1:A:100:ASP:HB2	1:A:136:GLN:HB2	1.75	0.68
1:A:9:ILE:HG22	1:A:13:MET:HE2	1.74	0.68
1:A:135:ARG:HG2	1:A:135:ARG:HH11	1.59	0.68
2:B:387:ILE:H	2:B:387:ILE:HD13	1.60	0.67
1:A:94:SER:HG	2:B:339:GLN:CD	2.02	0.67
2:B:384:LEU:O	2:B:386:ASP:N	2.28	0.67
2:B:110:ARG:HG3	2:B:135:ILE:HG23	1.76	0.66
1:A:13:MET:HE1	1:A:85:LEU:HD22	1.80	0.64
2:B:165:MET:HE3	2:B:170:TYR:HB2	1.79	0.64
1:A:109:ALA:HB3	1:A:110:PRO:HD3	1.82	0.60
1:A:134:ASP:HB2	1:A:207:GLN:O	2.02	0.60
1:A:71:LEU:HD12	1:A:71:LEU:N	2.16	0.60
2:B:29:GLU:OE2	2:B:64:LEU:HD23	2.03	0.58
2:B:101:THR:HA	2:B:152:PRO:O	2.03	0.58
2:B:331:LEU:HB3	2:B:334:ARG:HG3	1.86	0.57
1:A:135:ARG:HG2	1:A:135:ARG:NH1	2.19	0.57
2:B:110:ARG:NH2	2:B:111:GLU:HG2	2.21	0.56
2:B:356:TYR:CE2	2:B:360:ILE:HD11	2.41	0.55
2:B:211:LEU:C	2:B:211:LEU:HD23	2.33	0.54
1:A:134:ASP:HB3	1:A:208:GLN:HA	1.89	0.54
1:A:100:ASP:CB	1:A:136:GLN:HB2	2.38	0.53
2:B:144:VAL:HG22	2:B:165:MET:HE1	1.90	0.53
2:B:31:VAL:HG21	2:B:64:LEU:HD22	1.92	0.52
1:A:71:LEU:HD21	1:A:116:PHE:HB2	1.93	0.50
1:A:71:LEU:O	1:A:72:GLN:NE2	2.44	0.50
2:B:110:ARG:HH22	2:B:111:GLU:HG2	1.77	0.50
1:A:145:GLN:O	1:A:146:SER:C	2.53	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:131:CYS:HA	2:B:154:ILE:O	2.11	0.49
2:B:90:ILE:HD11	2:B:180:VAL:HG21	1.94	0.49
1:A:34:THR:HG23	2:B:194:ASP:OD2	2.12	0.49
2:B:108:PRO:HB3	2:B:187:MET:HG3	1.94	0.49
1:A:182:GLN:O	1:A:183:GLN:HG3	2.13	0.49
2:B:292:MET:O	2:B:297:PHE:HB2	2.14	0.48
1:A:27:LEU:HB3	1:A:63:TRP:CZ2	2.48	0.47
2:B:64:LEU:HB3	2:B:65:PRO:CD	2.44	0.47
2:B:383:THR:HG22	2:B:383:THR:O	2.14	0.47
2:B:277:ILE:HB	2:B:327:ILE:HG22	1.95	0.47
1:A:142:ARG:HB2	1:A:164:TYR:CZ	2.49	0.47
1:A:91:LEU:HG	1:A:94:SER:HB2	1.97	0.47
2:B:330:ASP:OD1	2:B:359:ARG:NE	2.44	0.46
2:B:75:GLN:HA	2:B:213:SER:O	2.15	0.46
2:B:291:LYS:O	2:B:295:ARG:HG3	2.15	0.46
2:B:255:ARG:HD2	2:B:257:GLU:OE2	2.15	0.45
1:A:151:ASN:OD1	1:A:153:SER:OG	2.31	0.45
2:B:286:ASP:O	2:B:289:THR:HG22	2.16	0.45
1:A:203:THR:HG22	1:A:204:TYR:N	2.32	0.45
2:B:340:GLN:O	2:B:341:VAL:HG13	2.17	0.45
1:A:10:ARG:NH1	1:A:83:ASP:O	2.51	0.44
2:B:143:GLU:HA	2:B:143:GLU:OE2	2.17	0.44
2:B:216:MET:N	2:B:217:PRO:CD	2.80	0.43
2:B:116:ILE:O	2:B:120:VAL:HG23	2.18	0.43
1:A:98:ALA:HB3	1:A:138:ALA:HB3	2.00	0.43
2:B:253:VAL:HG12	2:B:255:ARG:O	2.17	0.43
2:B:318:PHE:CD1	2:B:326:LEU:HB2	2.53	0.43
2:B:133:ALA:HA	2:B:156:VAL:O	2.18	0.43
1:A:116:PHE:O	1:A:117:TRP:C	2.59	0.43
2:B:266:LEU:HD22	2:B:270:LEU:HD11	2.00	0.43
1:A:86:PHE:O	1:A:175:GLY:HA2	2.18	0.43
1:A:113:ALA:HA	1:A:116:PHE:CE2	2.55	0.42
2:B:65:PRO:HB3	2:B:232:ILE:CD1	2.49	0.42
2:B:266:LEU:CD2	2:B:398:MET:HE1	2.50	0.42
2:B:380:ASP:O	2:B:384:LEU:HG	2.20	0.41
1:A:170:GLY:HA2	2:B:320:SER:O	2.20	0.41
1:A:62:LEU:HD12	1:A:184:MET:HE1	2.02	0.41
1:A:164:TYR:CD1	1:A:172:ALA:HB2	2.55	0.41
2:B:287:TRP:O	2:B:291:LYS:HG2	2.21	0.41
1:A:71:LEU:N	1:A:71:LEU:CD1	2.82	0.41
2:B:383:THR:O	2:B:383:THR:CG2	2.69	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:TRP:HE1	1:A:134:ASP:HA	1.86	0.40
2:B:298:THR:HB	2:B:323:SER:HA	2.02	0.40
2:B:302:MET:HB2	2:B:314:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/211 (99%)	195 (94%)	13 (6%)	0	100	100
2	B	383/394 (97%)	363 (95%)	17 (4%)	3 (1%)	16	50
All	All	591/605 (98%)	558 (94%)	30 (5%)	3 (0%)	24	60

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	385	ARG
2	B	254	GLU
2	B	273	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/174 (99%)	171 (99%)	2 (1%)	63	82
2	B	339/348 (97%)	318 (94%)	21 (6%)	16	49
All	All	512/522 (98%)	489 (96%)	23 (4%)	24	59

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	LEU
1	A	135	ARG
2	B	23	ILE
2	B	58	ILE
2	B	156	VAL
2	B	167	ASN
2	B	189	SER
2	B	219	ASP
2	B	235	LEU
2	B	298	THR
2	B	299	VAL
2	B	309	LYS
2	B	314	ILE
2	B	327	ILE
2	B	330	ASP
2	B	332	LEU
2	B	338	VAL
2	B	341	VAL
2	B	384	LEU
2	B	387	ILE
2	B	394	SER
2	B	400	LEU
2	B	402	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	136	GLN
1	A	145	GLN
1	A	207	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	77	GLN
2	B	167	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	210/211 (99%)	-0.21	2 (0%)	79 59	74, 111, 151, 196	0
2	B	385/394 (97%)	-0.10	7 (1%)	67 44	67, 87, 138, 164	0
All	All	595/605 (98%)	-0.14	9 (1%)	72 49	67, 96, 146, 196	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	402	VAL	3.6
2	B	403	ALA	3.1
2	B	404	ASP	2.8
2	B	332	LEU	2.4
2	B	400	LEU	2.2
2	B	138	THR	2.1
1	A	130	GLU	2.1
2	B	331	LEU	2.0
1	A	2	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.