



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 19, 2026 – 08:19 PM UTC

PDB ID : 7PR7 / pdb_00007pr7
Title : Crystal structure of human heparanase in complex with covalent inhibitor VL166
Authors : Wu, L.; Armstrong, Z.; Davies, G.J.
Deposited on : 2021-09-21
Resolution : 1.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	
2	B	74	
3	C	2	
4	D	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EDO	A	605	-	X	-	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7993 atoms, of which 3865 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heparanase 50 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	385	6212	1993	3125	526	556	12	89	4	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	159	LYS	-	expression tag	UNP Q9Y251
A	307	ARG	LYS	variant	UNP Q9Y251

- Molecule 2 is a protein called Heparanase 8 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	74	1204	388	609	97	110	14	1	0

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	2	48	14	24	1	9	5	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-(2R,3S,5R,6R)-2,3,4,5,6-pentakis(oxidanyl)cyclohexane-1-carboxylic acid.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
4	D	2	50	15	23	1	11	6	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	A	1	Total 28	C 8	H 14	N 1	O 5	3	0
5	A	1	Total 28	C 8	H 14	N 1	O 5	3	0
5	A	1	Total 28	C 8	H 14	N 1	O 5	3	0

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	1	0
			10	2	6	2		
6	A	1	Total	C	H	O	1	0
			10	2	6	2		
6	A	1	Total	C	H	O	1	0
			10	2	6	2		
6	A	1	Total	C	H	O	1	0
			10	2	6	2		
6	A	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	276	Total	O	0	0
			276	276		
8	B	48	Total	O	0	0
			48	48		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heparanase 50 kDa subunit



- Molecule 2: Heparanase 8 kDa subunit



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-(2R,3S,5R,6R)-2,3,4,5,6-pentakis (oxidanyl)cyclohexane-1-carboxylic acid



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.40Å 71.01Å 79.23Å 90.00° 95.14° 90.00°	Depositor
Resolution (Å)	46.26 – 1.52 46.26 – 1.52	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.26-1.52) 97.8 (46.26-1.52)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.52Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.166 , 0.188 0.165 , 0.186	Depositor DCC
R_{free} test set	3946 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtrriage
Anisotropy	0.224	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7993	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, EDO, CL, FUC, 8I4, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.03	3/3162 (0.1%)	1.23	4/4276 (0.1%)
2	B	1.06	0/609	1.22	1/826 (0.1%)
All	All	1.04	3/3771 (0.1%)	1.23	5/5102 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	343	GLU	CD-OE2	6.76	1.38	1.25
1	A	203	ASN	C-O	5.14	1.30	1.24
1	A	407	SER	CA-CB	-5.08	1.45	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	64	ASN	CA-CB-CG	5.83	118.44	112.60
1	A	425	GLY	CA-C-O	-5.71	118.51	122.22
1	A	271	PRO	N-CA-C	5.41	119.79	112.48
1	A	175	THR	CA-CB-OG1	-5.29	101.66	109.60
1	A	225	GLU	CB-CA-C	5.25	115.57	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3087	3125	3111	17	0
2	B	595	609	604	8	0
3	C	24	24	22	0	0
4	D	27	23	12	1	0
5	A	42	42	39	0	0
6	A	28	42	41	5	0
7	A	1	0	0	0	0
8	A	276	0	0	1	0
8	B	48	0	0	4	0
All	All	4128	3865	3829	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298[B]:TYR:CE1	1:A:345[B]:SER:OG	1.72	1.43
1:A:298[B]:TYR:HE1	1:A:345[B]:SER:OG	1.12	1.11
1:A:298[B]:TYR:CD1	1:A:345[B]:SER:OG	2.34	0.79
2:B:46[B]:GLN:NE2	8:B:201:HOH:O	2.00	0.79
1:A:489:LEU:HG	6:A:605:EDO:H11	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/385 (100%)	377 (97%)	10 (3%)	0	100	100
2	B	73/74 (99%)	73 (100%)	0	0	100	100
All	All	460/459 (100%)	450 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/335 (101%)	335 (99%)	3 (1%)	70	49
2	B	67/66 (102%)	67 (100%)	0	100	100
All	All	405/401 (101%)	402 (99%)	3 (1%)	76	57

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	LEU
1	A	391	TYR
1	A	392	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	GLN
1	A	472	ASN
1	A	494	GLN
2	B	64	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.88	0	17,19,21	1.80	2 (11%)
3	FUC	C	2	3	10,10,11	1.19	1 (10%)	14,14,16	1.55	4 (28%)
4	8I4	D	1	1,4	13,13,14	1.40	4 (30%)	18,19,21	1.66	5 (27%)
4	NDG	D	2	4	14,14,15	1.11	1 (7%)	17,19,21	1.35	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	FUC	C	2	3	-	-	0/1/1/1
4	8I4	D	1	1,4	-	2/4/24/28	0/1/1/1
4	NDG	D	2	4	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	2	NDG	O5-C5	-2.45	1.38	1.43
3	C	2	FUC	O5-C5	2.40	1.48	1.43
4	D	1	8I4	C1-C2	2.27	1.57	1.52
4	D	1	8I4	O6B-C6	-2.12	1.23	1.30
4	D	1	8I4	O6A-C6	2.10	1.28	1.22

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	O5-C1-C2	-5.77	102.36	111.29
4	D	1	8I4	O6B-C6-O6A	-3.05	117.16	124.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	FUC	C6-C5-C4	-2.91	107.75	113.08
4	D	1	8I4	O4-C4-C5	-2.79	103.47	109.80
3	C	1	NAG	C1-O5-C5	2.73	115.85	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	1	8I4	C4-C5-C6-O6A
4	D	1	8I4	C4-C5-C6-O6B


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	2	NDG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	609	-	3,3,3	0.29	0	2,2,2	0.49	0
6	EDO	A	605	-	3,3,3	1.29	1 (33%)	2,2,2	1.85	1 (50%)
5	NAG	A	602	1	14,14,15	1.04	1 (7%)	17,19,21	1.54	3 (17%)
6	EDO	A	606	-	3,3,3	0.20	0	2,2,2	0.35	0
6	EDO	A	604	-	3,3,3	0.49	0	2,2,2	0.45	0
5	NAG	A	601	1	14,14,15	0.92	1 (7%)	17,19,21	1.78	4 (23%)
6	EDO	A	608	-	3,3,3	0.46	0	2,2,2	0.11	0
6	EDO	A	607	-	3,3,3	0.17	0	2,2,2	0.32	0
6	EDO	A	610	-	3,3,3	0.28	0	2,2,2	0.18	0
5	NAG	A	603	1	14,14,15	0.66	0	17,19,21	1.43	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	609	-	-	1/1/1/1	-
6	EDO	A	605	-	-	1/1/1/1	-
5	NAG	A	602	1	-	2/6/23/26	0/1/1/1
6	EDO	A	606	-	-	0/1/1/1	-
6	EDO	A	604	-	-	0/1/1/1	-
5	NAG	A	601	1	-	2/6/23/26	0/1/1/1
6	EDO	A	608	-	-	1/1/1/1	-
6	EDO	A	607	-	-	1/1/1/1	-
6	EDO	A	610	-	-	0/1/1/1	-
5	NAG	A	603	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	602	NAG	C1-C2	2.64	1.55	1.52
5	A	601	NAG	C1-C2	2.39	1.55	1.52
6	A	605	EDO	O1-C1	-2.21	1.30	1.42

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	NAG	C1-O5-C5	3.97	117.50	112.19
5	A	601	NAG	C2-N2-C7	3.71	127.88	122.90
5	A	603	NAG	C1-O5-C5	3.60	117.02	112.19
5	A	602	NAG	O3-C3-C2	3.29	116.24	109.40
5	A	602	NAG	C4-C3-C2	-3.14	106.42	111.02

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	NAG	O5-C5-C6-O6
5	A	601	NAG	C4-C5-C6-O6
5	A	602	NAG	C4-C5-C6-O6
5	A	602	NAG	O5-C5-C6-O6
6	A	608	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	609	EDO	3	0
6	A	605	EDO	1	0
6	A	606	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/385 (100%)	-0.01	6 (1%) 70 76	10, 31, 56, 91	4 (1%)
2	B	74/74 (100%)	-0.03	1 (1%) 73 78	15, 32, 51, 80	1 (1%)
All	All	459/459 (100%)	-0.01	7 (1%) 72 77	10, 31, 56, 91	5 (1%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389[A]	GLY	4.9
1	A	258	PHE	3.7
1	A	543	ILE	3.2
2	B	36	GLN	2.2
1	A	160	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	14/15	0.74	0.14	51,61,82,82	2
3	FUC	C	2	10/11	0.87	0.11	48,52,59,59	3
4	NDG	D	2	14/15	0.96	0.06	23,28,34,39	3
4	8I4	D	1	13/14	0.97	0.05	23,26,27,29	3

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	601	14/15	0.70	0.14	74,87,92,101	3
6	EDO	A	608	4/4	0.73	0.18	51,55,58,61	1
5	NAG	A	602	14/15	0.84	0.11	49,60,66,71	3
6	EDO	A	610	4/4	0.84	0.11	49,57,58,60	1
5	NAG	A	603	14/15	0.85	0.11	56,61,69,79	3
6	EDO	A	606	4/4	0.87	0.13	47,51,54,55	1
6	EDO	A	609	4/4	0.89	0.11	51,52,55,55	1
6	EDO	A	604	4/4	0.95	0.07	32,33,34,34	1
6	EDO	A	607	4/4	0.96	0.07	41,47,49,51	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	EDO	A	605	4/4	0.97	0.06	25,29,31,34	1
7	CL	A	611	1/1	0.98	0.06	41,41,41,41	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.