



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:45 PM UTC

PDB ID : 7PRS / pdb_00007prs
Title : Crystal Structure of the B subunit of heat labile enterotoxin LT-IIc from Escherichia coli in complex with Sialyl-lacto-N-neotetraose d
Authors : Varrot, A.
Deposited on : 2021-09-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	DDD	104	2% 92% 5%
1	EEE	104	4% 90% 5% 5%
1	FFF	104	7% 91% 5%
1	GGG	104	3% 91% 5%
1	HHH	104	2% 90% 5% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	III	104	
1	JJJ	104	
1	KKK	104	
1	LLL	104	
1	MMM	104	
2	DaD	5	
3	GaG	4	
3	JaJ	4	
4	HaH	3	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat-labile enterotoxin IIA, B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	DDD	99	Total	C	N	O	S	0	0	0
			750	474	127	145	4			
1	EEE	99	Total	C	N	O	S	0	0	0
			748	473	126	145	4			
1	FFF	99	Total	C	N	O	S	0	1	0
			745	470	125	146	4			
1	GGG	99	Total	C	N	O	S	0	1	0
			749	475	126	144	4			
1	HHH	99	Total	C	N	O	S	0	0	0
			748	473	126	145	4			
1	III	99	Total	C	N	O	S	0	1	0
			752	477	127	144	4			
1	JJJ	99	Total	C	N	O	S	0	0	0
			758	480	129	145	4			
1	KKK	99	Total	C	N	O	S	0	0	0
			748	473	126	145	4			
1	LLL	99	Total	C	N	O	S	0	0	0
			748	473	126	145	4			
1	MMM	99	Total	C	N	O	S	0	0	0
			750	474	127	145	4			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
DDD	99	HIS	-	expression tag	UNP H6W8F2
DDD	100	HIS	-	expression tag	UNP H6W8F2
DDD	101	HIS	-	expression tag	UNP H6W8F2
DDD	102	HIS	-	expression tag	UNP H6W8F2
DDD	103	HIS	-	expression tag	UNP H6W8F2
DDD	104	HIS	-	expression tag	UNP H6W8F2
EEE	99	HIS	-	expression tag	UNP H6W8F2
EEE	100	HIS	-	expression tag	UNP H6W8F2
EEE	101	HIS	-	expression tag	UNP H6W8F2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
EEE	102	HIS	-	expression tag	UNP H6W8F2
EEE	103	HIS	-	expression tag	UNP H6W8F2
EEE	104	HIS	-	expression tag	UNP H6W8F2
FFF	99	HIS	-	expression tag	UNP H6W8F2
FFF	100	HIS	-	expression tag	UNP H6W8F2
FFF	101	HIS	-	expression tag	UNP H6W8F2
FFF	102	HIS	-	expression tag	UNP H6W8F2
FFF	103	HIS	-	expression tag	UNP H6W8F2
FFF	104	HIS	-	expression tag	UNP H6W8F2
GGG	99	HIS	-	expression tag	UNP H6W8F2
GGG	100	HIS	-	expression tag	UNP H6W8F2
GGG	101	HIS	-	expression tag	UNP H6W8F2
GGG	102	HIS	-	expression tag	UNP H6W8F2
GGG	103	HIS	-	expression tag	UNP H6W8F2
GGG	104	HIS	-	expression tag	UNP H6W8F2
HHH	99	HIS	-	expression tag	UNP H6W8F2
HHH	100	HIS	-	expression tag	UNP H6W8F2
HHH	101	HIS	-	expression tag	UNP H6W8F2
HHH	102	HIS	-	expression tag	UNP H6W8F2
HHH	103	HIS	-	expression tag	UNP H6W8F2
HHH	104	HIS	-	expression tag	UNP H6W8F2
III	99	HIS	-	expression tag	UNP H6W8F2
III	100	HIS	-	expression tag	UNP H6W8F2
III	101	HIS	-	expression tag	UNP H6W8F2
III	102	HIS	-	expression tag	UNP H6W8F2
III	103	HIS	-	expression tag	UNP H6W8F2
III	104	HIS	-	expression tag	UNP H6W8F2
JJJ	99	HIS	-	expression tag	UNP H6W8F2
JJJ	100	HIS	-	expression tag	UNP H6W8F2
JJJ	101	HIS	-	expression tag	UNP H6W8F2
JJJ	102	HIS	-	expression tag	UNP H6W8F2
JJJ	103	HIS	-	expression tag	UNP H6W8F2
JJJ	104	HIS	-	expression tag	UNP H6W8F2
KKK	99	HIS	-	expression tag	UNP H6W8F2
KKK	100	HIS	-	expression tag	UNP H6W8F2
KKK	101	HIS	-	expression tag	UNP H6W8F2
KKK	102	HIS	-	expression tag	UNP H6W8F2
KKK	103	HIS	-	expression tag	UNP H6W8F2
KKK	104	HIS	-	expression tag	UNP H6W8F2
LLL	99	HIS	-	expression tag	UNP H6W8F2
LLL	100	HIS	-	expression tag	UNP H6W8F2
LLL	101	HIS	-	expression tag	UNP H6W8F2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
LLL	102	HIS	-	expression tag	UNP H6W8F2
LLL	103	HIS	-	expression tag	UNP H6W8F2
LLL	104	HIS	-	expression tag	UNP H6W8F2
MMM	99	HIS	-	expression tag	UNP H6W8F2
MMM	100	HIS	-	expression tag	UNP H6W8F2
MMM	101	HIS	-	expression tag	UNP H6W8F2
MMM	102	HIS	-	expression tag	UNP H6W8F2
MMM	103	HIS	-	expression tag	UNP H6W8F2
MMM	104	HIS	-	expression tag	UNP H6W8F2

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	DaD	5	Total	C	N	O	0	0	0
			68	37	2	29			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	GaG	4	Total	C	N	O	0	0	0
			57	31	2	24			
3	JaJ	4	Total	C	N	O	0	0	0
			57	31	2	24			

- Molecule 4 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	HaH	3	Total	C	N	O	0	0	0
			45	25	2	18			

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	DDD	1	Total	O	S	0	0
			5	4	1		
5	EEE	1	Total	O	S	0	0
			5	4	1		
5	FFF	1	Total	O	S	0	0
			5	4	1		
5	HHH	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is N-acetyl-alpha-neuraminic acid (CCD ID: SIA) (formula: C₁₁H₁₉NO₉) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	III	1	Total	C	N	O	0	0
			21	11	1	9		
6	JJJ	1	Total	C	N	O	0	0
			21	11	1	9		
6	KKK	1	Total	C	N	O	0	0
			21	11	1	9		
6	LLL	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	DDD	71	Total	O	0	2
			73	73		
7	EEE	61	Total	O	0	3
			64	64		
7	FFF	45	Total	O	0	1
			46	46		
7	GGG	56	Total	O	0	2
			58	58		
7	HHH	73	Total	O	0	2
			75	75		
7	III	57	Total	O	0	2
			59	59		
7	JJJ	68	Total	O	0	1
			69	69		
7	KKK	69	Total	O	0	1
			70	70		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	LLL	54	Total	O	0	2
			56	56		
7	MMM	62	Total	O	0	4
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heat-labile enterotoxin IIA, B chain




- Molecule 1: Heat-labile enterotoxin IIA, B chain




- Molecule 1: Heat-labile enterotoxin IIA, B chain





- Molecule 1: Heat-labile enterotoxin IIA, B chain




- Molecule 1: Heat-labile enterotoxin IIA, B chain



- Molecule 1: Heat-labile enterotoxin IIA, B chain





- Molecule 1: Heat-labile enterotoxin IIA, B chain



- Molecule 1: Heat-labile enterotoxin IIA, B chain





- Molecule 1: Heat-labile enterotoxin IIA, B chain



- Molecule 1: Heat-labile enterotoxin IIA, B chain



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose

Chain JaJ:  100%

GAL1
NAG2
GAL3
SIA4

- Molecule 4: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain HaH:  100%

NAG1
GAL2
SIA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.09Å 121.27Å 126.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.54 – 2.00 38.54 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.54-2.00) 99.9 (38.54-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.188 , 0.230 0.196 , 0.235	Depositor DCC
R_{free} test set	4010 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.388	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8462	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1140e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GLC, NAG, SO4, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	DDD	1.21	0/766	1.30	1/1043 (0.1%)
1	EEE	1.21	1/764 (0.1%)	1.28	1/1041 (0.1%)
1	FFF	1.18	0/761	1.31	2/1041 (0.2%)
1	GGG	1.20	0/765	1.28	1/1044 (0.1%)
1	HHH	1.22	2/764 (0.3%)	1.29	1/1041 (0.1%)
1	III	1.21	0/768	1.29	0/1047
1	JJJ	1.22	1/774 (0.1%)	1.30	1/1051 (0.1%)
1	KKK	1.20	1/764 (0.1%)	1.27	1/1041 (0.1%)
1	LLL	1.23	1/764 (0.1%)	1.30	1/1041 (0.1%)
1	MMM	1.25	0/766	1.34	0/1043
All	All	1.21	6/7656 (0.1%)	1.30	9/10433 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	HHH	99	HIS	CE1-NE2	7.35	1.39	1.32
1	LLL	39	ILE	C-O	5.63	1.29	1.24
1	JJJ	94	ILE	C-O	5.28	1.29	1.24
1	HHH	21	VAL	C-O	5.27	1.30	1.24
1	KKK	78	VAL	C-O	5.20	1.29	1.23
1	EEE	53	GLN	C-O	-5.05	1.17	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	FFF	99	HIS	CA-C-O	6.08	131.13	120.80
1	JJJ	98	PRO	CB-CA-C	-5.83	102.50	112.64
1	EEE	98	PRO	CB-CA-C	-5.78	103.20	112.21
1	LLL	98	PRO	CB-CA-C	-5.68	103.24	112.62
1	KKK	98	PRO	CB-CA-C	-5.27	103.93	112.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	FFF	98	PRO	CB-CA-C	-5.25	103.96	112.62
1	GGG	98	PRO	CB-CA-C	-5.20	104.04	112.62
1	HHH	98	PRO	CB-CA-C	-5.14	104.14	112.62
1	DDD	89	ASN	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	DDD	750	0	734	2	0
1	EEE	748	0	727	3	0
1	FFF	745	0	710	1	0
1	GGG	749	0	724	3	0
1	HHH	748	0	727	2	0
1	III	752	0	733	3	0
1	JJJ	758	0	756	5	0
1	KKK	748	0	727	4	0
1	LLL	748	0	727	4	0
1	MMM	750	0	734	2	0
2	DaD	68	0	58	0	0
3	GaG	57	0	49	1	0
3	JaJ	57	0	49	0	0
4	HaH	45	0	38	0	0
5	DDD	5	0	0	0	0
5	EEE	5	0	0	0	0
5	FFF	5	0	0	0	0
5	HHH	5	0	0	0	0
6	III	21	0	18	0	0
6	JJJ	21	0	18	0	0
6	KKK	21	0	18	1	0
6	LLL	20	0	17	0	0
7	DDD	73	0	0	0	0
7	EEE	64	0	0	1	0
7	FFF	46	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	GGG	58	0	0	0	0
7	HHH	75	0	0	0	0
7	III	59	0	0	0	0
7	JJJ	69	0	0	1	0
7	KKK	70	0	0	0	0
7	LLL	56	0	0	0	0
7	MMM	66	0	0	0	0
All	All	8462	0	7564	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:KKK:57:ASP:OD1	1:LLL:35:LYS:HE3	1.84	0.76
1:JJJ:76:VAL:HG13	1:JJJ:98:PRO:HA	1.76	0.67
1:EEE:64:MET:CE	1:EEE:91:VAL:HG12	2.27	0.64
1:KKK:64:MET:CE	1:KKK:91:VAL:HG12	2.28	0.62
1:GGG:64:MET:CE	1:GGG:91:VAL:HG12	2.30	0.62
1:HHH:64:MET:CE	1:HHH:91:VAL:HG12	2.30	0.60
1:JJJ:64:MET:CE	1:JJJ:91:VAL:HG12	2.31	0.60
1:GGG:64:MET:HE1	1:GGG:91:VAL:HG12	1.84	0.60
1:GGG:76[B]:VAL:HG13	1:GGG:98:PRO:HA	1.84	0.59
1:MMM:64:MET:CE	1:MMM:91:VAL:HG12	2.31	0.59
1:DDD:64:MET:CE	1:DDD:91:VAL:HG12	2.32	0.59
1:III:64:MET:CE	1:III:91:VAL:HG12	2.32	0.59
1:LLL:64:MET:CE	1:LLL:91:VAL:HG12	2.33	0.59
1:EEE:64:MET:HE1	1:EEE:91:VAL:HG12	1.87	0.56
1:KKK:64:MET:HE1	1:KKK:91:VAL:HG12	1.87	0.56
1:III:64:MET:HE1	1:III:91:VAL:HG12	1.88	0.54
1:MMM:64:MET:HE1	1:MMM:91:VAL:HG12	1.90	0.54
1:III:76[B]:VAL:HG13	1:III:98:PRO:HA	1.92	0.52
1:JJJ:64:MET:HE1	1:JJJ:91:VAL:HG12	1.92	0.52
1:HHH:64:MET:HE1	1:HHH:91:VAL:HG12	1.91	0.51
1:LLL:64:MET:HE1	1:LLL:91:VAL:HG12	1.93	0.51
1:DDD:64:MET:HE1	1:DDD:91:VAL:HG12	1.93	0.50
1:EEE:22:GLN:HG2	7:EEE:353[B]:HOH:O	2.15	0.46
1:FFF:2:VAL:HG21	1:FFF:19:GLN:HG2	1.99	0.43
1:JJJ:2:VAL:HG21	1:JJJ:19:GLN:HG2	2.00	0.43
1:KKK:92:TRP:CH2	6:KKK:201:SIA:H6	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:JJJ:10:CYS:HB3	7:JJJ:323:HOH:O	2.21	0.41
1:LLL:76:VAL:HG13	1:LLL:98:PRO:HA	2.03	0.41
3:GaG:3:GAL:H4	3:GaG:4:SIA:C1	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	DDD	97/104 (93%)	97 (100%)	0	0	100	100
1	EEE	97/104 (93%)	97 (100%)	0	0	100	100
1	FFF	98/104 (94%)	98 (100%)	0	0	100	100
1	GGG	98/104 (94%)	98 (100%)	0	0	100	100
1	HHH	97/104 (93%)	97 (100%)	0	0	100	100
1	III	98/104 (94%)	98 (100%)	0	0	100	100
1	JJJ	97/104 (93%)	97 (100%)	0	0	100	100
1	KKK	97/104 (93%)	97 (100%)	0	0	100	100
1	LLL	97/104 (93%)	97 (100%)	0	0	100	100
1	MMM	97/104 (93%)	97 (100%)	0	0	100	100
All	All	973/1040 (94%)	973 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	DDD	82/89 (92%)	82 (100%)	0	100	100
1	EEE	81/89 (91%)	81 (100%)	0	100	100
1	FFF	79/89 (89%)	79 (100%)	0	100	100
1	GGG	80/89 (90%)	80 (100%)	0	100	100
1	HHH	81/89 (91%)	81 (100%)	0	100	100
1	III	81/89 (91%)	81 (100%)	0	100	100
1	JJJ	84/89 (94%)	84 (100%)	0	100	100
1	KKK	81/89 (91%)	81 (100%)	0	100	100
1	LLL	81/89 (91%)	81 (100%)	0	100	100
1	MMM	82/89 (92%)	82 (100%)	0	100	100
All	All	812/890 (91%)	812 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	DaD	1	2	12,12,12	0.44	0	17,17,17	1.30	2 (11%)
2	GAL	DaD	2	2	11,11,12	0.45	0	15,15,17	1.07	0
2	NAG	DaD	3	2	14,14,15	1.20	1 (7%)	17,19,21	1.37	3 (17%)
2	GAL	DaD	4	2	11,11,12	0.99	0	15,15,17	1.66	4 (26%)
2	SIA	DaD	5	2	20,20,21	0.72	0	21,28,31	1.31	3 (14%)
3	GAL	GaG	1	3	12,12,12	0.81	0	17,17,17	1.36	4 (23%)
3	NAG	GaG	2	3	14,14,15	0.97	0	17,19,21	1.44	3 (17%)
3	GAL	GaG	3	3	11,11,12	0.91	0	15,15,17	1.62	4 (26%)
3	SIA	GaG	4	3	20,20,21	0.64	1 (5%)	21,28,31	1.01	1 (4%)
4	NAG	HaH	1	4	14,14,15	1.45	1 (7%)	17,19,21	1.82	4 (23%)
4	GAL	HaH	2	4	11,11,12	0.83	0	15,15,17	1.87	2 (13%)
4	SIA	HaH	3	4	20,20,21	0.73	1 (5%)	21,28,31	1.38	3 (14%)
3	GAL	JaJ	1	3	12,12,12	0.61	0	17,17,17	1.27	3 (17%)
3	NAG	JaJ	2	3	14,14,15	1.23	1 (7%)	17,19,21	1.61	4 (23%)
3	GAL	JaJ	3	3	11,11,12	0.93	0	15,15,17	1.16	1 (6%)
3	SIA	JaJ	4	3	20,20,21	1.11	2 (10%)	21,28,31	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	DaD	1	2	-	0/2/22/22	0/1/1/1
2	GAL	DaD	2	2	-	1/2/19/22	0/1/1/1
2	NAG	DaD	3	2	-	0/6/23/26	0/1/1/1
2	GAL	DaD	4	2	-	2/2/19/22	0/1/1/1
2	SIA	DaD	5	2	-	4/18/34/38	0/1/1/1
3	GAL	GaG	1	3	-	1/2/22/22	0/1/1/1
3	NAG	GaG	2	3	-	0/6/23/26	0/1/1/1
3	GAL	GaG	3	3	-	2/2/19/22	0/1/1/1
3	SIA	GaG	4	3	-	0/18/34/38	0/1/1/1
4	NAG	HaH	1	4	-	0/6/23/26	0/1/1/1
4	GAL	HaH	2	4	-	2/2/19/22	0/1/1/1
4	SIA	HaH	3	4	-	4/18/34/38	0/1/1/1
3	GAL	JaJ	1	3	-	2/2/22/22	0/1/1/1
3	NAG	JaJ	2	3	-	0/6/23/26	0/1/1/1
3	GAL	JaJ	3	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIA	JaJ	4	3	-	4/18/34/38	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	JaJ	4	SIA	O1B-C1	-3.86	1.18	1.30
2	DaD	3	NAG	C1-C2	3.45	1.57	1.52
4	HaH	1	NAG	C1-C2	2.83	1.56	1.52
3	GaG	4	SIA	O1B-C1	-2.29	1.23	1.30
3	JaJ	4	SIA	C2-C1	2.26	1.55	1.52
4	HaH	3	SIA	O1B-C1	-2.23	1.23	1.30
3	JaJ	2	NAG	C2-N2	2.04	1.49	1.46

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	HaH	2	GAL	C1-O5-C5	5.94	120.14	112.19
4	HaH	1	NAG	O5-C1-C2	-4.64	104.11	111.29
2	DaD	1	GLC	C4-C3-C2	-3.95	103.90	110.83
2	DaD	5	SIA	O1B-C1-C2	3.77	122.51	112.71
4	HaH	1	NAG	C1-O5-C5	3.53	116.92	112.19
4	HaH	3	SIA	O1B-C1-C2	3.46	121.72	112.71
3	JaJ	2	NAG	O4-C4-C5	-3.33	101.13	109.32
3	GaG	2	NAG	C2-N2-C7	-3.23	118.57	122.90
2	DaD	4	GAL	C2-C3-C4	-3.19	105.26	110.86
3	GaG	3	GAL	O3-C3-C2	-3.09	103.76	110.05
3	GaG	1	GAL	C3-C4-C5	3.06	115.77	110.23
2	DaD	3	NAG	C1-O5-C5	2.89	116.06	112.19
2	DaD	3	NAG	O4-C4-C5	-2.85	102.32	109.32
2	DaD	4	GAL	O4-C4-C3	-2.83	103.69	110.38
3	JaJ	1	GAL	C1-O5-C5	2.79	119.05	113.65
3	GaG	2	NAG	C1-C2-N2	2.74	114.75	110.43
3	GaG	3	GAL	C3-C4-C5	2.68	115.09	110.23
4	HaH	1	NAG	O5-C5-C6	2.66	112.84	107.66
4	HaH	3	SIA	O6-C2-C1	2.65	112.72	107.72
3	JaJ	2	NAG	O5-C5-C6	2.60	112.73	107.66
4	HaH	1	NAG	C6-C5-C4	-2.59	106.67	113.02
2	DaD	5	SIA	O1A-C1-C2	-2.53	117.39	122.85
4	HaH	3	SIA	O1A-C1-C2	-2.51	117.43	122.85
3	GaG	1	GAL	O1-C1-C2	2.39	115.91	108.98
3	GaG	4	SIA	C8-C7-C6	-2.28	108.78	113.05
2	DaD	4	GAL	C1-C2-C3	2.27	112.96	109.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	JaJ	2	NAG	C8-C7-N2	2.26	119.87	116.12
3	JaJ	1	GAL	O5-C5-C6	2.25	112.02	106.44
4	HaH	2	GAL	O5-C5-C6	2.22	111.99	107.66
3	JaJ	1	GAL	O5-C1-C2	2.20	114.17	110.30
3	GaG	1	GAL	O5-C1-C2	-2.17	106.48	110.30
2	DaD	3	NAG	C1-C2-N2	-2.14	107.06	110.43
2	DaD	1	GLC	C6-C5-C4	-2.13	107.78	113.02
3	JaJ	2	NAG	O3-C3-C2	2.12	113.80	109.40
3	GaG	3	GAL	O2-C2-C1	2.10	114.02	109.22
3	GaG	1	GAL	O2-C2-C1	2.08	114.05	109.25
2	DaD	4	GAL	O3-C3-C2	-2.05	105.88	110.05
2	DaD	5	SIA	O6-C2-C1	2.03	111.55	107.72
3	GaG	2	NAG	O5-C1-C2	-2.02	108.17	111.29
3	JaJ	3	GAL	C6-C5-C4	2.00	117.94	113.02
3	GaG	3	GAL	O4-C4-C5	-2.00	104.40	109.32

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	HaH	2	GAL	O5-C5-C6-O6
4	HaH	2	GAL	C4-C5-C6-O6
3	JaJ	1	GAL	O5-C5-C6-O6
2	DaD	4	GAL	O5-C5-C6-O6
3	GaG	3	GAL	O5-C5-C6-O6
4	HaH	3	SIA	O7-C7-C8-C9
4	HaH	3	SIA	C6-C7-C8-C9
4	HaH	3	SIA	C6-C7-C8-O8
3	JaJ	1	GAL	C4-C5-C6-O6
2	DaD	4	GAL	C4-C5-C6-O6
3	GaG	3	GAL	C4-C5-C6-O6
4	HaH	3	SIA	O7-C7-C8-O8
2	DaD	5	SIA	C6-C7-C8-O8
3	JaJ	4	SIA	C6-C7-C8-O8
2	DaD	5	SIA	O7-C7-C8-C9
2	DaD	5	SIA	C6-C7-C8-C9
2	DaD	5	SIA	O7-C7-C8-O8
3	JaJ	4	SIA	C6-C7-C8-C9
3	GaG	1	GAL	C4-C5-C6-O6
3	JaJ	4	SIA	O7-C7-C8-C9
3	JaJ	4	SIA	O7-C7-C8-O8
2	DaD	2	GAL	C4-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	GaG	4	SIA	1	0
3	GaG	3	GAL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	DDD	201	-	4,4,4	0.41	0	6,6,6	0.25	0
5	SO4	EEE	201	-	4,4,4	0.40	0	6,6,6	0.16	0
6	SIA	KKK	201	-	21,21,21	0.77	1 (4%)	24,31,31	0.98	1 (4%)
6	SIA	LLL	201	-	20,20,21	0.80	1 (5%)	21,28,31	0.87	1 (4%)
5	SO4	FFF	201	-	4,4,4	0.38	0	6,6,6	0.44	0
6	SIA	III	201	-	21,21,21	0.62	0	24,31,31	0.98	1 (4%)
5	SO4	HHH	201	-	4,4,4	0.46	0	6,6,6	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SIA	JJJ	201	-	21,21,21	0.72	1 (4%)	24,31,31	1.00	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SIA	LLL	201	-	-	8/18/34/38	0/1/1/1
6	SIA	JJJ	201	-	-	8/20/38/38	0/1/1/1
6	SIA	KKK	201	-	-	5/20/38/38	0/1/1/1
6	SIA	III	201	-	-	8/20/38/38	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	LLL	201	SIA	O1B-C1	-2.76	1.21	1.30
6	JJJ	201	SIA	O1B-C1	-2.37	1.21	1.30
6	KKK	201	SIA	O1B-C1	-2.31	1.22	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	III	201	SIA	O1A-C1-C2	-3.83	117.46	123.85
6	KKK	201	SIA	O1A-C1-C2	-2.99	118.87	123.85
6	LLL	201	SIA	O1B-C1-C2	2.61	119.50	112.71
6	JJJ	201	SIA	O1A-C1-C2	-2.50	119.69	123.85
6	JJJ	201	SIA	O2-C2-O6	-2.42	103.42	109.51

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	III	201	SIA	C6-C7-C8-O8
6	III	201	SIA	O7-C7-C8-C9
6	III	201	SIA	O7-C7-C8-O8
6	III	201	SIA	C7-C8-C9-O9
6	III	201	SIA	O8-C8-C9-O9
6	KKK	201	SIA	C6-C7-C8-C9
6	KKK	201	SIA	O7-C7-C8-C9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	LLL	201	SIA	C6-C7-C8-C9
6	LLL	201	SIA	C6-C7-C8-O8
6	LLL	201	SIA	O7-C7-C8-C9
6	LLL	201	SIA	O7-C7-C8-O8
6	KKK	201	SIA	C6-C7-C8-O8
6	III	201	SIA	C6-C7-C8-C9
6	LLL	201	SIA	C11-C10-N5-C5
6	LLL	201	SIA	O10-C10-N5-C5
6	KKK	201	SIA	O7-C7-C8-O8
6	JJJ	201	SIA	C6-C7-C8-O8
6	LLL	201	SIA	C7-C8-C9-O9
6	JJJ	201	SIA	C7-C8-C9-O9
6	LLL	201	SIA	O8-C8-C9-O9
6	JJJ	201	SIA	C6-C7-C8-C9
6	JJJ	201	SIA	O8-C8-C9-O9
6	JJJ	201	SIA	O7-C7-C8-O8
6	KKK	201	SIA	O1B-C1-C2-C3
6	III	201	SIA	O1A-C1-C2-O2
6	JJJ	201	SIA	O1A-C1-C2-O2
6	JJJ	201	SIA	O7-C7-C8-C9
6	III	201	SIA	O1B-C1-C2-C3
6	JJJ	201	SIA	O1B-C1-C2-C3


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	KKK	201	SIA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand SIA KKK 201



Ligand SIA LLL 201







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	DDD	99/104 (95%)	-0.03	2 (2%) 65 64	22, 33, 47, 53	0
1	EEE	99/104 (95%)	0.24	4 (4%) 42 41	24, 37, 56, 60	2 (2%)
1	FFF	99/104 (95%)	0.59	7 (7%) 22 20	20, 42, 59, 71	1 (1%)
1	GGG	99/104 (95%)	0.27	3 (3%) 52 51	20, 40, 57, 63	1 (1%)
1	HHH	99/104 (95%)	0.14	2 (2%) 65 64	23, 34, 48, 56	0
1	III	99/104 (95%)	0.26	5 (5%) 33 32	19, 38, 58, 64	1 (1%)
1	JJJ	99/104 (95%)	0.20	2 (2%) 65 64	25, 36, 52, 67	0
1	KKK	99/104 (95%)	0.08	0 100 100	25, 36, 53, 59	0
1	LLL	99/104 (95%)	0.10	1 (1%) 79 79	23, 35, 54, 61	0
1	MMM	99/104 (95%)	0.12	1 (1%) 79 79	23, 34, 49, 60	0
All	All	990/1040 (95%)	0.20	27 (2%) 56 55	19, 37, 55, 71	5 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	FFF	86	TYR	4.3
1	FFF	11	ALA	3.6
1	MMM	86	TYR	3.5
1	JJJ	99	HIS	3.5
1	III	86	TYR	3.0
1	JJJ	43	ALA	2.9
1	DDD	43	ALA	2.8
1	FFF	43	ALA	2.8
1	GGG	11	ALA	2.7
1	EEE	54	TYR	2.5
1	III	87	THR	2.4
1	EEE	7	LYS	2.4
1	DDD	54	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	GGG	87	THR	2.3
1	III	11	ALA	2.3
1	FFF	20	SER	2.3
1	EEE	86	TYR	2.3
1	III	33	ASP	2.2
1	HHH	43	ALA	2.2
1	LLL	86	TYR	2.2
1	GGG	2	VAL	2.1
1	EEE	85	ALA	2.1
1	FFF	87	THR	2.0
1	FFF	1	GLY	2.0
1	HHH	2	VAL	2.0
1	FFF	42	SER	2.0
1	III	99	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	DaD	1	12/12	-	-	49,52,55,57	12
2	GAL	DaD	2	11/12	-	-	42,54,61,61	11
2	NAG	DaD	3	14/15	-	-	35,42,47,51	0
2	GAL	DaD	4	11/12	-	-	39,50,59,61	0
2	SIA	DaD	5	20/21	-	-	35,42,49,50	0
3	GAL	GaG	1	12/12	-	-	44,69,80,81	0
3	NAG	GaG	2	14/15	-	-	34,44,49,50	0
3	GAL	GaG	3	11/12	-	-	49,55,69,71	0
3	SIA	GaG	4	20/21	-	-	39,44,50,55	0
3	GAL	JaJ	1	12/12	-	-	40,54,64,67	12
3	NAG	JaJ	2	14/15	-	-	33,38,44,47	0
3	GAL	JaJ	3	11/12	-	-	39,43,47,47	0
3	SIA	JaJ	4	20/21	-	-	33,42,45,48	0
4	NAG	HaH	1	14/15	-	-	46,52,58,64	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GAL	HaH	2	11/12	-	-	42,50,62,63	0
4	SIA	HaH	3	20/21	-	-	39,46,52,52	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain DaD:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain GaG:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain JaJ:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SIA	III	201	21/21	0.67	0.17	45,57,62,69	21
5	SO4	FFF	201	5/5	0.73	0.21	31,35,46,48	5
6	SIA	JJJ	201	21/21	0.75	0.15	51,63,67,69	20
6	SIA	LLL	201	20/21	0.81	0.15	47,58,62,68	20
6	SIA	KKK	201	21/21	0.82	0.11	56,63,70,95	0
5	SO4	EEE	201	5/5	0.91	0.13	43,44,48,58	5
5	SO4	DDD	201	5/5	0.92	0.12	27,31,39,44	5
5	SO4	HHH	201	5/5	0.96	0.07	27,29,42,55	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SIA III 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA JJJ 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA LLL 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SIA KKK 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.