



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:12 AM UTC

PDB ID : 2PSO / pdb_00002pso
Title : Human StarD13 (DLC2) lipid transfer and protein localization domain
Authors : Lehtio, L.; Busam, R.; Arrowsmith, C.H.; Berglund, H.; Collins, R.; Dahlgren, L.G.; Edwards, A.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hallberg, B.M.; Holmberg-Schiavone, L.; Johansson, I.; Kallas, A.; Karlberg, T.; Kotenyova, T.; Moche, M.; Nordlund, P.; Nyman, T.; Ogg, D.; Sagemark, J.; Stenmark, P.; Sundstrom, M.; Thorsell, A.G.; Van den Berg, S.; Weigelt, J.; Welin, M.; Persson, C.; Structural Genomics Consortium (SGC)
Deposited on : 2007-05-07
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	 2% 67% 14% 17%
1	B	237	 2% 65% 14% 16%
1	C	237	 11% 60% 17% 21%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 4477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called StAR-related lipid transfer protein 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	197	1517	962	268	279	8	0	0	0
1	B	198	1535	979	267	282	7	0	0	0
1	C	188	1425	911	250	258	6	0	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	877	MET	-	cloning artifact	UNP Q9Y3M8
A	878	HIS	-	cloning artifact	UNP Q9Y3M8
A	879	HIS	-	cloning artifact	UNP Q9Y3M8
A	880	HIS	-	cloning artifact	UNP Q9Y3M8
A	881	HIS	-	cloning artifact	UNP Q9Y3M8
A	882	HIS	-	cloning artifact	UNP Q9Y3M8
A	883	HIS	-	cloning artifact	UNP Q9Y3M8
A	884	SER	-	cloning artifact	UNP Q9Y3M8
A	885	SER	-	cloning artifact	UNP Q9Y3M8
A	886	GLY	-	cloning artifact	UNP Q9Y3M8
A	887	VAL	-	cloning artifact	UNP Q9Y3M8
A	888	ASP	-	cloning artifact	UNP Q9Y3M8
A	889	LEU	-	cloning artifact	UNP Q9Y3M8
A	890	GLY	-	cloning artifact	UNP Q9Y3M8
A	891	THR	-	cloning artifact	UNP Q9Y3M8
A	892	GLU	-	cloning artifact	UNP Q9Y3M8
A	893	ASN	-	cloning artifact	UNP Q9Y3M8
A	894	LEU	-	cloning artifact	UNP Q9Y3M8
A	895	TYR	-	cloning artifact	UNP Q9Y3M8
A	896	PHE	-	cloning artifact	UNP Q9Y3M8
A	897	GLN	-	cloning artifact	UNP Q9Y3M8
A	898	SER	-	cloning artifact	UNP Q9Y3M8
A	899	MET	-	cloning artifact	UNP Q9Y3M8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	900	GLU	-	cloning artifact	UNP Q9Y3M8
A	901	GLU	-	cloning artifact	UNP Q9Y3M8
A	902	SER	-	cloning artifact	UNP Q9Y3M8
B	877	MET	-	cloning artifact	UNP Q9Y3M8
B	878	HIS	-	cloning artifact	UNP Q9Y3M8
B	879	HIS	-	cloning artifact	UNP Q9Y3M8
B	880	HIS	-	cloning artifact	UNP Q9Y3M8
B	881	HIS	-	cloning artifact	UNP Q9Y3M8
B	882	HIS	-	cloning artifact	UNP Q9Y3M8
B	883	HIS	-	cloning artifact	UNP Q9Y3M8
B	884	SER	-	cloning artifact	UNP Q9Y3M8
B	885	SER	-	cloning artifact	UNP Q9Y3M8
B	886	GLY	-	cloning artifact	UNP Q9Y3M8
B	887	VAL	-	cloning artifact	UNP Q9Y3M8
B	888	ASP	-	cloning artifact	UNP Q9Y3M8
B	889	LEU	-	cloning artifact	UNP Q9Y3M8
B	890	GLY	-	cloning artifact	UNP Q9Y3M8
B	891	THR	-	cloning artifact	UNP Q9Y3M8
B	892	GLU	-	cloning artifact	UNP Q9Y3M8
B	893	ASN	-	cloning artifact	UNP Q9Y3M8
B	894	LEU	-	cloning artifact	UNP Q9Y3M8
B	895	TYR	-	cloning artifact	UNP Q9Y3M8
B	896	PHE	-	cloning artifact	UNP Q9Y3M8
B	897	GLN	-	cloning artifact	UNP Q9Y3M8
B	898	SER	-	cloning artifact	UNP Q9Y3M8
B	899	MET	-	cloning artifact	UNP Q9Y3M8
B	900	GLU	-	cloning artifact	UNP Q9Y3M8
B	901	GLU	-	cloning artifact	UNP Q9Y3M8
B	902	SER	-	cloning artifact	UNP Q9Y3M8
C	877	MET	-	cloning artifact	UNP Q9Y3M8
C	878	HIS	-	cloning artifact	UNP Q9Y3M8
C	879	HIS	-	cloning artifact	UNP Q9Y3M8
C	880	HIS	-	cloning artifact	UNP Q9Y3M8
C	881	HIS	-	cloning artifact	UNP Q9Y3M8
C	882	HIS	-	cloning artifact	UNP Q9Y3M8
C	883	HIS	-	cloning artifact	UNP Q9Y3M8
C	884	SER	-	cloning artifact	UNP Q9Y3M8
C	885	SER	-	cloning artifact	UNP Q9Y3M8
C	886	GLY	-	cloning artifact	UNP Q9Y3M8
C	887	VAL	-	cloning artifact	UNP Q9Y3M8
C	888	ASP	-	cloning artifact	UNP Q9Y3M8
C	889	LEU	-	cloning artifact	UNP Q9Y3M8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	890	GLY	-	cloning artifact	UNP Q9Y3M8
C	891	THR	-	cloning artifact	UNP Q9Y3M8
C	892	GLU	-	cloning artifact	UNP Q9Y3M8
C	893	ASN	-	cloning artifact	UNP Q9Y3M8
C	894	LEU	-	cloning artifact	UNP Q9Y3M8
C	895	TYR	-	cloning artifact	UNP Q9Y3M8
C	896	PHE	-	cloning artifact	UNP Q9Y3M8
C	897	GLN	-	cloning artifact	UNP Q9Y3M8
C	898	SER	-	cloning artifact	UNP Q9Y3M8
C	899	MET	-	cloning artifact	UNP Q9Y3M8
C	900	GLU	-	cloning artifact	UNP Q9Y3M8
C	901	GLU	-	cloning artifact	UNP Q9Y3M8
C	902	SER	-	cloning artifact	UNP Q9Y3M8

4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	78.24Å 78.24Å 212.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.79 – 2.80 19.79 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.79-2.80) 99.5 (19.79-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.79Å)	Xtrriage
Refinement program	REFMAC 5.3.0032	Depositor
R, R_{free}	0.210 , 0.244 0.206 , 0.242	Depositor DCC
R_{free} test set	1571 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	75.1	Xtrriage
Anisotropy	0.248	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.055 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4477	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/1554	1.09	4/2118 (0.2%)
1	B	0.76	0/1573	1.10	6/2143 (0.3%)
1	C	0.57	0/1458	1.01	8/1990 (0.4%)
All	All	0.70	0/4585	1.07	18/6251 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	982	VAL	N-CA-C	7.41	118.18	111.81
1	C	948	ASN	CA-C-N	7.09	126.77	119.82
1	C	948	ASN	C-N-CA	7.09	126.77	119.82
1	B	913	LEU	N-CA-C	-6.50	105.35	113.28
1	C	1022	LEU	CA-C-N	6.40	127.84	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	945	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1517	0	1440	13	0
1	B	1535	0	1471	18	0
1	C	1425	0	1349	15	0
All	All	4477	0	4260	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1005:ALA:HB1	1:B:1006:PRO:HD3	1.50	0.90
1:B:957:VAL:HG13	1:B:1094:VAL:HG13	1.68	0.74
1:B:1005:ALA:HB1	1:B:1006:PRO:CD	2.18	0.73
1:B:1005:ALA:CB	1:B:1006:PRO:CD	2.72	0.67
1:A:983:GLN:O	1:A:999:TYR:HA	1.95	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	195/237 (82%)	170 (87%)	21 (11%)	4 (2%)	5 20
1	B	196/237 (83%)	177 (90%)	17 (9%)	2 (1%)	12 38
1	C	184/237 (78%)	162 (88%)	21 (11%)	1 (0%)	24 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	575/711 (81%)	509 (88%)	59 (10%)	7 (1%)	10	34

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1005	ALA
1	A	1041	LEU
1	A	1103	PRO
1	C	925	PHE
1	B	1076	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/209 (76%)	138 (87%)	20 (13%)	4	15
1	B	161/209 (77%)	139 (86%)	22 (14%)	3	13
1	C	145/209 (69%)	128 (88%)	17 (12%)	5	18
All	All	464/627 (74%)	405 (87%)	59 (13%)	4	15

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	976	LEU
1	C	1075	LYS
1	B	1030	VAL
1	C	1074	LEU
1	C	1014	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1040	GLN
1	C	1002	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	936	ASN
1	B	912	HIS
1	B	993	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/237 (83%)	-0.30	4 (2%) 65 56	7, 37, 82, 100	0
1	B	198/237 (83%)	-0.26	4 (2%) 65 56	12, 34, 71, 88	0
1	C	188/237 (79%)	0.72	26 (13%) 6 5	17, 65, 93, 108	0
All	All	583/711 (81%)	0.04	34 (5%) 29 22	7, 44, 88, 108	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	909	TYR	10.6
1	C	910	LEU	8.5
1	C	1024	LYS	5.1
1	C	951	LYS	4.4
1	C	911	ASN	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.