



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:35 AM UTC

PDB ID : 1PT7 / pdb_00001pt7
Title : Crystal structure of the apo-form of the yfdW gene product of E. coli
Authors : Gruez, A.; Roig-Zamboni, V.; Valencia, C.; Campanacci, V.; Cambillau, C.
Deposited on : 2003-06-23
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	 2% 82% 12% • 5%
1	B	437	 % 78% 16% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	602	-	X	-	-
2	PO4	B	603	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7024 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein yfdW.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	415	Total	C	N	O	S	51	0	0
			3216	2040	549	610	17			
1	B	415	Total	C	N	O	S	27	0	0
			3216	2040	549	610	17			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	SER	-	cloning artifact	UNP P69902
A	-19	TYR	-	cloning artifact	UNP P69902
A	-18	TYR	-	cloning artifact	UNP P69902
A	-17	HIS	-	expression tag	UNP P69902
A	-16	HIS	-	expression tag	UNP P69902
A	-15	HIS	-	expression tag	UNP P69902
A	-14	HIS	-	expression tag	UNP P69902
A	-13	HIS	-	expression tag	UNP P69902
A	-12	HIS	-	expression tag	UNP P69902
A	-11	LEU	-	cloning artifact	UNP P69902
A	-10	GLU	-	cloning artifact	UNP P69902
A	-9	SER	-	cloning artifact	UNP P69902
A	-8	THR	-	cloning artifact	UNP P69902
A	-7	SER	-	cloning artifact	UNP P69902
A	-6	LEU	-	cloning artifact	UNP P69902
A	-5	TYR	-	cloning artifact	UNP P69902
A	-4	LYS	-	cloning artifact	UNP P69902
A	-3	LYS	-	cloning artifact	UNP P69902
A	-2	ALA	-	cloning artifact	UNP P69902
A	-1	GLY	-	cloning artifact	UNP P69902
A	0	LEU	-	cloning artifact	UNP P69902
B	-20	SER	-	cloning artifact	UNP P69902
B	-19	TYR	-	cloning artifact	UNP P69902
B	-18	TYR	-	cloning artifact	UNP P69902
B	-17	HIS	-	expression tag	UNP P69902

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP P69902
B	-15	HIS	-	expression tag	UNP P69902
B	-14	HIS	-	expression tag	UNP P69902
B	-13	HIS	-	expression tag	UNP P69902
B	-12	HIS	-	expression tag	UNP P69902
B	-11	LEU	-	cloning artifact	UNP P69902
B	-10	GLU	-	cloning artifact	UNP P69902
B	-9	SER	-	cloning artifact	UNP P69902
B	-8	THR	-	cloning artifact	UNP P69902
B	-7	SER	-	cloning artifact	UNP P69902
B	-6	LEU	-	cloning artifact	UNP P69902
B	-5	TYR	-	cloning artifact	UNP P69902
B	-4	LYS	-	cloning artifact	UNP P69902
B	-3	LYS	-	cloning artifact	UNP P69902
B	-2	ALA	-	cloning artifact	UNP P69902
B	-1	GLY	-	cloning artifact	UNP P69902
B	0	LEU	-	cloning artifact	UNP P69902

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	279	Total	O	0	0
			279	279		
4	B	261	Total	O	0	0
			261	261		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hypothetical protein yfdW



• Molecule 1: Hypothetical protein yfdW



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.92Å 118.63Å 136.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80 20.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	89.5 (20.00-1.80) 89.4 (20.00-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.07 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.159 , 0.191 0.171 , 0.202	Depositor DCC
R_{free} test set	2468 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.708	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7024	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2371e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	6/3292 (0.2%)	1.16	9/4469 (0.2%)
1	B	1.18	2/3292 (0.1%)	1.09	10/4469 (0.2%)
All	All	1.20	8/6584 (0.1%)	1.13	19/8938 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	114	GLU	CG-CD	-15.26	1.13	1.52
1	B	195	ARG	CD-NE	-8.96	1.33	1.46
1	A	149	ALA	C-O	-6.50	1.16	1.24
1	A	102	ILE	C-N	6.47	1.42	1.33
1	B	263	ASP	C-O	-6.14	1.20	1.25
1	A	36	ILE	CA-CB	-5.84	1.47	1.54
1	A	410	MET	CG-SD	5.07	1.93	1.80
1	A	57	ALA	CA-CB	5.06	1.61	1.53

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	GLU	CB-CG-CD	32.52	167.89	112.60
1	B	134	VAL	CB-CA-C	-12.16	96.11	112.04
1	A	114	GLU	CG-CD-OE2	-9.52	96.50	118.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	225	GLU	CB-CG-CD	9.51	128.77	112.60
1	A	114	GLU	CG-CD-OE1	7.58	135.83	118.40
1	B	413	ASN	CB-CA-C	-6.14	99.27	110.63
1	B	134	VAL	N-CA-C	6.08	117.56	110.62
1	B	114	GLU	CB-CG-CD	5.66	122.22	112.60
1	A	38	ARG	CB-CG-CD	5.65	124.30	111.30
1	B	412	GLN	CA-C-N	5.53	128.25	120.28
1	B	412	GLN	C-N-CA	5.53	128.25	120.28
1	A	192	ARG	CG-CD-NE	5.46	124.02	112.00
1	B	80	LYS	CB-CG-CD	5.42	123.76	111.30
1	B	388	LEU	N-CA-C	-5.37	103.30	110.55
1	A	193	GLY	N-CA-C	-5.19	104.66	112.60
1	B	109	TRP	N-CA-C	-5.19	105.70	111.36
1	B	137	LYS	CG-CD-CE	5.04	122.89	111.30
1	A	102	ILE	CA-C-N	5.04	127.34	120.54
1	A	102	ILE	C-N-CA	5.04	127.34	120.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	GLU	Sidechain
1	A	192	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3216	0	3159	41	0
1	B	3216	0	3159	56	0
2	A	20	0	0	1	0
2	B	20	0	0	1	0
3	A	6	0	7	0	0
3	B	6	0	8	0	0
4	A	279	0	0	8	1
4	B	261	0	0	3	1
All	All	7024	0	6333	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:MET:HE1	1:B:105:MET:SD	1.89	1.12
1:A:137:LYS:HG2	4:A:972:HOH:O	1.58	1.03
1:B:74:THR:HG21	1:B:83:MET:HE3	1.41	1.01
1:A:175:HIS:HE1	1:A:206:ASN:HD22	1.19	0.89
1:B:13:PHE:CZ	1:B:83:MET:HE2	2.11	0.86
1:B:229:GLN:HE21	1:B:229:GLN:H	1.26	0.82
1:B:323:GLU:OE1	4:B:836:HOH:O	2.01	0.78
1:B:66:ASN:HD21	1:B:384:LYS:H	1.32	0.78
1:A:368:LEU:H	1:B:216:GLN:HE22	1.33	0.76
1:A:276:ASN:HD21	1:A:279:ASN:HD22	1.31	0.76
1:A:216:GLN:HE22	1:B:368:LEU:H	1.31	0.75
2:A:602:PO4:O4	2:A:605:PO4:O1	2.07	0.72
1:A:273:GLN:HE22	1:A:275:GLN:NE2	1.86	0.72
1:A:188:GLU:OE1	1:B:188:GLU:OE2	2.08	0.72
1:B:319:ILE:CD1	1:B:324:ALA:HB2	2.19	0.71
1:A:73:ASN:C	1:A:73:ASN:HD22	1.98	0.71
1:A:154:GLY:HA2	1:B:161:LEU:HD21	1.73	0.71
1:A:373:PRO:HB2	1:B:198:MET:HE1	1.74	0.70
1:A:276:ASN:ND2	1:A:279:ASN:HD22	1.90	0.69
1:B:319:ILE:HD13	1:B:324:ALA:HB2	1.75	0.69
1:B:13:PHE:HZ	1:B:83:MET:HE2	1.56	0.67
1:B:286:LYS:NZ	1:B:311:GLU:OE1	2.27	0.67
1:B:66:ASN:ND2	1:B:384:LYS:H	1.93	0.65
1:A:274:GLU:CD	4:A:856:HOH:O	2.39	0.64
1:B:229:GLN:H	1:B:229:GLN:NE2	1.93	0.64
1:A:216:GLN:NE2	1:B:368:LEU:H	1.96	0.63
1:B:38:ARG:HB3	1:B:41:VAL:HG13	1.79	0.63
1:A:175:HIS:CE1	1:A:206:ASN:HD22	2.10	0.61
1:B:74:THR:CG2	1:B:83:MET:HE3	2.25	0.60
1:A:69:SER:H	1:A:392:HIS:HD2	1.49	0.60
1:A:175:HIS:HE1	1:A:206:ASN:ND2	1.96	0.60
1:B:69:SER:H	1:B:392:HIS:HD2	1.48	0.59
1:A:368:LEU:H	1:B:216:GLN:NE2	2.01	0.57
1:B:69:SER:H	1:B:392:HIS:CD2	2.23	0.56
1:A:11:LEU:HD21	4:A:844:HOH:O	2.05	0.56
1:B:273:GLN:HE22	1:B:275:GLN:HB2	1.71	0.56
1:B:70:ILE:HG13	1:B:393:THR:HG23	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:MET:HA	1:A:83:MET:HE2	1.88	0.55
1:A:70:ILE:HG22	1:A:393:THR:HA	1.88	0.54
1:B:360:GLU:OE1	1:B:366:LYS:NZ	2.40	0.54
1:B:15:GLY:H	1:B:96:ASN:HD21	1.56	0.54
1:B:78:GLU:OE1	1:B:413:ASN:ND2	2.41	0.54
1:B:206:ASN:ND2	1:B:209:ARG:HH11	2.06	0.54
1:B:65:SER:HB3	1:B:66:ASN:HD22	1.74	0.52
1:A:69:SER:H	1:A:392:HIS:CD2	2.27	0.52
1:B:175:HIS:HE1	1:B:206:ASN:OD1	1.93	0.51
1:A:215:GLN:HE21	1:B:341:MET:HB2	1.76	0.50
1:B:11:LEU:HG	1:B:90:ALA:HB2	1.93	0.50
1:B:11:LEU:HD11	1:B:86:LEU:C	2.37	0.49
1:B:215:GLN:HE22	1:B:238:ALA:HA	1.76	0.49
1:B:260:TRP:HA	1:B:266:ALA:HB3	1.94	0.49
2:B:603:PO4:O2	2:B:606:PO4:O1	2.31	0.47
1:A:137:LYS:NZ	4:A:972:HOH:O	2.37	0.47
1:A:341:MET:HB2	1:B:215:GLN:HE21	1.80	0.47
1:A:167:LEU:HD21	1:B:167:LEU:HD21	1.97	0.47
1:B:100:GLY:O	1:B:104:HIS:ND1	2.49	0.46
1:A:13:PHE:HZ	1:A:83:MET:SD	2.39	0.46
1:B:11:LEU:HG	1:B:90:ALA:CB	2.46	0.45
1:A:9:LYS:HD2	1:A:89:GLU:O	2.17	0.45
1:B:206:ASN:HD21	1:B:209:ARG:HH11	1.63	0.45
1:A:89:GLU:HG3	4:A:844:HOH:O	2.17	0.45
1:B:273:GLN:NE2	1:B:275:GLN:H	2.15	0.45
1:A:137:LYS:CE	4:A:972:HOH:O	2.64	0.44
1:B:37:GLU:HB3	1:B:42:GLY:HA2	1.99	0.44
1:A:215:GLN:HE22	1:A:238:ALA:HA	1.83	0.44
1:B:392:HIS:HE1	4:B:951:HOH:O	2.00	0.44
1:B:195:ARG:HD3	4:B:955:HOH:O	2.17	0.44
1:B:93:LEU:C	1:B:93:LEU:HD23	2.43	0.43
1:B:210:VAL:O	1:B:213:ARG:HB3	2.17	0.43
1:A:73:ASN:HD21	1:A:75:LYS:HB2	1.83	0.43
1:A:35:LYS:HE2	1:A:60:PHE:HE2	1.84	0.42
1:A:11:LEU:HG	1:A:90:ALA:HB2	2.01	0.42
1:B:215:GLN:HE22	1:B:239:VAL:H	1.67	0.42
1:B:124:ILE:HG13	1:B:176:LEU:HD22	2.00	0.42
1:A:299:HIS:ND1	1:A:299:HIS:C	2.77	0.42
1:A:68:ARG:HA	1:A:392:HIS:CD2	2.54	0.41
1:A:215:GLN:HE22	1:A:239:VAL:H	1.67	0.41
1:B:103:ASP:OD1	1:B:108:THR:HA	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:LYS:HE2	1:B:260:TRP:CZ3	2.55	0.41
1:B:17:GLN:HG3	1:B:63:LEU:HD11	2.01	0.41
1:B:68:ARG:HA	1:B:392:HIS:CD2	2.55	0.41
1:A:89:GLU:CG	4:A:844:HOH:O	2.68	0.41
1:A:188:GLU:CD	1:B:188:GLU:OE2	2.63	0.41
1:B:257:CYS:SG	1:B:268:ILE:HG23	2.61	0.41
1:A:124:ILE:HG13	1:A:176:LEU:HD22	2.03	0.40
1:A:230:TYR:HA	1:A:231:PRO:HA	1.93	0.40
1:A:330:GLN:NE2	4:A:886:HOH:O	2.54	0.40
1:B:84:GLU:OE1	1:B:88:ARG:NH2	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:947:HOH:O	4:B:959:HOH:O[2_554]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/437 (94%)	406 (98%)	6 (2%)	1 (0%)	43	31
1	B	413/437 (94%)	407 (98%)	5 (1%)	1 (0%)	43	31
All	All	826/874 (94%)	813 (98%)	11 (1%)	2 (0%)	43	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	VAL
1	B	16	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/361 (94%)	333 (98%)	8 (2%)	44	33
1	B	341/361 (94%)	334 (98%)	7 (2%)	47	36
All	All	682/722 (94%)	667 (98%)	15 (2%)	45	34

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	114	GLU
1	A	129	GLU
1	A	137	LYS
1	A	161	LEU
1	A	288	GLU
1	A	389	LEU
1	A	411	LYS
1	B	41	VAL
1	B	75	LYS
1	B	96	ASN
1	B	134	VAL
1	B	225	GLU
1	B	229	GLN
1	B	319	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	175	HIS
1	A	215	GLN
1	A	216	GLN
1	A	275	GLN
1	A	276	ASN
1	A	330	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	GLN
1	A	392	HIS
1	A	398	GLN
1	B	64	ASN
1	B	66	ASN
1	B	73	ASN
1	B	96	ASN
1	B	111	HIS
1	B	135	ASN
1	B	175	HIS
1	B	206	ASN
1	B	215	GLN
1	B	216	GLN
1	B	229	GLN
1	B	273	GLN
1	B	299	HIS
1	B	392	HIS
1	B	398	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	702	-	5,5,5	0.65	0	5,5,5	1.14	0
2	PO4	A	602	-	4,4,4	1.76	1 (25%)	6,6,6	2.54	3 (50%)
2	PO4	B	606	-	4,4,4	1.05	0	6,6,6	0.81	0
2	PO4	B	603	-	4,4,4	1.71	1 (25%)	6,6,6	2.61	3 (50%)
2	PO4	A	605	-	4,4,4	0.85	0	6,6,6	1.00	0
2	PO4	A	607	-	4,4,4	0.46	0	6,6,6	0.94	0
2	PO4	B	604	-	4,4,4	1.53	1 (25%)	6,6,6	1.76	2 (33%)
2	PO4	A	601	-	4,4,4	1.22	1 (25%)	6,6,6	1.68	2 (33%)
2	PO4	B	608	-	4,4,4	0.77	0	6,6,6	1.25	1 (16%)
3	GOL	A	701	-	5,5,5	1.04	0	5,5,5	1.19	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	701	-	-	0/4/4/4	-
3	GOL	B	702	-	-	0/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	602	PO4	P-O4	-3.00	1.45	1.54
2	B	603	PO4	P-O1	2.71	1.56	1.50
2	B	604	PO4	P-O2	2.28	1.61	1.54
2	A	601	PO4	P-O1	2.24	1.55	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	603	PO4	O4-P-O3	5.02	123.54	107.91
2	A	602	PO4	O4-P-O3	4.88	123.09	107.91
2	B	603	PO4	O2-P-O1	-2.72	101.33	110.95
2	A	601	PO4	O4-P-O1	-2.52	102.03	110.95
2	B	604	PO4	O3-P-O2	2.49	115.66	107.91
2	A	602	PO4	O3-P-O2	-2.46	100.26	107.91
2	A	601	PO4	O2-P-O1	2.45	119.60	110.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	602	PO4	O2-P-O1	-2.42	102.41	110.95
2	B	608	PO4	O4-P-O3	2.33	115.16	107.91
2	B	603	PO4	O4-P-O1	-2.30	102.81	110.95
3	A	701	GOL	C3-C2-C1	-2.24	103.59	111.80
2	B	604	PO4	O4-P-O2	2.08	114.38	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	602	PO4	1	0
2	B	606	PO4	1	0
2	B	603	PO4	1	0
2	A	605	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	415/437 (94%)	-0.22	7 (1%) 69 69	11, 19, 31, 39	24 (5%)
1	B	415/437 (94%)	-0.23	6 (1%) 73 73	10, 19, 32, 41	22 (5%)
All	All	830/874 (94%)	-0.23	13 (1%) 70 71	10, 19, 32, 41	46 (5%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	102	ILE	5.3
1	A	105	MET	3.8
1	A	99	PRO	3.2
1	A	104	HIS	2.7
1	A	103	ASP	2.6
1	B	98	HIS	2.5
1	B	104	HIS	2.4
1	B	83	MET	2.4
1	A	98	HIS	2.3
1	B	135	ASN	2.2
1	A	74	THR	2.2
1	B	414	HIS	2.1
1	B	137	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	B	608	5/5	0.85	0.15	18,23,30,31	5
3	GOL	B	702	6/6	0.88	0.11	26,32,34,38	0
2	PO4	A	607	5/5	0.91	0.12	24,29,33,36	5
2	PO4	B	606	5/5	0.95	0.10	44,45,48,50	0
2	PO4	A	605	5/5	0.95	0.09	39,39,41,43	0
3	GOL	A	701	6/6	0.95	0.07	23,27,35,38	0
2	PO4	A	601	5/5	0.95	0.13	22,28,32,37	0
2	PO4	B	604	5/5	0.96	0.10	17,28,28,29	0
2	PO4	B	603	5/5	0.99	0.06	21,23,26,28	0
2	PO4	A	602	5/5	0.99	0.05	22,23,24,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.