



wwPDB EM Validation Summary Report ⓘ

Mar 27, 2026 – 11:00 AM UTC

PDB ID : 8PTZ / pdb_00008ptz
EMDB ID : EMD-17926
Title : Cryo-EM structure of human Elp123 in complex with tRNA, S-ethyl-CoA, 5'-deoxyadenosine and methionine
Authors : Abbassi, N.; Jaciuk, M.; Lin, T.-Y.; Glatt, S.
Deposited on : 2023-07-16
Resolution : 3.35 Å(reported)
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMD archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	14390 (2.85 - 3.85)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1332	
2	B	826	
3	C	581	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	X	75	44% 35% 43% 23%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 17194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongator complex protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	685	Total	C	N	O	S	0	0
			5357	3396	915	1017	29		

- Molecule 2 is a protein called Elongator complex protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	741	Total	C	N	O	S	0	0
			5861	3732	1016	1074	39		

- Molecule 3 is a protein called Elongator complex protein 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	538	Total	C	N	O	S	0	0
			4296	2726	763	780	27		

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	548	GLY	-	expression tag	UNP Q9H9T3
C	549	LEU	-	expression tag	UNP Q9H9T3
C	550	GLU	-	expression tag	UNP Q9H9T3
C	551	GLY	-	expression tag	UNP Q9H9T3
C	552	SER	-	expression tag	UNP Q9H9T3
C	553	ALA	-	expression tag	UNP Q9H9T3
C	554	TRP	-	expression tag	UNP Q9H9T3
C	555	SER	-	expression tag	UNP Q9H9T3
C	556	HIS	-	expression tag	UNP Q9H9T3
C	557	PRO	-	expression tag	UNP Q9H9T3
C	558	GLN	-	expression tag	UNP Q9H9T3
C	559	PHE	-	expression tag	UNP Q9H9T3
C	560	GLU	-	expression tag	UNP Q9H9T3
C	561	LYS	-	expression tag	UNP Q9H9T3
C	562	GLY	-	expression tag	UNP Q9H9T3

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	563	GLY	-	expression tag	UNP Q9H9T3
C	564	GLY	-	expression tag	UNP Q9H9T3
C	565	SER	-	expression tag	UNP Q9H9T3
C	566	GLY	-	expression tag	UNP Q9H9T3
C	567	GLY	-	expression tag	UNP Q9H9T3
C	568	GLY	-	expression tag	UNP Q9H9T3
C	569	SER	-	expression tag	UNP Q9H9T3
C	570	GLY	-	expression tag	UNP Q9H9T3
C	571	GLY	-	expression tag	UNP Q9H9T3
C	572	SER	-	expression tag	UNP Q9H9T3
C	573	ALA	-	expression tag	UNP Q9H9T3
C	574	TRP	-	expression tag	UNP Q9H9T3
C	575	SER	-	expression tag	UNP Q9H9T3
C	576	HIS	-	expression tag	UNP Q9H9T3
C	577	PRO	-	expression tag	UNP Q9H9T3
C	578	GLN	-	expression tag	UNP Q9H9T3
C	579	PHE	-	expression tag	UNP Q9H9T3
C	580	GLU	-	expression tag	UNP Q9H9T3
C	581	LYS	-	expression tag	UNP Q9H9T3

- Molecule 4 is a RNA chain called tRNA Gln.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	X	75	Total	C	N	O	P	0	0
			1593	711	278	529	75		

- Molecule 5 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
5	C	1	Total	Fe	S	0
			8	4	4	

- Molecule 6 is 5'-DEOXYADENOSINE (CCD ID: 5AD) (formula: $C_{10}H_{13}N_5O_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
6	C	1	Total	C	N	O	0
			18	10	5	3	

- Molecule 7 is S-Ethyl-CoA (CCD ID: A2U) (formula: $C_{23}H_{40}N_7O_{16}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
			Total	C	N	O	P	S	
7	C	1	50	23	7	16	3	1	0

- Molecule 8 is METHIONINE (CCD ID: MET) (formula: $C_5H_{11}NO_2S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	S	
8	C	1	9	5	1	2	1	0

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
9	X	2	Total	Mg	0
			2	2	

ILE	VAL	SER	ALA	LEU	LEU	ILE
MET	THR	GLU	GLN	VAL	GLN	ASP
ALA	HIS	MET	LYS	GLU	LYS	ASP
SER	ILE	SER	ASN	GLN	GLU	TYR
TYR	LEU	GLY	TYR	ARG	HIS	LEU
GLN	LYS	LYS	MET	LYS	MET	LYS
GLN	VAL	TYR	ALA	HIS	TYR	ARG
GLN	LEU	SER	PHE	ILE	GLU	TYR
LYS	PHE	HIS	LEU	ASP	PRO	GLU
THR	LEU	SER	ASP	ALA	ALA	LYS
SER	PHE	ASN	SER	ALA	GLY	ALA
VAL	GLU	SER	GLN	MET	LEU	ILE
PRO	PHE	ARG	THR	VAL	MET	GLY
VAL	ASP	ILE	ALA	LEU	PHE	HIS
LEU	GLU	SER	THR	GLU	ALA	LEU
ASP	GLN	ALA	PHE	GLU	GLU	SER
ALA	GLY	ARG	SER	CYS	CYS	LYS
GLU	ARG	SER	ARG	ALA	GLY	CYS
LEU	GLU	SER	HIS	GLN	ALA	GLY
PHE	LEU	LYS	LYS	ASP	HIS	PRO
ILE	GLN	ASN	LYS	TYR	GLU	GLU
PRO	LYS	ARG	ARG	GLU	LYS	TYR
PRO	ALA	ARG	LEU	GLU	ALA	PHE
LYS	PHE	LYS	LEU	ALA	LEU	PRO
ILE	GLU	ALA	VAL	VAL	SER	GLU
ASN	ASP	GLU	VAL	LEU	ALA	CYS
ARG	THR	ARG	ARG	LEU	PHE	LEU
ARG	LEU	LYS	GLU	LEU	ASN	LEU
THR	GLN	LYS	LEU	LEU	THR	LEU
GLN	LEU	HIS	LYS	GLU	CYS	ILE
TRP	MET	SER	GLU	GLY	GLY	LYS
LYS	GLU	LEU	GLN	ALA	ASN	ASP
LEU	ARG	LYS	ALA	ALA	TRP	LYS
SER	SER	GLU	GLN	TRP	LYS	ASN
LEU	LEU	GLY	GLN	GLU	GLN	LEU
LEU	PRO	SER	ALA	GLU	ALA	TYR
ASP	GLU	PRO	GLY	ALA	LEU	ASN
	TRP	ILE	LEU	LEU	CYS	GLU
	TRP	GLU	ASP	ARG	VAL	ALA
	THR	ASP	ASP	LEU	ALA	LYS
	LEU	LEU	GLU	VAL	ALA	LYS
	THR	ALA	VAL	TYR	GLN	LEU
	TYR	LEU	PRO	LYS	LEU	TYR
	GLN	LEU	HIS	TYR	ASN	SER
	ASN	GLU	GLY	ASN	PHE	PRO
	ASN	ALA	GLN	ARG	THR	SER
	ALA	SER	SER	LEU	LYS	SER
	THR	GLU	ASP	ILE	ASP	GLN
	PRO	VAL	PHE	ILE	GLN	GLN
	VAL	VAL	LEU	GLU	VAL	TYR
	LEU	GLN	SER	THR	GLY	ASP
	GLY	ASN	GLU	ASN	LEU	ILE
	PRO	THR	THR	VAL	GLY	SER
	ASN	GLU	SER	LYS	ARG	ILE
	SER	ASN	SER	PRO	THR	ALA
	THR	LEU	VAL	SER	LEU	TYR
	ALA	LYS	VAL	ILE	ALA	GLY
	ASN	ASP	SER	LEU	GLY	ILE
	SER	GLU	CYS	THR	LYS	HIS


- Molecule 2: Elongator complex protein 2






• Molecule 3: Elongator complex protein 3

Chain C: 67% 25% 7%



• Molecule 4: tRNA Gln

Chain X: 35% 44% 43% 23%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	31251	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2700	Depositor
Magnification	96000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	2.128	Depositor
Minimum map value	-0.014	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.018	Depositor
Recommended contour level	0.11	Depositor
Map size (Å)	391.0, 391.0, 391.0	wwPDB
Map dimensions	460, 460, 460	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.85, 0.85, 0.85	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MG, 5AD, A2U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/5483	0.37	0/7466
2	B	0.17	0/6018	0.40	0/8196
3	C	0.16	0/4391	0.42	1/5933 (0.0%)
4	X	0.37	0/1778	0.53	0/2766
All	All	0.19	0/17670	0.41	1/24361 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	250	VAL	N-CA-C	-5.16	107.27	112.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5357	0	5268	104	0
2	B	5861	0	5704	140	0
3	C	4296	0	4324	98	0
4	X	1593	0	807	30	0
5	C	8	0	0	0	0
6	C	18	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	50	0	0	0	0
8	C	9	0	8	0	0
9	X	2	0	0	0	0
All	All	17194	0	16124	362	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 362 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:350:ASN:HD22	2:B:355:MET:H	1.18	0.91
4:X:12:G:O2'	4:X:13:U:OP1	1.91	0.87
2:B:574:THR:HG21	2:B:621:ALA:HA	1.59	0.83
1:A:311:GLU:HB3	1:A:314:SER:HB3	1.65	0.78
2:B:679:PRO:HG2	2:B:730:PRO:HA	1.65	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	681/1332 (51%)	625 (92%)	56 (8%)	0	100	100
2	B	735/826 (89%)	662 (90%)	73 (10%)	0	100	100
3	C	536/581 (92%)	497 (93%)	39 (7%)	0	100	100
All	All	1952/2739 (71%)	1784 (91%)	168 (9%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	608/1183 (51%)	607 (100%)	1 (0%)	87	85
2	B	647/722 (90%)	646 (100%)	1 (0%)	87	85
3	C	468/498 (94%)	466 (100%)	2 (0%)	84	82
All	All	1723/2403 (72%)	1719 (100%)	4 (0%)	85	85

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	573	LEU
2	B	376	ARG
3	C	250	VAL
3	C	434	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	196	HIS
3	C	528	ASN
2	B	114	HIS
2	B	370	GLN
2	B	410	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	X	74/75 (98%)	28 (37%)	2 (2%)

5 of 28 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	X	3	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	X	4	C
4	X	10	G
4	X	13	U
4	X	14	A

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	X	12	G
4	X	57	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	MET	C	604	5	7,8,8	0.89	1 (14%)	5,9,9	1.21	1 (20%)
7	A2U	C	603	-	50,52,52	2.96	15 (30%)	72,77,77	2.12	14 (19%)
5	SF4	C	601	3,8	0,12,12	-	-	-	-	-
6	5AD	C	602	-	20,20,20	0.56	0	28,30,30	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	MET	C	604	5	-	0/8/8/8	-
7	A2U	C	603	-	-	19/50/66/66	0/3/3/3
5	SF4	C	601	3,8	-	-	0/6/5/5
6	5AD	C	602	-	-	0/4/20/20	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	603	A2U	C45-C30	-7.85	1.32	1.52
7	C	603	A2U	O31-C30	6.90	1.60	1.45
7	C	603	A2U	P21-O24	6.78	1.66	1.59
7	C	603	A2U	P25-O24	6.77	1.66	1.59
7	C	603	A2U	C12-N11	6.53	1.48	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	603	A2U	O31-C32-N33	11.00	129.22	108.09
7	C	603	A2U	C43-C32-N33	-5.30	100.14	113.30
7	C	603	A2U	C36-C42-N41	-5.23	119.51	126.72
7	C	603	A2U	N41-C40-N39	-4.12	122.34	128.58
7	C	603	A2U	N41-C42-N33	3.60	133.30	127.17

There are no chirality outliers.

5 of 19 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	C	603	A2U	C17-C16-C19-O20
7	C	603	A2U	C14-C16-C19-O20
7	C	603	A2U	C18-C16-C19-O20
7	C	603	A2U	C19-O20-P21-O22
7	C	603	A2U	C19-O20-P21-O24


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17926. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 230




Y Index: 230




Z Index: 230


6.2.2 Raw map



X Index: 230



Y Index: 230



Z Index: 230

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 246




Y Index: 247




Z Index: 189


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 459


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.11. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_17926_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 90 nm³; this corresponds to an approximate mass of 81 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.299 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.299 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.35	-	-
Author-provided FSC curve	3.35	4.13	3.43
Unmasked-calculated*	8.94	17.83	9.34

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.94 differs from the reported value 3.35 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17926 and PDB model 8PTZ. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.11 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.11).

9.4 Atom inclusion [i](#)



At the recommended contour level, 78% of all backbone atoms, 70% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.11) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7010	0.3860
A	0.7170	0.4060
B	0.6800	0.3740
C	0.7880	0.4360
X	0.4850	0.2240

