



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:40 PM UTC

PDB ID : 4PU6 / pdb_00004pu6
Title : Crystal structure of potassium-dependent plant-type L-asparaginase from *Phaseolus vulgaris* in complex with K⁺ cations
Authors : Bejger, M.; Gilski, M.; Imiolczyk, B.; Jaskolski, M.
Deposited on : 2014-03-12
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	0% 65% 14% 20%
1	C	197	3% 64% 12% 20%
2	B	131	74% 25%
2	D	131	3% 67% 29%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-ASPARAGINASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	0	0
			1173	731	205	231	6			
1	C	157	Total	C	N	O	S	0	0	0
			1173	731	205	231	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP V7CU13
A	0	ALA	-	expression tag	UNP V7CU13
A	17	THR	ASN	SEE REMARK 999	UNP V7CU13
C	-1	GLY	-	expression tag	UNP V7CU13
C	0	ALA	-	expression tag	UNP V7CU13
C	17	THR	ASN	SEE REMARK 999	UNP V7CU13


- Molecule 2 is a protein called L-ASPARAGINASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	0	0
			944	580	167	184	13			
2	D	131	Total	C	N	O	S	0	0	0
			944	580	167	184	13			

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	C	2	Total	K	0	0
			2	2		

- Molecule 4 is ASPARTIC ACID (CCD ID: ASP) (formula: C₄H₇NO₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	O	
			9	4	1	4	
							0
							0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	48	Total	O		
			48	48	0	0
5	B	22	Total	O		
			22	22	0	0
5	C	34	Total	O		
			34	34	0	0
5	D	10	Total	O		
			10	10	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: L-ASPARAGINASE ALPHA SUBUNIT




• Molecule 1: L-ASPARAGINASE ALPHA SUBUNIT




• Molecule 2: L-ASPARAGINASE BETA SUBUNIT



• Molecule 2: L-ASPARAGINASE BETA SUBUNIT





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.94Å 102.67Å 127.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.60 – 2.30 47.60 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.3 (47.60-2.30) 96.2 (47.60-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.176 , 0.231 0.181 , 0.235	Depositor DCC
R_{free} test set	1009 reflections (3.10%)	wwPDB-VP
Wilson B-factor (Å ²)	47.7	Xtriage
Anisotropy	0.651	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 67.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4361	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/1190	1.03	0/1613
1	C	0.76	0/1190	1.01	2/1613 (0.1%)
2	B	0.79	0/954	1.01	2/1283 (0.2%)
2	D	0.69	0/954	1.00	5/1283 (0.4%)
All	All	0.75	0/4288	1.01	9/5792 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	228	SER	CA-C-N	-6.21	113.86	120.94
2	D	228	SER	C-N-CA	-6.21	113.86	120.94
1	C	18	LEU	CA-C-N	5.88	125.84	119.78
1	C	18	LEU	C-N-CA	5.88	125.84	119.78
2	D	318	GLY	N-CA-C	-5.45	108.13	115.21
2	D	257	LEU	N-CA-C	5.34	117.52	111.11
2	B	228	SER	CA-C-N	-5.31	114.89	120.94
2	B	228	SER	C-N-CA	-5.31	114.89	120.94
2	D	317	ASP	N-CA-C	-5.15	106.97	113.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1173	0	1176	21	0
1	C	1173	0	1176	22	0
2	B	944	0	930	23	0
2	D	944	0	930	27	0
3	A	2	0	0	0	0
3	C	2	0	0	0	0
4	B	9	0	3	1	0
5	A	48	0	0	4	0
5	B	22	0	0	0	0
5	C	34	0	0	0	0
5	D	10	0	0	0	0
All	All	4361	0	4215	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:14:VAL:O	2:D:307:ASN:ND2	2.20	0.71
2:B:311:ARG:HH22	2:B:326:ASP:HB3	1.55	0.71
1:A:2:GLY:N	5:A:336:HOH:O	2.28	0.66
2:B:259:ARG:HD3	2:D:256:THR:HG21	1.76	0.66
2:B:259:ARG:CD	2:D:256:THR:HG21	2.29	0.62
2:B:221:MET:HE3	1:C:117:HIS:CD2	2.34	0.62
1:A:22:ARG:HH22	1:A:63:LEU:HD11	1.67	0.59
1:C:40:ALA:O	1:C:45:VAL:HG13	2.03	0.59
1:A:23:GLN:HG3	2:B:324:ILE:HD12	1.85	0.58
2:D:315:THR:OG1	2:D:319:PHE:HB3	2.04	0.57
1:A:2:GLY:N	5:A:335:HOH:O	2.37	0.57
1:C:24:GLU:O	1:C:28:GLN:HG2	2.04	0.56
1:A:30:LEU:HD11	2:B:310:PHE:HB3	1.86	0.56
1:A:133:GLY:HA2	5:A:327:HOH:O	2.04	0.56
2:D:255:GLY:O	2:D:283:ARG:NH1	2.41	0.54
1:C:15:ASP:OD2	1:C:17:THR:HG22	2.09	0.52
1:A:74:GLU:O	1:A:155:LYS:NZ	2.36	0.52
1:A:81:GLU:HB3	1:A:96:SER:HA	1.91	0.52
2:D:205:GLU:OE1	2:D:205:GLU:N	2.39	0.52
1:A:72:LEU:HD21	1:A:143:VAL:HG13	1.91	0.52
2:D:284:LEU:HD13	2:D:305:ASN:ND2	2.25	0.51
2:D:315:THR:C	2:D:317:ASP:H	2.18	0.51
1:A:14:VAL:O	2:B:307:ASN:ND2	2.27	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:300:VAL:HG21	2:B:313:CYS:HB2	1.92	0.50
1:A:90:ARG:HH12	2:D:283:ARG:NH1	2.11	0.49
4:B:501:ASP:N	4:B:501:ASP:OD1	2.45	0.48
2:D:272:LEU:HD22	2:D:295:SER:HB3	1.94	0.48
2:B:229:PRO:HG2	2:B:230:LEU:HD13	1.95	0.48
1:A:117:HIS:CG	2:D:224:ARG:HG3	2.49	0.47
1:A:90:ARG:HH12	2:D:283:ARG:HH12	1.61	0.47
1:A:83:SER:HB3	2:B:229:PRO:HA	1.96	0.47
2:D:241:CYS:SG	2:D:262:ALA:HB2	2.55	0.47
2:B:204:ARG:NH2	2:B:316:GLU:OE2	2.45	0.47
1:C:20:LEU:HD13	1:C:21:GLU:N	2.30	0.47
2:B:224:ARG:HG3	1:C:117:HIS:CG	2.50	0.46
1:A:124:GLY:HA3	2:D:222:THR:O	2.16	0.46
2:D:256:THR:HG23	2:D:260:GLU:HG2	1.97	0.46
1:C:8:VAL:HG22	2:D:312:ALA:HB3	1.97	0.45
2:D:288:LYS:HA	2:D:305:ASN:OD1	2.16	0.45
2:D:255:GLY:HA2	2:D:283:ARG:HH12	1.82	0.45
1:A:46:PRO:O	1:A:50:VAL:HG23	2.16	0.45
1:A:66:SER:O	1:A:67:GLY:C	2.59	0.45
2:B:221:MET:CE	1:C:116:PRO:HD2	2.47	0.45
2:B:311:ARG:NH2	2:B:326:ASP:HB3	2.29	0.45
1:A:154:ALA:HA	5:A:344:HOH:O	2.17	0.45
2:B:252:ILE:HG23	2:B:257:LEU:HB2	1.98	0.44
2:B:276:VAL:HG21	2:B:301:ALA:HB3	2.00	0.44
2:B:284:LEU:HD13	2:B:284:LEU:HA	1.79	0.44
2:D:260:GLU:O	2:D:264:VAL:HG13	2.18	0.44
2:B:280:ILE:HA	2:B:284:LEU:HD23	1.99	0.44
2:B:204:ARG:NE	2:B:316:GLU:OE2	2.41	0.44
2:D:255:GLY:O	2:D:256:THR:HG22	2.17	0.44
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.91	0.44
1:C:4:TRP:CE2	2:D:316:GLU:HB2	2.52	0.44
2:B:243:VAL:HG21	2:B:261:VAL:HB	1.98	0.44
2:B:325:TRP:O	2:B:326:ASP:HB3	2.18	0.44
1:C:66:SER:O	1:C:67:GLY:C	2.61	0.43
2:B:221:MET:HE1	1:C:115:SER:HB2	2.00	0.43
2:D:264:VAL:O	2:D:268:LYS:HB2	2.19	0.43
2:D:281:LYS:HD3	2:D:282:HIS:NE2	2.34	0.43
1:C:15:ASP:CG	1:C:17:THR:HG22	2.44	0.43
1:C:18:LEU:HA	1:C:19:PRO:HD3	1.85	0.43
1:C:109:ARG:HD2	1:C:112:MET:HE2	2.00	0.43
1:C:150:MET:HB3	2:D:218:MET:HE3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83:SER:HB3	2:D:229:PRO:HA	2.01	0.42
2:D:259:ARG:O	2:D:259:ARG:HD3	2.19	0.42
1:C:140:GLU:H	1:C:140:GLU:CD	2.27	0.42
1:C:46:PRO:HD2	1:C:49:ASP:OD2	2.21	0.41
1:C:107:LEU:HD12	1:C:107:LEU:HA	1.84	0.41
1:C:114:LYS:HB3	1:C:114:LYS:HE3	1.83	0.41
2:D:285:ASP:C	2:D:287:GLY:H	2.27	0.41
2:B:265:MET:O	2:B:269:GLY:HA2	2.21	0.41
1:C:20:LEU:HD13	1:C:21:GLU:H	1.85	0.41
1:A:107:LEU:HA	1:A:107:LEU:HD12	1.72	0.41
1:A:112:MET:HE2	1:A:112:MET:HB3	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/197 (79%)	149 (96%)	6 (4%)	0	100	100
1	C	155/197 (79%)	149 (96%)	5 (3%)	1 (1%)	21	27
2	B	129/131 (98%)	120 (93%)	9 (7%)	0	100	100
2	D	129/131 (98%)	118 (92%)	11 (8%)	0	100	100
All	All	568/656 (87%)	536 (94%)	31 (6%)	1 (0%)	43	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	67	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/160 (79%)	124 (98%)	3 (2%)	43	62
1	C	127/160 (79%)	118 (93%)	9 (7%)	13	19
2	B	96/96 (100%)	87 (91%)	9 (9%)	8	11
2	D	96/96 (100%)	81 (84%)	15 (16%)	2	3
All	All	446/512 (87%)	410 (92%)	36 (8%)	10	14

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	VAL
1	A	18	LEU
1	A	153	LEU
2	B	217	LEU
2	B	230	LEU
2	B	240	VAL
2	B	246	THR
2	B	271	LYS
2	B	281	LYS
2	B	286	GLU
2	B	302	CYS
2	B	322	VAL
1	C	17	THR
1	C	20	LEU
1	C	21	GLU
1	C	39	SER
1	C	45	VAL
1	C	63	LEU
1	C	115	SER
1	C	146	ASP
1	C	158	ASN
2	D	202	VAL
2	D	217	LEU
2	D	230	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	246	THR
2	D	257	LEU
2	D	264	VAL
2	D	281	LYS
2	D	283	ARG
2	D	286	GLU
2	D	294	VAL
2	D	300	VAL
2	D	302	CYS
2	D	306	CYS
2	D	313	CYS
2	D	322	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	131	GLN
1	C	103	ASN
1	C	132	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ASP	B	501	-	7,8,8	1.17	0	6,10,10	1.58	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ASP	B	501	-	-	3/8/8/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	ASP	OXT-C-O	-3.15	116.92	124.08

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	501	ASP	O-C-CA-N
4	B	501	ASP	OXT-C-CA-N
4	B	501	ASP	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	501	ASP	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/197 (79%)	-0.23	2 (1%) 75 76	20, 41, 69, 105	2 (1%)
1	C	157/197 (79%)	-0.12	6 (3%) 44 46	22, 48, 82, 109	5 (3%)
2	B	131/131 (100%)	-0.06	0 100 100	20, 52, 76, 103	1 (0%)
2	D	131/131 (100%)	0.19	4 (3%) 51 53	27, 65, 92, 106	8 (6%)
All	All	576/656 (87%)	-0.07	12 (2%) 63 65	20, 49, 86, 109	16 (2%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	20	LEU	4.5
1	A	3	GLY	3.3
1	A	158	ASN	3.2
1	C	158	ASN	2.8
1	C	150	MET	2.5
2	D	281	LYS	2.5
1	C	153	LEU	2.4
1	C	157	ALA	2.3
2	D	315	THR	2.3
1	C	2	GLY	2.2
2	D	308	GLY	2.1
2	D	320	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ASP	B	501	9/9	0.87	0.11	25,37,47,50	9
3	K	C	201	1/1	0.97	0.07	57,57,57,57	0
3	K	C	202	1/1	0.97	0.07	47,47,47,47	0
3	K	A	201	1/1	0.97	0.07	41,41,41,41	0
3	K	A	202	1/1	0.98	0.04	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.