



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:28 AM UTC

PDB ID : 2PVV / pdb_00002pvv
Title : Structure of human glutamate carboxypeptidase II (GCPII) in complex with L-serine-O-sulfate
Authors : Barinka, C.; Lubkowski, J.
Deposited on : 2007-05-10
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	709	2% 82% 15%
2	B	2	100%
2	C	2	50% 50%
2	D	2	50% 50%
3	E	4	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	E	3	X	-	-	-
8	OSE	A	1768	X	-	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	17	0
			5628	3619	944	1046	19			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ARG	-	cloning artifact	UNP Q04609
A	43	SER	-	cloning artifact	UNP Q04609

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0

- Molecule 8 is O-SULFO-L-SERINE (CCD ID: OSE) (formula: $C_3H_7NO_6S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	N	O	S	0	0
			11	3	1	6	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	510	Total O 510 510	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate carboxypeptidase 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

- Molecule 3: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

100%

MAG1
MAG2
MAN3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.62Å 130.22Å 158.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.11 30.00 – 2.11	Depositor EDS
% Data completeness (in resolution range)	98.6 (30.00-2.11) 98.8 (30.00-2.11)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.49 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.174 , 0.212 0.171 , 0.209	Depositor DCC
R_{free} test set	3061 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.603	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6329	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, OSE, ZN, NAG, CL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	8/5826 (0.1%)	1.06	9/7895 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	705	PHE	N-CA	7.33	1.51	1.46
1	A	635	ALA	CA-CB	6.44	1.63	1.53
1	A	741	ALA	CA-CB	5.89	1.62	1.53
1	A	359	ILE	CA-CB	5.71	1.61	1.53
1	A	611	ILE	CA-CB	5.67	1.61	1.54
1	A	693	ALA	CA-CB	-5.59	1.46	1.53
1	A	148	PRO	CA-C	5.36	1.57	1.52
1	A	411	ARG	CA-C	-5.21	1.47	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	652	ASP	N-CA-C	7.31	121.91	112.92
1	A	147	PRO	O-C-N	6.90	124.39	121.15
1	A	548	GLY	N-CA-C	-6.04	105.89	112.04
1	A	266	ASP	CA-C-N	5.72	126.10	119.47
1	A	266	ASP	C-N-CA	5.72	126.10	119.47
1	A	309	MET	N-CA-C	5.71	118.52	109.96
1	A	251	GLY	CA-C-N	5.19	125.60	120.31
1	A	251	GLY	C-N-CA	5.19	125.60	120.31
1	A	142	SER	N-CA-C	5.07	117.25	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5628	0	5471	63	0
2	B	28	0	25	0	0
2	C	28	0	25	1	0
2	D	28	0	25	2	0
3	E	50	0	43	0	0
4	A	42	0	39	2	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	11	0	4	1	0
9	A	510	0	0	15	0
All	All	6329	0	5632	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.32	1.22
1:A:486:GLU:HG2	9:A:2025:HOH:O	1.68	0.94
1:A:620:GLN:NE2	9:A:2116:HOH:O	2.10	0.84
1:A:506:PHE:HB2	1:A:509:MET:HG3	1.69	0.74
1:A:177:VAL:HG12	1:A:188:LEU:HD11	1.73	0.71
1:A:367:GLU:OE1	1:A:662[A]:ARG:NH1	2.29	0.65
1:A:660[A]:VAL:O	1:A:664[A]:MET:CG	2.27	0.64
9:A:2140:HOH:O	2:C:2:NAG:H83	1.98	0.62
1:A:412:PRO:HA	1:A:589:ASN:OD1	2.02	0.59
1:A:506:PHE:CB	1:A:509:MET:HG3	2.32	0.59
1:A:169:MET:HE3	1:A:347:HIS:NE2	2.18	0.59
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.37	0.59
1:A:443:GLN:CG	9:A:2278:HOH:O	2.52	0.58
1:A:443:GLN:HG2	9:A:2278:HOH:O	2.05	0.57
1:A:152:GLU:OE2	9:A:1982:HOH:O	2.17	0.56
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.40	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:GLU:O	1:A:72:LYS:HG3	2.06	0.55
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.88	0.55
1:A:273:PRO:HB3	1:A:437:GLU:HB2	1.90	0.53
1:A:229:SER:O	1:A:299:TYR:HB3	2.09	0.53
9:A:1986:HOH:O	2:D:2:NAG:H83	2.08	0.53
1:A:310:GLY:O	1:A:328:ASN:HB3	2.09	0.53
1:A:303:GLN:OE1	1:A:325:VAL:HG13	2.09	0.52
1:A:295:HIS:HB2	1:A:346:ILE:HD11	1.92	0.52
1:A:169:MET:HE2	1:A:345:HIS:HD2	1.75	0.51
1:A:463:ARG:HH21	1:A:545:LYS:HE2	1.76	0.51
1:A:347:HIS:HE1	9:A:2104:HOH:O	1.94	0.51
1:A:169:MET:HA	1:A:344:MET:O	2.11	0.51
9:A:1986:HOH:O	2:D:2:NAG:C8	2.58	0.51
1:A:131:ILE:HG12	1:A:135:GLY:HA2	1.95	0.49
4:A:1757:NAG:H83	9:A:1984:HOH:O	2.12	0.49
1:A:453:ASP:HB3	1:A:536[B]:ARG:HH12	1.76	0.49
1:A:536[B]:ARG:NH2	9:A:2272:HOH:O	2.39	0.48
1:A:506:PHE:HB2	1:A:509:MET:CG	2.41	0.48
1:A:186:PHE:O	1:A:190:ARG:HB2	2.13	0.47
1:A:132:ASN:HD21	1:A:136:ASN:HD22	1.62	0.47
1:A:132:ASN:ND2	1:A:136:ASN:HB2	2.29	0.47
1:A:489:GLU:H	1:A:489:GLU:CD	2.24	0.46
1:A:485:ASP:O	9:A:1902:HOH:O	2.21	0.46
1:A:129:SER:HA	1:A:139:PHE:O	2.16	0.46
1:A:258:ILE:HD13	1:A:294:VAL:HB	1.98	0.45
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.17	0.45
1:A:691:ILE:O	1:A:704:SER:HA	2.17	0.45
1:A:369:ASP:OD2	1:A:662[B]:ARG:NH2	2.31	0.45
1:A:656[B]:SER:O	1:A:658[B]:PRO:HD3	2.16	0.45
1:A:246:TRP:CD1	4:A:1760:NAG:H83	2.51	0.45
1:A:443:GLN:HG3	9:A:2278:HOH:O	2.17	0.44
1:A:132:ASN:OD1	1:A:132:ASN:C	2.61	0.44
1:A:439:SER:O	1:A:443:GLN:HB2	2.17	0.44
1:A:358:VAL:HB	1:A:420:SER:HB3	1.99	0.44
1:A:110:LEU:CD2	1:A:358:VAL:HG22	2.47	0.44
1:A:234:TYR:O	1:A:550:PRO:HB3	2.18	0.44
1:A:463:ARG:NH2	1:A:545:LYS:HE2	2.33	0.44
1:A:659[B]:ILE:HD13	1:A:659[B]:ILE:HA	1.86	0.43
1:A:325:VAL:HB	1:A:326:PRO:HD2	1.99	0.43
1:A:427:GLY:HA2	8:A:1768:OSE:O3S	2.19	0.43
1:A:258:ILE:CD1	1:A:290:PRO:HG3	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:GLU:OE2	1:A:72:LYS:HE3	2.19	0.42
1:A:505:GLU:HG2	1:A:506:PHE:CE1	2.54	0.42
1:A:609:ASP:OD2	9:A:2126:HOH:O	2.21	0.42
1:A:469:LEU:O	1:A:595:PHE:HA	2.19	0.42
1:A:154:VAL:HG22	9:A:2266:HOH:O	2.19	0.41
1:A:171:GLU:HA	1:A:342:VAL:O	2.21	0.41
1:A:198:GLY:O	1:A:223:LYS:HE2	2.20	0.41
1:A:240:LYS:O	1:A:245:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	706/709 (100%)	679 (96%)	24 (3%)	3 (0%)	30	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	541	TRP
1	A	542	GLU
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	608/605 (100%)	593 (98%)	15 (2%)	42	47

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	MET
1	A	88[A]	GLN
1	A	88[B]	GLN
1	A	174	LEU
1	A	239	VAL
1	A	351	GLU
1	A	388	PRO
1	A	405	LEU
1	A	465	ASP
1	A	489	GLU
1	A	519	ASN
1	A	536[A]	ARG
1	A	536[B]	ARG
1	A	589	ASN
1	A	600	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	136	ASN
1	A	540	ASN
1	A	618	HIS
1	A	667	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	B	1	1,2	14,14,15	0.65	0	17,19,21	1.54	4 (23%)
2	NAG	B	2	2	14,14,15	0.69	0	17,19,21	2.09	4 (23%)
2	NAG	C	1	1,2	14,14,15	0.83	0	17,19,21	1.13	1 (5%)
2	NAG	C	2	2	14,14,15	0.41	0	17,19,21	1.58	5 (29%)
2	NAG	D	1	1,2	14,14,15	1.11	1 (7%)	17,19,21	1.68	4 (23%)
2	NAG	D	2	2	14,14,15	0.68	0	17,19,21	1.19	1 (5%)
3	NAG	E	1	1,3	14,14,15	0.75	0	17,19,21	1.76	3 (17%)
3	NAG	E	2	3	14,14,15	0.83	1 (7%)	17,19,21	1.34	3 (17%)
3	MAN	E	3	3	11,11,12	1.07	1 (9%)	15,15,17	1.33	1 (6%)
3	MAN	E	4	3	11,11,12	0.62	0	15,15,17	1.21	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	MAN	E	3	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	MAN	O5-C1	-2.98	1.38	1.43
3	E	2	NAG	O5-C1	-2.11	1.40	1.43
2	D	1	NAG	C1-C2	2.05	1.55	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	5.73	130.58	122.90
3	E	1	NAG	C1-O5-C5	4.80	118.62	112.19
2	B	2	NAG	C1-O5-C5	4.08	117.66	112.19
2	D	1	NAG	O5-C1-C2	-4.01	105.09	111.29
3	E	1	NAG	O5-C1-C2	-3.58	105.75	111.29
2	C	2	NAG	C8-C7-N2	3.49	121.90	116.12
2	B	2	NAG	C4-C3-C2	3.44	116.06	111.02
3	E	4	MAN	O5-C5-C6	3.42	114.31	107.66
2	B	1	NAG	O4-C4-C3	-3.28	102.65	110.38
2	D	2	NAG	C8-C7-N2	3.23	121.48	116.12
2	D	1	NAG	O5-C5-C4	-3.14	103.18	110.83
3	E	3	MAN	C1-O5-C5	2.93	116.11	112.19
3	E	2	NAG	C3-C4-C5	-2.75	105.24	110.23
3	E	2	NAG	C8-C7-N2	2.53	120.31	116.12
2	C	2	NAG	C2-N2-C7	2.49	126.23	122.90
3	E	1	NAG	C2-N2-C7	-2.39	119.70	122.90
2	D	1	NAG	C2-N2-C7	2.38	126.09	122.90
2	B	1	NAG	O5-C5-C6	2.33	112.20	107.66
2	C	2	NAG	O4-C4-C5	2.31	115.02	109.32
2	B	1	NAG	O5-C1-C2	-2.29	107.75	111.29
3	E	2	NAG	O7-C7-C8	-2.25	118.05	122.05
2	C	2	NAG	O7-C7-C8	-2.23	118.09	122.05
2	B	2	NAG	C3-C4-C5	2.20	114.22	110.23
2	C	2	NAG	C1-C2-N2	-2.17	107.02	110.43
2	B	1	NAG	C2-N2-C7	-2.10	120.09	122.90
2	C	1	NAG	O4-C4-C3	-2.03	105.60	110.38
2	D	1	NAG	O6-C6-C5	-2.02	104.45	111.33

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	3	MAN	C1

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
3	E	1	NAG	C4-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	NAG	2	0
2	C	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1759	1	14,14,15	0.44	0	17,19,21	1.14	1 (5%)
8	OSE	A	1768	-	9,10,10	1.31	1 (11%)	9,14,14	4.14	3 (33%)
4	NAG	A	1757	1	14,14,15	0.70	0	17,19,21	2.02	4 (23%)
4	NAG	A	1760	1	14,14,15	0.71	0	17,19,21	1.64	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1759	1	-	3/6/23/26	0/1/1/1
8	OSE	A	1768	-	1/1/3/3	2/10/10/10	-
4	NAG	A	1757	1	-	4/6/23/26	0/1/1/1
4	NAG	A	1760	1	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	1768	OSE	CA-N	-2.50	1.35	1.48

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	1768	OSE	OG-CB-CA	-11.33	94.33	107.48
4	A	1757	NAG	C1-O5-C5	6.45	120.83	112.19
4	A	1760	NAG	C1-O5-C5	4.32	117.97	112.19
8	A	1768	OSE	O3S-S-O2S	3.68	121.45	108.56
4	A	1759	NAG	C1-O5-C5	3.35	116.67	112.19
4	A	1757	NAG	C3-C4-C5	2.56	114.87	110.23
8	A	1768	OSE	O3S-S-OG	-2.48	100.67	106.37
4	A	1760	NAG	O7-C7-C8	-2.45	117.69	122.05
4	A	1757	NAG	C8-C7-N2	2.26	119.86	116.12
4	A	1757	NAG	C2-N2-C7	-2.13	120.04	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
8	A	1768	OSE	CA

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1759	NAG	C8-C7-N2-C2
4	A	1759	NAG	O7-C7-N2-C2
8	A	1768	OSE	C-CA-CB-OG
4	A	1757	NAG	C8-C7-N2-C2
4	A	1757	NAG	O7-C7-N2-C2
4	A	1757	NAG	O5-C5-C6-O6
4	A	1759	NAG	O5-C5-C6-O6
4	A	1757	NAG	C4-C5-C6-O6
4	A	1760	NAG	O5-C5-C6-O6
8	A	1768	OSE	O-C-CA-N

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	1768	OSE	1	0
4	A	1757	NAG	1	0
4	A	1760	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/709 (97%)	-0.21	17 (2%) 59 62	8, 26, 50, 71	18 (2%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	TRP	5.8
1	A	656[A]	SER	4.7
1	A	546	PHE	3.3
1	A	152	GLU	3.0
1	A	544	ASN	2.7
1	A	653	PHE	2.6
1	A	153	ASN	2.6
1	A	543	THR	2.5
1	A	657[A]	ASN	2.4
1	A	155[A]	SER	2.4
1	A	506	PHE	2.3
1	A	313	ALA	2.3
1	A	332	GLY	2.2
1	A	124	HIS	2.2
1	A	548	GLY	2.1
1	A	547	SER	2.1
1	A	154	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.69	0.14	61,64,68,69	0
2	NAG	B	2	14/15	0.73	0.15	46,55,60,61	0
2	NAG	C	1	14/15	0.84	0.12	47,49,54,59	0
3	NAG	E	2	14/15	0.85	0.13	37,42,46,54	0
2	NAG	D	2	14/15	0.87	0.10	35,42,48,49	0
3	MAN	E	4	11/12	0.88	0.11	50,52,55,57	0
3	MAN	E	3	11/12	0.89	0.09	42,46,47,49	0
2	NAG	D	1	14/15	0.93	0.08	28,31,37,40	0
2	NAG	B	1	14/15	0.94	0.09	37,42,46,50	0
3	NAG	E	1	14/15	0.97	0.05	16,26,37,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	1757	14/15	0.69	0.15	51,57,64,64	0
4	NAG	A	1759	14/15	0.69	0.15	68,73,76,76	0
4	NAG	A	1760	14/15	0.83	0.12	36,52,56,59	0
6	CA	A	1753	1/1	0.99	0.02	16,16,16,16	0
8	OSE	A	1768	11/11	0.99	0.04	22,22,27,32	0
5	ZN	A	1751	1/1	1.00	0.03	24,24,24,24	0
7	CL	A	1754	1/1	1.00	0.01	20,20,20,20	0
5	ZN	A	1752	1/1	1.00	0.03	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.