



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:35 PM UTC

PDB ID : 3PV6 / pdb_00003pv6
Title : Crystal structure of NKp30 bound to its ligand B7-H6
Authors : Li, Y.
Deposited on : 2010-12-06
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	
2	B	117	
3	C	2	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig-like domain-containing protein DKFZp686O24166/DKFZp686I21167.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1681	1075	277	319	10			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	-	expression tag	UNP Q68D85
A	263	ALA	-	expression tag	UNP Q68D85
A	264	ALA	-	expression tag	UNP Q68D85
A	265	ALA	-	expression tag	UNP Q68D85
A	266	HIS	-	expression tag	UNP Q68D85
A	267	HIS	-	expression tag	UNP Q68D85
A	268	HIS	-	expression tag	UNP Q68D85
A	269	HIS	-	expression tag	UNP Q68D85
A	270	HIS	-	expression tag	UNP Q68D85
A	271	HIS	-	expression tag	UNP Q68D85

- Molecule 2 is a protein called Natural cytotoxicity triggering receptor 3.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	112	Total	C	N	O	S	0	0	0
			859	538	162	157	2			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	9	Total O 9 9	0	0
5	B	6	Total O 6 6	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig-like domain-containing protein DKFZp686O24166/DKFZp686I21167



- Molecule 2: Natural cytotoxicity triggering receptor 3



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.87Å 74.87Å 125.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-2.30) 93.1 (50.00-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.42 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.239 , 0.289 0.239 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.701	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2611	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	3/1721 (0.2%)	1.31	22/2339 (0.9%)
2	B	1.06	0/878	1.26	6/1191 (0.5%)
All	All	1.08	3/2599 (0.1%)	1.29	28/3530 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	200	ILE	CA-CB	5.52	1.63	1.55
1	A	31	MET	SD-CE	5.34	1.93	1.79
1	A	112	ILE	CA-CB	5.19	1.60	1.54

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	ILE	N-CA-C	8.00	118.91	107.88
1	A	144	ALA	N-CA-C	-7.75	96.77	109.40
1	A	26	LEU	N-CA-C	7.11	120.77	108.76
1	A	25	ASP	N-CA-C	7.09	120.70	109.50
2	B	48	LEU	N-CA-C	-6.94	99.15	109.15
2	B	42	ASN	N-CA-C	6.77	120.35	107.75
1	A	236	HIS	N-CA-C	-6.76	103.53	111.03
1	A	44	VAL	N-CA-C	6.73	118.84	109.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	SER	CA-C-N	6.51	126.74	119.32
1	A	96	SER	C-N-CA	6.51	126.74	119.32
1	A	219	GLN	N-CA-C	-6.23	103.73	113.02
1	A	90	ARG	N-CA-C	-6.03	97.67	108.47
1	A	230	VAL	N-CA-C	5.91	116.38	107.75
1	A	126	VAL	N-CA-C	-5.78	97.70	106.42
2	B	47	ARG	N-CA-C	-5.72	98.96	108.75
1	A	166	SER	N-CA-C	5.62	117.73	109.24
1	A	160	LYS	N-CA-C	5.55	118.86	109.76
1	A	223	GLY	N-CA-C	-5.53	108.29	115.59
1	A	186	HIS	CA-C-N	5.41	125.67	119.83
1	A	186	HIS	C-N-CA	5.41	125.67	119.83
1	A	47	PHE	N-CA-C	5.36	118.67	110.20
1	A	74	LYS	N-CA-C	5.33	122.14	110.80
1	A	118	GLY	CA-C-N	-5.31	115.50	122.99
1	A	118	GLY	C-N-CA	-5.31	115.50	122.99
2	B	101	HIS	N-CA-C	-5.29	106.60	113.16
1	A	89	PHE	N-CA-C	-5.23	105.64	112.23
2	B	110	VAL	N-CA-C	5.07	115.77	108.48
2	B	102	ASP	N-CA-C	-5.05	106.97	113.23

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	169	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1681	0	1647	56	0
2	B	859	0	840	29	0
3	C	28	0	25	0	0
4	A	28	0	26	0	0
5	A	9	0	0	0	0
5	B	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2611	0	2538	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:MET:HE3	1:A:31:MET:HA	1.51	0.91
1:A:146:ARG:HH22	1:A:148:LEU:HD11	1.41	0.86
1:A:162:MET:HE2	1:A:212:CYS:SG	2.21	0.81
1:A:160:LYS:HG2	1:A:214:LYS:HG3	1.65	0.79
1:A:160:LYS:HE3	1:A:214:LYS:HD3	1.66	0.77
1:A:128:PRO:HD2	1:A:129:LEU:HD13	1.67	0.76
1:A:149:LEU:HD12	1:A:160:LYS:O	1.85	0.76
2:B:59:GLU:HG2	2:B:64:LYS:HG3	1.70	0.72
1:A:179:LYS:HB3	1:A:190:ILE:HD11	1.71	0.72
1:A:146:ARG:NH2	1:A:148:LEU:HD11	2.04	0.72
2:B:37:LEU:HD12	2:B:93:LEU:HD23	1.72	0.70
1:A:192:GLU:HG3	1:A:193:ASP:H	1.61	0.65
1:A:216:ASN:OD1	1:A:219:GLN:HB2	1.97	0.65
2:B:65:GLU:O	2:B:68:ASN:HB2	1.98	0.64
2:B:66:VAL:HG23	2:B:67:ARG:N	2.12	0.63
1:A:179:LYS:HE2	1:A:188:ILE:HD11	1.80	0.63
2:B:76:ARG:NH2	2:B:102:ASP:OD1	2.30	0.63
1:A:31:MET:HA	1:A:31:MET:CE	2.28	0.62
1:A:225:VAL:HG22	1:A:244:THR:HG22	1.80	0.61
1:A:72:PHE:CE2	1:A:74:LYS:HB2	2.35	0.61
2:B:56:PHE:CE2	2:B:65:GLU:HG2	2.35	0.61
2:B:84:ARG:O	2:B:88:ASP:HB2	2.03	0.59
2:B:33:SER:O	2:B:98:VAL:HG13	2.03	0.58
2:B:24:PRO:HG3	2:B:38:PRO:HG2	1.85	0.58
2:B:66:VAL:C	2:B:68:ASN:H	2.12	0.57
2:B:88:ASP:O	2:B:89:HIS:HB2	2.04	0.57
1:A:192:GLU:O	1:A:193:ASP:HB2	2.03	0.57
1:A:71:THR:OG1	1:A:72:PHE:HD1	1.88	0.56
1:A:71:THR:HG1	1:A:72:PHE:HD1	1.52	0.56
2:B:45:GLN:HG3	2:B:46:GLY:N	2.21	0.55
2:B:45:GLN:HG3	2:B:46:GLY:H	1.72	0.55
1:A:222:PRO:C	1:A:224:THR:H	2.15	0.54
1:A:192:GLU:HG3	1:A:193:ASP:N	2.22	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:THR:O	1:A:245:LEU:CG	2.57	0.53
1:A:160:LYS:HE3	1:A:214:LYS:CD	2.38	0.52
2:B:66:VAL:CG2	2:B:67:ARG:N	2.72	0.52
1:A:188:ILE:O	1:A:188:ILE:HD12	2.10	0.51
2:B:44:SER:HB2	2:B:47:ARG:HB3	1.92	0.51
1:A:39:PRO:HG2	1:A:42:ASP:CG	2.36	0.50
1:A:80:GLU:OE2	1:A:85:HIS:HE1	1.95	0.50
2:B:41:PHE:CZ	2:B:89:HIS:HA	2.47	0.50
1:A:99:ARG:HB3	1:A:104:ASP:HB3	1.94	0.49
1:A:94:ILE:CG1	1:A:95:VAL:N	2.77	0.48
2:B:76:ARG:HH22	2:B:102:ASP:CG	2.22	0.48
1:A:194:VAL:O	1:A:195:ILE:HD13	2.13	0.48
1:A:58:ILE:HA	1:A:61:MET:SD	2.55	0.47
1:A:94:ILE:HG12	1:A:95:VAL:N	2.28	0.47
2:B:48:LEU:HD13	2:B:86:LEU:HB3	1.96	0.47
1:A:39:PRO:O	1:A:42:ASP:HB2	2.14	0.47
1:A:186:HIS:HD2	1:A:187:PRO:O	1.98	0.47
1:A:56:LEU:HD22	1:A:61:MET:HE3	1.97	0.47
1:A:68:LYS:NZ	1:A:73:ASP:O	2.42	0.47
1:A:161:TYR:CE1	1:A:215:LEU:HD12	2.50	0.47
1:A:188:ILE:HD12	1:A:188:ILE:C	2.40	0.47
2:B:24:PRO:CG	2:B:38:PRO:HG2	2.44	0.47
1:A:186:HIS:HA	1:A:187:PRO:HD3	1.75	0.46
1:A:244:THR:O	1:A:245:LEU:CB	2.63	0.46
2:B:74:ARG:HB3	2:B:74:ARG:CZ	2.45	0.46
1:A:160:LYS:CG	1:A:214:LYS:HG3	2.43	0.46
1:A:215:LEU:HB3	1:A:216:ASN:H	1.57	0.45
1:A:67:TRP:CH2	1:A:69:SER:HB2	2.51	0.45
1:A:222:PRO:C	1:A:224:THR:N	2.75	0.45
1:A:50:ILE:HG22	1:A:51:PHE:N	2.31	0.45
2:B:59:GLU:HG3	2:B:61:VAL:HG22	1.99	0.45
1:A:147:LEU:HD13	1:A:243:PHE:CD2	2.51	0.45
1:A:126:VAL:O	1:A:127:THR:C	2.60	0.45
1:A:70:LEU:HA	1:A:70:LEU:HD23	1.70	0.44
2:B:40:SER:HB3	2:B:90:GLN:HE22	1.83	0.44
1:A:220:GLU:HB3	1:A:226:TYR:OH	2.17	0.44
2:B:35:ALA:HB2	2:B:98:VAL:HG11	2.00	0.44
1:A:227:GLN:HB3	1:A:242:ASN:HB3	1.99	0.43
1:A:244:THR:O	1:A:245:LEU:HG	2.18	0.43
1:A:145:SER:HA	1:A:164:GLU:O	2.19	0.43
1:A:142:SER:HA	1:A:234:SER:OG	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:THR:O	1:A:245:LEU:HD23	2.19	0.43
2:B:66:VAL:C	2:B:68:ASN:N	2.77	0.42
1:A:132:GLN:HE21	1:A:132:GLN:HB3	1.69	0.42
2:B:40:SER:HB3	2:B:90:GLN:NE2	2.34	0.42
2:B:48:LEU:CD1	2:B:86:LEU:HB3	2.50	0.42
2:B:73:PHE:O	2:B:74:ARG:C	2.63	0.42
2:B:56:PHE:CE2	2:B:65:GLU:CG	3.01	0.41
1:A:120:TYR:O	1:A:134:THR:HA	2.21	0.41
1:A:220:GLU:C	1:A:221:ASP:O	2.64	0.40
2:B:112:VAL:HG11	2:B:115:LEU:HD22	2.04	0.40
1:A:40:LEU:HD12	1:A:41:ASN:N	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/248 (85%)	188 (90%)	18 (9%)	4 (2%)		
2	B	110/117 (94%)	96 (87%)	14 (13%)	0		
All	All	320/365 (88%)	284 (89%)	32 (10%)	4 (1%)		

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	LYS
1	A	159	ASP
1	A	216	ASN
1	A	197	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/218 (87%)	171 (90%)	18 (10%)	8	10
2	B	91/97 (94%)	81 (89%)	10 (11%)	6	7
All	All	280/315 (89%)	252 (90%)	28 (10%)	7	9

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	VAL
1	A	31	MET
1	A	41	ASN
1	A	42	ASP
1	A	58	ILE
1	A	71	THR
1	A	73	ASP
1	A	97	PRO
1	A	98	TRP
1	A	102	SER
1	A	125	VAL
1	A	128	PRO
1	A	132	GLN
1	A	159	ASP
1	A	200	ILE
1	A	214	LYS
1	A	219	GLN
1	A	245	LEU
2	B	33	SER
2	B	47	ARG
2	B	48	LEU
2	B	68	ASN
2	B	94	HIS
2	B	98	VAL
2	B	99	ARG
2	B	104	SER
2	B	113	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	115	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	54	GLN
1	A	85	HIS
1	A	132	GLN
1	A	174	ASN
1	A	182	GLN
1	A	186	HIS
2	B	68	ASN
2	B	90	GLN
2	B	121	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.96	0	17,19,21	0.87	1 (5%)
3	NAG	C	2	3	14,14,15	1.04	1 (7%)	17,19,21	1.51	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	C4-C5	2.33	1.58	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C3-C4-C5	2.85	115.39	110.23
3	C	2	NAG	C4-C3-C2	2.59	114.81	111.02
3	C	2	NAG	C2-N2-C7	-2.58	119.44	122.90
3	C	2	NAG	O5-C1-C2	-2.44	107.52	111.29
3	C	1	NAG	O5-C1-C2	-2.24	107.82	111.29

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	830	1	14,14,15	1.21	1 (7%)	17,19,21	1.04	2 (11%)
4	NAG	A	820	1	14,14,15	1.05	1 (7%)	17,19,21	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	830	1	-	5/6/23/26	0/1/1/1
4	NAG	A	820	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	830	NAG	C1-C2	3.15	1.56	1.52
4	A	820	NAG	C3-C2	2.51	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	830	NAG	C4-C3-C2	2.23	114.29	111.02
4	A	830	NAG	C1-O5-C5	-2.21	109.22	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	830	NAG	C3-C2-N2-C7
4	A	830	NAG	C8-C7-N2-C2
4	A	830	NAG	O7-C7-N2-C2
4	A	820	NAG	C4-C5-C6-O6
4	A	830	NAG	O5-C5-C6-O6
4	A	820	NAG	O5-C5-C6-O6
4	A	830	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/248 (86%)	0.92	27 (12%) 8 9	32, 48, 71, 81	0
2	B	112/117 (95%)	0.94	15 (13%) 7 8	35, 49, 78, 87	0
All	All	326/365 (89%)	0.93	42 (12%) 7 8	32, 48, 75, 87	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	SER	5.7
1	A	245	LEU	4.8
1	A	217	SER	4.8
1	A	149	LEU	4.5
1	A	158	GLU	4.3
2	B	36	PHE	4.0
2	B	42	ASN	4.0
1	A	24	ALA	3.9
1	A	98	TRP	3.5
1	A	72	PHE	3.3
2	B	71	PRO	3.3
1	A	51	PHE	3.3
2	B	130	GLU	3.1
1	A	148	LEU	3.1
1	A	161	TYR	3.0
2	B	67	ARG	3.0
1	A	216	ASN	3.0
2	B	45	GLN	2.9
1	A	159	ASP	2.8
1	A	188	ILE	2.8
1	A	84	ASP	2.7
1	A	108	ARG	2.7
2	B	46	GLY	2.6
1	A	212	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	53	SER	2.5
2	B	68	ASN	2.5
1	A	28	VAL	2.4
2	B	66	VAL	2.4
1	A	56	LEU	2.4
1	A	138	GLU	2.4
1	A	236	HIS	2.4
2	B	44	SER	2.3
2	B	20	TRP	2.3
2	B	73	PHE	2.2
1	A	74	LYS	2.2
2	B	127	VAL	2.1
2	B	47	ARG	2.1
1	A	25	ASP	2.1
2	B	101	HIS	2.1
1	A	71	THR	2.1
1	A	222	PRO	2.0
1	A	47	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	2	14/15	0.77	0.14	56,60,62,63	0
3	NAG	C	1	14/15	0.94	0.08	41,45,48,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	830	14/15	0.61	0.16	75,78,79,79	0
4	NAG	A	820	14/15	0.73	0.17	60,64,65,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.