



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:46 PM UTC

PDB ID : 4PV9 / pdb_00004pv9
Title : Crystal Structure of H2Kb-Q600V complex
Authors : Twist, K.-A.; Rossjohn, J.; Gras, S.
Deposited on : 2014-03-15
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	4% 93% 7%
1	C	278	5% 93% 5%
2	B	99	4% 89% 11%
2	D	99	2% 90% 10%
3	E	8	88% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	8	 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	S	0	12	0
			2332	1479	412	432	9			
1	C	277	Total	C	N	O	S	0	8	0
			2300	1457	405	429	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	2	0
			835	534	140	153	8			
2	D	99	Total	C	N	O	S	0	1	0
			829	530	139	153	7			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	ASP	ALA	conflict	UNP P01887
D	85	ASP	ALA	conflict	UNP P01887

- Molecule 3 is a protein called S598 peptide modified Q600V.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	8	Total	C	N	O	0	1	0
			73	48	15	10			
3	F	8	Total	C	N	O	0	0	0
			65	43	12	10			

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	295	Total 295	O 295	0	0
6	B	109	Total 109	O 109	0	0
6	C	253	Total 253	O 253	0	0
6	D	106	Total 106	O 106	0	0
6	E	10	Total 10	O 10	0	0
6	F	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: S598 peptide modified Q600V



- Molecule 3: S598 peptide modified Q600V

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.47Å 90.75Å 89.52Å 90.00° 111.58° 90.00°	Depositor
Resolution (Å)	61.81 – 2.00 61.81 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.9 (61.81-2.00) 97.9 (61.81-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.00Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.10.0	Depositor
R, R_{free}	0.173 , 0.218 0.192 , 0.237	Depositor DCC
R_{free} test set	3309 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.560	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 74.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7236	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0088e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ABA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/2432	1.16	2/3303 (0.1%)
1	C	0.81	0/2378	1.19	5/3230 (0.2%)
2	B	0.81	0/864	1.12	0/1170
2	D	0.81	0/855	1.12	4/1159 (0.3%)
3	E	0.72	0/68	0.82	0/85
3	F	0.93	0/58	0.96	0/74
All	All	0.82	0/6655	1.16	11/9021 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	197	ASP	CA-CB-CG	7.89	120.49	112.60
1	C	40	ALA	CA-C-N	5.67	127.89	120.28
1	C	40	ALA	C-N-CA	5.67	127.89	120.28
1	A	220	ASN	CA-CB-CG	5.57	118.17	112.60
1	C	161	GLU	CB-CG-CD	5.49	121.93	112.60
1	C	129	ASP	CA-CB-CG	5.42	118.02	112.60
2	D	71	THR	CB-CA-C	5.39	115.83	110.33
1	A	277	PRO	N-CA-C	5.20	117.04	110.70
2	D	75	THR	N-CA-C	5.09	119.76	113.50
2	D	84	HIS	CA-C-N	5.07	127.78	120.38
2	D	84	HIS	C-N-CA	5.07	127.78	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2332	0	2271	14	0
1	C	2300	0	2219	7	0
2	B	835	0	817	10	0
2	D	829	0	806	4	0
3	E	73	0	87	0	0
3	F	65	0	72	0	0
4	C	4	0	3	0	0
4	D	4	0	3	0	0
5	C	6	0	8	0	0
6	A	295	0	0	5	0
6	B	109	0	0	0	0
6	C	253	0	0	0	0
6	D	106	0	0	1	0
6	E	10	0	0	0	0
6	F	15	0	0	0	0
All	All	7236	0	6286	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:MET:HG2	6:A:553:HOH:O	1.89	0.72
1:A:138:MET:CG	6:A:553:HOH:O	2.39	0.70
2:D:37:ILE:HB	2:D:51:MET:HE1	1.79	0.65
1:A:94[B]:THR:HG21	2:B:31:HIS:CE1	2.37	0.59
1:C:172:LEU:O	1:C:176:ASN:HB2	2.03	0.59
2:D:67:HIS:HD2	6:D:289:HOH:O	1.88	0.56
2:B:33:PRO:O	2:B:54:MET:HE1	2.09	0.52
1:A:138:MET:HG3	6:A:553:HOH:O	2.08	0.51
2:B:39:MET:H	2:B:45:LYS:HZ2	1.60	0.50
2:B:36:GLU:HB2	2:B:83:LYS:HB3	1.94	0.49
1:A:94[B]:THR:CG2	2:B:31:HIS:HE1	2.26	0.48
1:A:45:TYR:HE1	1:A:67:ALA:HB2	1.79	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:215:LEU:HD22	1:C:261:VAL:HG22	1.96	0.48
1:A:172:LEU:O	1:A:176:ASN:HB2	2.14	0.47
1:C:81:LEU:HD13	1:C:118:TYR:CD1	2.50	0.46
1:A:94[B]:THR:HG21	2:B:31:HIS:HE1	1.77	0.45
1:A:10[A]:THR:HG22	6:A:313:HOH:O	2.15	0.45
1:C:189:VAL:HG23	1:C:272[A]:LEU:HD13	1.98	0.45
2:B:39:MET:H	2:B:45:LYS:NZ	2.13	0.45
1:A:24:GLU:HB2	1:A:36:PHE:HB3	2.00	0.44
1:A:187:ALA:HA	1:A:204:TRP:O	2.17	0.43
2:D:7:ILE:HD12	2:D:91:LYS:HE3	2.00	0.43
2:B:37:ILE:HD12	2:B:51[B]:MET:HE1	2.00	0.43
2:B:37:ILE:HD12	2:B:51[A]:MET:HE1	2.00	0.43
2:B:32:PRO:HD2	2:B:85:ASP:OD1	2.20	0.42
1:A:273:ARG:NH2	6:A:477:HOH:O	2.50	0.41
1:A:13:SER:HB3	1:A:78:LEU:HD13	2.01	0.41
1:C:19:GLU:OE1	1:C:75:ARG:NH1	2.53	0.41
1:C:23:MET:HE2	1:C:23:MET:HB2	1.92	0.41
1:C:45:TYR:HE1	1:C:67:ALA:HB2	1.86	0.41
2:D:36:GLU:HB2	2:D:83:LYS:HB3	2.02	0.41
1:A:160:LEU:O	1:A:165:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/278 (104%)	282 (98%)	6 (2%)	0	100	100
1	C	282/278 (101%)	274 (97%)	8 (3%)	0	100	100
2	B	99/99 (100%)	96 (97%)	3 (3%)	0	100	100
2	D	98/99 (99%)	94 (96%)	4 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	5/8 (62%)	5 (100%)	0	0	100	100
3	F	5/8 (62%)	5 (100%)	0	0	100	100
All	All	777/770 (101%)	756 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/236 (105%)	245 (99%)	3 (1%)	63	70
1	C	242/236 (102%)	238 (98%)	4 (2%)	53	60
2	B	96/94 (102%)	96 (100%)	0	100	100
2	D	95/94 (101%)	95 (100%)	0	100	100
3	E	7/6 (117%)	7 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	694/672 (103%)	687 (99%)	7 (1%)	68	75

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ARG
1	A	45	TYR
1	A	103	VAL
1	C	14	ARG
1	C	23	MET
1	C	45	TYR
1	C	197	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	72	GLN
1	A	86	ASN
1	A	220	ASN
2	B	67	HIS
1	C	65	GLN
1	C	87	GLN
2	D	13	HIS
2	D	34	HIS
2	D	38	GLN
2	D	67	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ABA	E	2	3	4,5,6	2.25	1 (25%)	1,5,7	0.37	0
3	ABA	F	2	3	4,5,6	1.11	0	1,5,7	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ABA	E	2	3	-	0/3/4/6	-
3	ABA	F	2	3	-	0/3/4/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	ABA	CB-CA	4.30	1.64	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	C	302	-	5,5,5	0.12	0	5,5,5	0.62	0
4	ACT	C	301	-	3,3,3	1.16	0	3,3,3	0.94	0
4	ACT	D	101	-	3,3,3	1.25	0	3,3,3	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	302	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	302	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	278/278 (100%)	0.14	10 (3%)	46	45	11, 31, 54, 82	17 (6%)
1	C	277/278 (99%)	0.17	14 (5%)	33	32	15, 33, 53, 70	19 (6%)
2	B	99/99 (100%)	0.22	4 (4%)	42	41	21, 33, 52, 70	8 (8%)
2	D	99/99 (100%)	0.10	2 (2%)	65	64	17, 31, 52, 70	9 (9%)
3	E	7/8 (87%)	-0.14	0	100	100	20, 23, 32, 34	1 (14%)
3	F	7/8 (87%)	0.13	0	100	100	25, 26, 40, 47	0
All	All	767/770 (99%)	0.15	30 (3%)	43	42	11, 32, 53, 82	54 (7%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	10.4
2	D	1	ILE	7.3
1	C	225	ILE	5.9
1	A	138	MET	4.3
2	B	85	ASP	3.8
1	C	227	ASP	3.8
1	A	225	ILE	3.5
1	A	278	PRO	3.4
1	A	89	LYS	3.3
1	A	40	ALA	3.1
1	C	277	PRO	3.0
1	C	221	GLY	2.9
1	C	226	GLN	2.8
2	B	48	LYS	2.8
2	D	85	ASP	2.6
1	A	170[A]	ARG	2.6
2	B	19[A]	LYS	2.5
1	C	41	GLU	2.5
1	C	224	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	40	ALA	2.4
1	C	195	PRO	2.4
1	A	16	GLY	2.3
1	C	197	ASP	2.3
1	A	195	PRO	2.3
1	C	196	GLU	2.2
1	C	89	LYS	2.2
1	A	212	ASP	2.1
1	C	128	GLU	2.0
1	A	42	ASN	2.0
1	C	181	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ABA	E	2	6/7	0.95	0.07	17,18,19,19	0
3	ABA	F	2	6/7	0.96	0.06	20,21,22,25	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	D	101	4/4	0.54	0.17	64,66,66,67	0
4	ACT	C	301	4/4	0.81	0.25	47,51,53,56	0
5	GOL	C	302	6/6	0.82	0.15	54,59,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.