



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:51 PM UTC

PDB ID : 2PWV / pdb_00002pwv
Title : tRNA guanine transglycosylase E235Q mutant in complex with preQ0
Authors : Tidten, N.; Stengl, B.; Heine, A.; Reuter, K.; Klebe, G.
Deposited on : 2007-05-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PQ0	A	500	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Queuine tRNA-ribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	2	0
			2809	1762	508	518	21			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	235	GLN	GLU	engineered mutation	UNP P28720
A	312	LYS	THR	SEE REMARK 999	UNP P28720

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 2-AMINO-4-OXO-4,7-DIHYDRO-3H-PYRROLO[2,3-D]PYRIMIDINE-5-CARBONITRILE (CCD ID: PQ0) (formula: C₇H₅N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	7	5	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	228	Total	O	0	0
			228	228		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Queuine tRNA-ribosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.76Å 65.17Å 70.45Å 90.00° 96.12° 90.00°	Depositor
Resolution (Å)	10.00 – 1.70 10.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	90.0 (10.00-1.70) 94.3 (10.00-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.70Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.201 , 0.245 0.200 , 0.240	Depositor DCC
R_{free} test set	2133 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.434	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 74.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3087	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PQ0, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2873	1.51	32/3871 (0.8%)

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ARG	CD-NE-CZ	8.89	136.84	124.40
1	A	246	PHE	CA-CB-CG	8.31	122.11	113.80
1	A	199	PHE	CA-CB-CG	7.07	120.87	113.80
1	A	42	PHE	CA-CB-CG	-6.75	107.05	113.80
1	A	226	TYR	CA-C-O	6.72	127.37	120.24
1	A	202	GLN	CA-C-N	6.44	133.84	121.54
1	A	202	GLN	C-N-CA	6.44	133.84	121.54
1	A	138	GLU	CA-C-N	6.40	129.37	120.29
1	A	138	GLU	C-N-CA	6.40	129.37	120.29
1	A	138	GLU	O-C-N	-6.39	115.49	122.07
1	A	151	ILE	CA-C-N	6.29	130.34	121.66
1	A	151	ILE	C-N-CA	6.29	130.34	121.66
1	A	166	SER	CA-C-N	6.23	128.62	120.28
1	A	166	SER	C-N-CA	6.23	128.62	120.28
1	A	177	ARG	CD-NE-CZ	6.04	132.85	124.40
1	A	186	PHE	CA-C-N	6.04	128.97	120.28
1	A	186	PHE	C-N-CA	6.04	128.97	120.28
1	A	362	ARG	NE-CZ-NH2	5.97	124.57	119.20
1	A	43	MET	O-C-N	5.81	126.65	121.19
1	A	195	ASN	CA-CB-CG	-5.73	106.87	112.60
1	A	272	VAL	O-C-N	-5.58	116.08	121.83
1	A	16	PHE	CA-CB-CG	5.56	119.36	113.80
1	A	88	GLY	CA-C-O	-5.45	118.22	122.52
1	A	336	ARG	NE-CZ-NH1	-5.45	116.05	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	349	HIS	CA-CB-CG	-5.42	108.38	113.80
1	A	19	ALA	CA-C-O	5.31	125.81	119.60
1	A	85	LYS	CA-C-O	5.31	126.18	120.55
1	A	315	ASP	CA-C-N	5.29	127.89	120.28
1	A	315	ASP	C-N-CA	5.29	127.89	120.28
1	A	301	ASN	N-CA-CB	5.28	119.08	110.37
1	A	195	ASN	OD1-CG-ND2	5.20	127.80	122.60
1	A	186	PHE	CA-C-O	-5.00	115.55	120.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2809	0	2742	22	0
2	A	1	0	0	0	0
3	A	13	0	5	2	0
4	A	36	0	48	6	0
5	A	228	0	0	5	0
All	All	3087	0	2795	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:500:PQ0:N3	3:A:500:PQ0:C2	1.73	1.51
1:A:191:GLU:HG3	4:A:705:GOL:H12	1.69	0.72
1:A:72:TYR:O	1:A:75:MET:HG2	1.92	0.69
1:A:252:PRO:HA	4:A:717:GOL:H2	1.84	0.59
1:A:145:HIS:CD2	1:A:189:ARG:HD3	2.40	0.57
3:A:500:PQ0:N3	3:A:500:PQ0:N2	2.46	0.52
1:A:321:ALA:O	1:A:325:LYS:HD2	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:LYS:HB2	1:A:326:TRP:CE3	2.47	0.50
1:A:145:HIS:CG	1:A:189:ARG:HD3	2.46	0.50
1:A:106:TYR:HB3	5:A:750:HOH:O	2.12	0.50
1:A:372:GLN:OE1	1:A:376:ASP:OD1	2.30	0.49
1:A:303:ARG:HD2	5:A:913:HOH:O	2.15	0.47
1:A:280:ASP:OD2	4:A:703:GOL:H32	2.15	0.45
1:A:273:GLU:HA	4:A:716:GOL:H31	1.98	0.45
1:A:107:GLN:NE2	4:A:703:GOL:O2	2.50	0.45
1:A:191:GLU:OE1	4:A:705:GOL:O2	2.29	0.45
1:A:242:ARG:O	1:A:245:ASP:HB2	2.16	0.45
1:A:255:LYS:NZ	5:A:847:HOH:O	2.50	0.44
1:A:52:LYS:HD2	1:A:289:ARG:CZ	2.48	0.43
1:A:285:THR:O	1:A:289:ARG:HG3	2.19	0.42
1:A:21[A]:ARG:HD2	5:A:912:HOH:O	2.19	0.42
1:A:320:CYS:SG	1:A:322:VAL:HB	2.60	0.41
1:A:169:ALA:O	1:A:173:GLU:HG3	2.21	0.40
1:A:338:GLY:HA2	5:A:875:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/386 (92%)	349 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/315 (92%)	281 (97%)	9 (3%)	35	18

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	71	THR
1	A	85	LYS
1	A	110	SER
1	A	117	GLN
1	A	220	GLU
1	A	242	ARG
1	A	317	GLU
1	A	371	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	202	GLN
1	A	359	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	600	-	5,5,5	0.32	0	5,5,5	0.34	0
4	GOL	A	705	-	5,5,5	0.30	0	5,5,5	0.49	0
4	GOL	A	717	-	5,5,5	0.32	0	5,5,5	0.30	0
4	GOL	A	716	-	5,5,5	0.29	0	5,5,5	0.25	0
4	GOL	A	703	-	5,5,5	0.32	0	5,5,5	0.33	0
3	PQ0	A	500	-	14,14,14	9.46	12 (85%)	14,20,20	9.25	13 (92%)
4	GOL	A	706	-	5,5,5	0.31	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	600	-	-	2/4/4/4	-
4	GOL	A	705	-	-	0/4/4/4	-
4	GOL	A	717	-	-	2/4/4/4	-
4	GOL	A	716	-	-	4/4/4/4	-
4	GOL	A	703	-	-	0/4/4/4	-
3	PQ0	A	500	-	-	0/0/2/2	0/2/2/2
4	GOL	A	706	-	-	2/4/4/4	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	PQ0	C2-N3	16.42	1.73	1.33
3	A	500	PQ0	C10-N11	15.61	1.42	1.14
3	A	500	PQ0	O6-C6	15.02	1.52	1.23
3	A	500	PQ0	C4-N3	13.04	1.56	1.36
3	A	500	PQ0	C5-C4	-10.39	1.21	1.45
3	A	500	PQ0	C8-N9	8.95	1.54	1.36
3	A	500	PQ0	C10-C7	-6.69	1.31	1.43
3	A	500	PQ0	C2-N1	6.57	1.53	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	PQ0	C8-C7	5.73	1.44	1.38
3	A	500	PQ0	C4-N9	-3.87	1.28	1.36
3	A	500	PQ0	C6-N1	2.93	1.44	1.38
3	A	500	PQ0	C2-N2	-2.67	1.27	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	PQ0	C7-C8-N9	-14.53	91.34	107.45
3	A	500	PQ0	O6-C6-N1	-12.49	96.63	120.11
3	A	500	PQ0	N2-C2-N1	11.56	141.16	116.76
3	A	500	PQ0	N1-C2-N3	-10.84	103.48	123.32
3	A	500	PQ0	C5-C6-N1	10.27	143.39	115.10
3	A	500	PQ0	C5-C4-N9	9.86	123.81	107.89
3	A	500	PQ0	C7-C5-C4	-9.36	96.64	105.51
3	A	500	PQ0	C2-N1-C6	-8.91	108.95	125.11
3	A	500	PQ0	C7-C10-N11	-8.41	162.35	177.68
3	A	500	PQ0	C2-N3-C4	7.97	127.42	113.36
3	A	500	PQ0	N9-C4-N3	-5.81	118.92	125.99
3	A	500	PQ0	N2-C2-N3	-5.37	109.20	119.67
3	A	500	PQ0	O6-C6-C5	-3.26	115.91	125.69

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	600	GOL	C1-C2-C3-O3
4	A	600	GOL	O2-C2-C3-O3
4	A	706	GOL	O1-C1-C2-O2
4	A	706	GOL	O1-C1-C2-C3
4	A	716	GOL	O1-C1-C2-O2
4	A	716	GOL	O1-C1-C2-C3
4	A	716	GOL	C1-C2-C3-O3
4	A	717	GOL	O1-C1-C2-C3
4	A	716	GOL	O2-C2-C3-O3
4	A	717	GOL	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	705	GOL	2	0
4	A	717	GOL	1	0
4	A	716	GOL	1	0
4	A	703	GOL	2	0
3	A	500	PQ0	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	361/386 (93%)	0.04	14 (3%)	43 47	13, 20, 41, 62	2 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	116	LYS	3.9
1	A	105	GLY	3.7
1	A	106	TYR	3.3
1	A	134	MET	3.3
1	A	383	ALA	2.9
1	A	107	GLN	2.8
1	A	125	LYS	2.7
1	A	158	CYS	2.6
1	A	111	LEU	2.6
1	A	47	THR	2.3
1	A	286	ARG	2.2
1	A	382	PHE	2.2
1	A	104	GLY	2.1
1	A	72	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	703	6/6	0.70	0.15	31,67,71,74	0
4	GOL	A	717	6/6	0.71	0.19	40,58,64,82	0
4	GOL	A	716	6/6	0.78	0.17	27,46,60,62	0
4	GOL	A	706	6/6	0.79	0.15	33,51,59,74	0
4	GOL	A	705	6/6	0.88	0.12	23,42,45,46	0
4	GOL	A	600	6/6	0.89	0.09	22,26,32,33	0
3	PQ0	A	500	13/13	0.90	0.07	20,23,29,30	0
2	ZN	A	400	1/1	0.99	0.02	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.