



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 04:54 PM UTC

PDB ID : 5PWW / pdb_00005pww
Title : PanDDA analysis group deposition – Crystal Structure of SP100 after initial refinement with no ligand modelled (structure 20)
Authors : Pearce, N.M.; Krojer, T.; Talon, R.; Bradley, A.R.; Fairhead, M.; Sethi, R.; Wright, N.; MacLean, E.; Collins, P.; Brandao-Neto, J.; Douangamath, A.; Renjie, Z.; Dias, A.; Ng, J.; Brennan, P.E.; Cox, O.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; von Delft, F.
Deposited on : 2017-02-08
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	 6% 86% 13% ..
1	B	180	 12% 81% 17% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nuclear autoantigen Sp-100.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	10	0
			1536	971	268	278	19			
1	B	177	Total	C	N	O	S	0	7	0
			1480	940	258	263	19			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	312	Total	O	0	3
			312	312		
5	B	267	Total	O	0	0
			267	267		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nuclear autoantigen Sp-100

Chain A: 



- Molecule 1: Nuclear autoantigen Sp-100

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	127.70Å 45.38Å 83.33Å 90.00° 102.24° 90.00°	Depositor
Resolution (Å)	40.72 – 1.59 40.72 – 1.59	Depositor EDS
% Data completeness (in resolution range)	97.2 (40.72-1.59) 97.2 (40.72-1.59)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.59Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.174 , 0.205 0.183 , 0.214	Depositor DCC
R_{free} test set	3150 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3620	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.48	10/1603 (0.6%)	1.24	2/2154 (0.1%)
1	B	1.59	15/1538 (1.0%)	1.27	2/2071 (0.1%)
All	All	1.54	25/3141 (0.8%)	1.25	4/4225 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	837	MET	N-CA	7.50	1.55	1.46
1	B	830	VAL	N-CA	7.05	1.54	1.46
1	B	724	PHE	CA-C	-6.99	1.43	1.52
1	B	847	PHE	N-CA	-6.72	1.37	1.46
1	A	760	GLY	C-O	6.51	1.30	1.23
1	B	845	LYS	N-CA	6.48	1.54	1.46
1	B	784	LEU	N-CA	6.26	1.53	1.46
1	B	864	PHE	N-CA	5.99	1.53	1.46
1	B	843	ASN	N-CA	-5.98	1.39	1.46
1	B	801	TYR	CA-C	5.92	1.55	1.52
1	B	858	ILE	CA-CB	5.83	1.61	1.54
1	A	849	ARG	C-O	-5.82	1.19	1.24
1	A	741	SER	CA-C	5.79	1.59	1.52
1	A	730	ILE	CA-C	5.62	1.58	1.52
1	A	859	GLN	N-CA	-5.59	1.39	1.46
1	A	779	CYS	N-CA	5.41	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	855	ARG	C-O	5.29	1.30	1.24
1	B	853	PHE	N-CA	5.25	1.53	1.46
1	A	732	SER	N-CA	5.22	1.52	1.46
1	B	788	CYS	N-CA	5.21	1.52	1.46
1	B	868	PHE	C-O	-5.20	1.18	1.24
1	A	763	GLU	CD-OE1	-5.17	1.15	1.25
1	A	832	GLY	C-O	5.14	1.30	1.23
1	B	789	ASP	CA-C	-5.14	1.46	1.52
1	B	785	LYS	N-CA	5.02	1.52	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	849	ARG	CB-CA-C	-6.48	108.41	117.23
1	A	796	ALA	N-CA-C	5.73	117.33	111.14
1	B	756	GLU	N-CA-C	-5.56	99.46	108.41
1	B	832	GLY	O-C-N	5.34	127.37	122.19

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	755	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1497	13	0
1	B	1480	0	1414	13	0
2	A	3	0	0	0	0
2	B	2	0	0	0	0
3	A	12	0	12	3	0
4	A	4	0	6	2	0
4	B	4	0	6	1	0
5	A	312	0	0	12	1
5	B	267	0	0	7	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3620	0	2935	30	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:803[A]:ARG:NH2	5:A:1001:HOH:O	1.73	1.21
4:B:903:EDO:O2	5:B:1001:HOH:O	1.58	1.21
1:A:829[A]:ARG:NH1	5:A:1002:HOH:O	1.88	1.07
1:A:803[B]:ARG:NH1	5:A:1003:HOH:O	1.99	0.93
1:B:826[B]:MET:SD	5:B:1016:HOH:O	2.32	0.87
1:B:716[A]:CYS:SG	5:B:1170:HOH:O	2.35	0.83
3:A:904:MES:H81	5:A:1244:HOH:O	1.85	0.77
1:A:756:GLU:HG3	5:A:1153:HOH:O	1.82	0.76
1:B:826[B]:MET:HE2	5:B:1202:HOH:O	1.86	0.76
1:A:824:GLU:OE1	1:A:826[B]:MET:HE2	1.93	0.67
5:A:1001:HOH:O	1:B:773:PRO:HG2	1.95	0.67
1:A:802:ASN:O	3:A:904:MES:H32	1.99	0.62
1:B:750:ILE:HD11	5:B:1001:HOH:O	1.99	0.61
1:B:849:ARG:O	1:B:854[B]:THR:HG21	2.05	0.57
1:A:762:GLN:NE2	5:A:1007:HOH:O	2.39	0.55
1:A:810:GLN:NE2	5:A:1008:HOH:O	2.41	0.53
1:B:817:LYS:NZ	5:B:1006:HOH:O	2.42	0.53
1:B:757:SER:O	1:B:759:SER:N	2.39	0.51
1:A:803[A]:ARG:NE	3:A:904:MES:O1S	2.32	0.49
1:B:757:SER:C	1:B:759:SER:H	2.20	0.49
4:A:905:EDO:C2	5:A:1087:HOH:O	2.60	0.49
1:B:757:SER:C	1:B:759:SER:N	2.71	0.48
1:A:862[A]:ASP:OD1	5:A:1004:HOH:O	2.20	0.47
1:B:800:TYR:HE1	1:B:803:ARG:CB	2.28	0.46
1:A:849:ARG:O	1:A:854[B]:THR:HG21	2.16	0.46
1:B:800:TYR:CE1	1:B:803:ARG:CB	2.99	0.45
4:A:905:EDO:H21	5:A:1087:HOH:O	2.18	0.44
1:A:821:SER:HA	1:A:826[B]:MET:HE3	2.01	0.41
1:B:842:HIS:ND1	5:B:1004:HOH:O	2.37	0.41
1:A:748:LYS:HE3	5:A:1270:HOH:O	2.20	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1045:HOH:O	5:B:1045:HOH:O[2_556]	0.87	1.33
5:B:1201:HOH:O	5:B:1201:HOH:O[2_555]	0.98	1.22
5:A:1078:HOH:O	5:A:1199:HOH:O[1_565]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/180 (103%)	182 (98%)	4 (2%)	0	100	100
1	B	182/180 (101%)	173 (95%)	7 (4%)	2 (1%)	11	2
All	All	368/360 (102%)	355 (96%)	11 (3%)	2 (0%)	21	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	758	GLN
1	B	757	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/171 (104%)	177 (99%)	1 (1%)	78	66
1	B	165/171 (96%)	163 (99%)	2 (1%)	63	43
All	All	343/342 (100%)	340 (99%)	3 (1%)	78	55

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	738	ASN
1	B	862[A]	ASP
1	B	862[B]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	807	GLN
1	A	816	ASN
1	B	701	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	903	-	3,3,3	0.43	0	2,2,2	0.59	0
3	MES	A	904	-	12,12,12	1.18	1 (8%)	15,16,16	2.37	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	905	-	3,3,3	0.48	0	2,2,2	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	903	-	-	0/1/1/1	-
3	MES	A	904	-	-	1/6/14/14	0/1/1/1
4	EDO	A	905	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	904	MES	C8-S	-3.34	1.72	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	MES	C6-O1-C2	4.99	126.02	109.88
3	A	904	MES	C5-N4-C3	4.33	118.17	108.84
3	A	904	MES	C6-C5-N4	3.73	115.80	110.12
3	A	904	MES	C7-N4-C3	3.72	121.16	111.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	904	MES	C8-C7-N4-C3
4	A	905	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	903	EDO	1	0
3	A	904	MES	3	0
4	A	905	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/180 (98%)	0.18	11 (6%) 26 27	12, 22, 53, 102	10 (5%)
1	B	177/180 (98%)	0.52	22 (12%) 8 8	12, 24, 77, 133	7 (3%)
All	All	355/360 (98%)	0.35	33 (9%) 14 14	12, 23, 72, 133	17 (4%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	735	ALA	6.1
1	B	711	TRP	4.6
1	B	757	SER	4.6
1	B	800	TYR	4.5
1	B	759	SER	3.9
1	B	806	SER	3.8
1	B	756	GLU	3.7
1	B	803	ARG	3.4
1	A	878	SER	3.3
1	A	738	ASN	3.3
1	B	758	GLN	3.3
1	A	736	ASN	3.1
1	B	735	ALA	3.1
1	B	804	GLU	3.0
1	B	805	GLY	2.8
1	B	760	GLY	2.8
1	B	737	LYS	2.8
1	A	739	PRO	2.7
1	A	737	LYS	2.6
1	B	801	TYR	2.6
1	B	755	PRO	2.6
1	B	874	ILE	2.6
1	B	701	ASN	2.5
1	B	754	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	738	ASN	2.5
1	B	876	GLU	2.5
1	B	807	GLN	2.4
1	A	703	ASN	2.3
1	A	719	THR	2.3
1	B	736	ASN	2.1
1	A	849	ARG	2.1
1	A	701	ASN	2.1
1	A	715	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	903	1/1	0.76	0.20	95,95,95,95	1
4	EDO	A	905	4/4	0.80	0.15	45,55,60,68	0
2	ZN	A	902	1/1	0.94	0.06	29,29,29,29	0
3	MES	A	904	12/12	0.95	0.10	20,22,26,28	12
4	EDO	B	903	4/4	0.95	0.08	36,39,41,43	0
2	ZN	A	901	1/1	0.97	0.04	22,22,22,22	0
2	ZN	B	902	1/1	0.98	0.04	24,24,24,24	0
2	ZN	B	901	1/1	0.98	0.03	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.