



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 04:42 PM UTC

PDB ID : 8PWW / pdb_00008pww
Title : PfRH5 bound to monoclonal antibody MAD8-151
Authors : Farrell, B.; Higgins, M.K.
Deposited on : 2023-07-21
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	
1	C	338	
2	B	250	
2	D	250	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticulocyte-binding protein homolog 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	0	0
			2535	1632	429	459	15			
1	C	283	Total	C	N	O	S	0	1	0
			2403	1549	405	436	13			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	TYR	CYS	engineered mutation	UNP Q8IFM5
A	216	ALA	THR	engineered mutation	UNP Q8IFM5
A	250	ALA	THR	engineered mutation	UNP Q8IFM5
C	203	TYR	CYS	engineered mutation	UNP Q8IFM5
C	216	ALA	THR	engineered mutation	UNP Q8IFM5
C	250	ALA	THR	engineered mutation	UNP Q8IFM5

- Molecule 2 is a protein called scFv fragment for antibody MAD8-151.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	1	0
			1738	1097	286	348	7			
2	D	225	Total	C	N	O	S	0	1	0
			1734	1095	285	347	7			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	C	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	145	Total	O	0	0
			145	145		
6	B	162	Total	O	0	0
			162	162		
6	C	123	Total	O	0	0
			123	123		
6	D	103	Total	O	0	0
			103	103		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Reticulocyte-binding protein homolog 5




• Molecule 1: Reticulocyte-binding protein homolog 5



• Molecule 2: scFv fragment for antibody MAD8-151



• Molecule 2: scFv fragment for antibody MAD8-151





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.30Å 121.38Å 79.00Å 90.00° 90.18° 90.00°	Depositor
Resolution (Å)	79.00 – 1.95 79.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.5 (79.00-1.95) 96.6 (79.00-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.94Å)	Xtriage
Refinement program	BUSTER 2.10.4 (20-OCT-2021)	Depositor
R, R_{free}	0.230 , 0.252 (Not available) , 0.227	Depositor DCC
R_{free} test set	5117 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.456	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.014 for l,k,-h 0.257 for h,-k,-l 0.025 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8983	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EDO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	2/2588 (0.1%)	1.05	5/3473 (0.1%)
1	C	0.67	0/2454	1.06	4/3292 (0.1%)
2	B	0.74	0/1779	0.97	2/2420 (0.1%)
2	D	0.74	0/1772	0.98	3/2411 (0.1%)
All	All	0.71	2/8593 (0.0%)	1.02	14/11596 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	255	MET	SD-CE	-8.04	1.59	1.79
1	A	382	ILE	CG1-CD1	-7.86	1.21	1.51

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	135	ASP	CA-CB-CG	7.13	119.73	112.60
2	B	107	ASP	CA-CB-CG	6.77	119.37	112.60
1	A	452	LEU	N-CA-C	-6.76	100.47	110.48
2	D	107	ASP	CA-CB-CG	6.54	119.14	112.60
1	A	207	ASP	CA-CB-CG	6.29	118.89	112.60
1	C	207	ASP	CA-CB-CG	6.27	118.87	112.60
2	D	117	VAL	N-CA-CB	6.18	118.41	110.99
1	A	175	ILE	CA-C-N	5.55	125.75	120.43
1	A	175	ILE	C-N-CA	5.55	125.75	120.43
2	B	164	ASN	CA-CB-CG	5.34	117.94	112.60
1	C	175	ILE	CA-C-N	5.17	125.39	120.43
1	C	175	ILE	C-N-CA	5.17	125.39	120.43
1	A	164	LEU	CD1-CG-CD2	5.16	122.16	110.80
1	C	402	GLN	N-CA-C	-5.10	103.31	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2535	0	2557	21	0
1	C	2403	0	2430	24	0
2	B	1738	0	1695	13	0
2	D	1734	0	1687	18	0
3	A	4	0	6	0	0
3	C	4	0	6	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	D	6	0	8	0	0
5	A	7	0	10	0	0
5	C	7	0	10	0	0
6	A	145	0	0	0	0
6	B	162	0	0	1	0
6	C	123	0	0	1	0
6	D	103	0	0	0	0
All	All	8983	0	8425	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:238:LEU:HD21	1:C:254:MET:HB3	1.63	0.80
1:A:382:ILE:O	1:A:386:THR:HG23	1.88	0.73
1:C:257:GLU:O	1:C:261:LYS:HG2	1.89	0.73
2:B:23:THR:HG23	6:B:412:HOH:O	1.95	0.66
1:C:348:LEU:HD22	1:C:358:ILE:HD13	1.78	0.65
1:A:417:TYR:O	1:A:421:ILE:HG12	1.96	0.65
1:C:417:TYR:O	1:C:421:ILE:HG12	1.97	0.64
1:A:378:HIS:O	1:A:382:ILE:HG12	1.96	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:382:ILE:O	1:C:386:THR:HG23	1.97	0.64
1:C:378:HIS:O	1:C:382:ILE:HG13	1.99	0.62
1:C:238:LEU:HD23	1:C:254:MET:HE2	1.81	0.62
2:D:96:ALA:HB1	2:D:106:ILE:CG2	2.30	0.61
1:A:322:VAL:HG11	1:A:386:THR:HG21	1.85	0.59
2:B:37:ILE:CD1	2:B:47:TRP:HD1	2.16	0.59
2:D:4:LEU:CD2	2:D:24:VAL:HG22	2.33	0.58
1:A:266:ILE:HG22	1:A:270:LYS:HE2	1.86	0.57
2:D:37:ILE:CD1	2:D:47:TRP:HD1	2.17	0.57
2:B:37:ILE:HD11	2:B:47:TRP:HD1	1.69	0.57
2:D:37:ILE:HD11	2:D:47:TRP:HD1	1.70	0.57
1:A:267:LYS:HA	1:A:270:LYS:HE3	1.86	0.56
1:C:164:LEU:HD22	1:C:429:MET:CB	2.35	0.56
1:A:158:ALA:HB2	1:A:270:LYS:HD3	1.88	0.56
1:C:267:LYS:HA	1:C:270:LYS:HE3	1.86	0.56
1:A:164:LEU:HD22	1:A:429:MET:CB	2.36	0.55
1:C:266:ILE:HG22	1:C:270:LYS:HE2	1.87	0.55
2:B:138:MET:HE3	2:B:157:CYS:SG	2.47	0.55
2:D:138:MET:HE3	2:D:157:CYS:SG	2.46	0.55
1:C:158:ALA:HB2	1:C:270:LYS:HD3	1.88	0.54
1:A:164:LEU:HD22	1:A:429:MET:HB3	1.90	0.53
2:B:96:ALA:HB1	2:B:106:ILE:CG2	2.39	0.53
2:B:105:LEU:HD13	2:B:180:LEU:HD22	1.91	0.53
2:D:96:ALA:HB1	2:D:106:ILE:HG21	1.92	0.51
1:A:345:LEU:HD21	1:A:362:LYS:HG2	1.93	0.50
1:A:382:ILE:HD11	1:A:423:SER:C	2.35	0.50
1:C:164:LEU:HD22	1:C:429:MET:HB3	1.92	0.50
1:C:252:LYS:HD3	1:C:255:MET:HB3	1.92	0.50
2:D:105:LEU:HD13	2:D:180:LEU:HD22	1.92	0.50
2:D:39:GLN:HB3	2:D:92:VAL:CG1	2.42	0.49
1:C:345:LEU:HD21	1:C:362:LYS:HG2	1.95	0.49
1:A:405:GLU:O	1:A:409:ARG:HG2	2.13	0.48
1:C:284:LYS:O	1:C:288:THR:HG23	2.14	0.47
1:C:361:ILE:HG23	1:C:441:LEU:HD11	1.95	0.47
1:A:386:THR:HG22	1:A:420:PHE:CE1	2.49	0.47
1:C:367:GLU:HG3	6:C:602:HOH:O	2.15	0.47
1:C:405:GLU:O	1:C:409:ARG:HG2	2.15	0.47
2:B:159:ALA:O	2:B:203:THR:OG1	2.33	0.47
2:B:97:ARG:HB3	2:B:108[A]:SER:HB2	1.98	0.46
1:A:241:PRO:HG2	1:A:357:TYR:CE2	2.51	0.46
2:B:164:ASN:ND2	2:B:226:GLU:OE1	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:ASP:OD2	1:A:187:HIS:NE2	2.49	0.45
1:C:237:LYS:HG2	1:C:254:MET:HE3	1.99	0.44
1:C:164:LEU:HD22	1:C:429:MET:HB2	1.99	0.44
2:D:18:LEU:HD13	2:D:115:VAL:HG11	1.99	0.44
1:A:361:ILE:HG23	1:A:441:LEU:HD11	1.99	0.44
2:D:164:ASN:ND2	2:D:226:GLU:OE1	2.49	0.44
1:A:448:LEU:HD23	1:A:452:LEU:HD12	2.00	0.44
2:B:39:GLN:HB2	2:B:45:LEU:HD23	2.00	0.43
2:B:172:GLN:HB2	2:B:178:LEU:HG	2.01	0.43
2:D:39:GLN:HB2	2:D:45:LEU:HD23	2.01	0.43
1:C:238:LEU:HD21	1:C:254:MET:CB	2.40	0.43
2:D:39:GLN:HB3	2:D:92:VAL:HG13	2.01	0.42
2:D:78:PHE:CZ	2:D:95:CYS:HB2	2.55	0.42
1:C:318:LEU:HD13	1:C:421:ILE:CD1	2.49	0.42
1:C:241:PRO:HB2	1:C:357:TYR:CE1	2.55	0.42
1:A:157:ILE:O	1:A:161:ILE:HG13	2.20	0.41
2:B:78:PHE:CZ	2:B:95:CYS:HB2	2.56	0.41
1:A:206:VAL:O	1:A:210:ILE:HG12	2.21	0.41
1:A:403:LYS:O	1:A:407:LEU:HG	2.21	0.41
2:B:40:PRO:HB2	2:B:43:LYS:HD2	2.03	0.41
2:D:12:VAL:HG21	2:D:18:LEU:HD13	2.03	0.41
2:D:13:LYS:O	2:D:16:GLU:HB2	2.20	0.41
1:A:164:LEU:HD22	1:A:429:MET:HB2	2.03	0.41
2:D:37:ILE:CD1	2:D:47:TRP:CD1	3.02	0.41
1:C:183:ASP:OD2	1:C:187:HIS:NE2	2.49	0.41
2:D:4:LEU:HD21	2:D:34:TRP:HZ3	1.85	0.41
2:D:40:PRO:HB2	2:D:43:LYS:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/338 (88%)	292 (98%)	5 (2%)	0	100	100
1	C	278/338 (82%)	276 (99%)	2 (1%)	0	100	100
2	B	224/250 (90%)	216 (96%)	7 (3%)	1 (0%)	30	22
2	D	222/250 (89%)	214 (96%)	7 (3%)	1 (0%)	24	15
All	All	1021/1176 (87%)	998 (98%)	21 (2%)	2 (0%)	43	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	55	GLY
2	D	55	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/327 (88%)	280 (97%)	8 (3%)	38	25
1	C	275/327 (84%)	269 (98%)	6 (2%)	45	35
2	B	198/208 (95%)	193 (98%)	5 (2%)	42	29
2	D	197/208 (95%)	191 (97%)	6 (3%)	36	22
All	All	958/1070 (90%)	933 (97%)	25 (3%)	40	28

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164	LEU
1	A	263	LYS
1	A	308	ARG
1	A	322	VAL
1	A	406	LEU
1	A	438	GLU
1	A	442	ASN
1	A	454	MET
2	B	85	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	135	ASP
2	B	178	LEU
2	B	179	LYS
2	B	203	THR
1	C	164	LEU
1	C	252	LYS
1	C	342	GLU
1	C	406	LEU
1	C	438	GLU
1	C	442	ASN
2	D	92	VAL
2	D	106	ILE
2	D	113	MET
2	D	117	VAL
2	D	135	ASP
2	D	194	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	ASN
1	A	293	GLN
1	A	447	HIS
2	B	187	ASN
2	B	213	GLN
1	C	159	ASN
1	C	293	GLN
1	C	326	ASN
2	D	171	GLN
2	D	187	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	501	-	3,3,3	0.49	0	2,2,2	0.31	0
3	EDO	C	502	-	3,3,3	0.56	0	2,2,2	0.18	0
4	GOL	B	301	-	5,5,5	0.15	0	5,5,5	0.39	0
5	PEG	C	501	-	6,6,6	0.18	0	5,5,5	0.14	0
5	PEG	A	503	-	6,6,6	0.19	0	5,5,5	0.09	0
4	GOL	D	301	-	5,5,5	0.11	0	5,5,5	0.67	0
4	GOL	A	502	-	5,5,5	0.07	0	5,5,5	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	501	-	-	0/1/1/1	-
3	EDO	C	502	-	-	0/1/1/1	-
4	GOL	B	301	-	-	2/4/4/4	-
5	PEG	C	501	-	-	1/4/4/4	-
5	PEG	A	503	-	-	3/4/4/4	-
4	GOL	D	301	-	-	2/4/4/4	-
4	GOL	A	502	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	301	GOL	C1-C2-C3-O3
4	B	301	GOL	O2-C2-C3-O3
4	D	301	GOL	O1-C1-C2-O2
5	A	503	PEG	C1-C2-O2-C3
5	A	503	PEG	C4-C3-O2-C2
5	C	501	PEG	O2-C3-C4-O4
4	D	301	GOL	O2-C2-C3-O3
5	A	503	PEG	O2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/338 (88%)	-1.05	0 100 100	20, 33, 50, 70	0
1	C	283/338 (83%)	-0.99	0 100 100	18, 33, 56, 77	1 (0%)
2	B	227/250 (90%)	-1.10	0 100 100	18, 28, 42, 57	1 (0%)
2	D	225/250 (90%)	-1.11	0 100 100	18, 27, 40, 61	1 (0%)
All	All	1034/1176 (87%)	-1.06	0 100 100	18, 30, 50, 77	3 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	501	4/4	0.98	0.11	47,47,47,48	0
3	EDO	C	502	4/4	0.98	0.07	54,54,54,54	0
4	GOL	A	502	6/6	0.98	0.06	56,56,56,56	0
4	GOL	D	301	6/6	0.98	0.07	42,43,43,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PEG	A	503	7/7	0.98	0.06	54,55,55,55	0
5	PEG	C	501	7/7	0.98	0.06	51,51,52,52	0
4	GOL	B	301	6/6	0.99	0.05	47,47,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.