



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 10:07 AM UTC

PDB ID : 1PYW / pdb_00001pyw
Title : Human class II MHC protein HLA-DR1 bound to a designed peptide related to influenza virus hemagglutinin, FVKQNA(MAA)AL, in complex with staphylococcal enterotoxin C3 variant 3B2 (SEC3-3B2)
Authors : Zavala-Ruiz, Z.; Sundberg, E.J.; Stone, J.D.; DeOliveira, D.B.; Chan, I.C.; Svendsen, J.; Mariuzza, R.A.; Stern, L.J.
Deposited on : 2003-07-09
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	12% 77% 20% ..
2	B	190	6% 71% 25% .
3	C	10	10% 60% 30% 10%
4	D	239	7% 77% 17% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	0	0	0
			1470	952	239	274	5			

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DR-1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	190	Total	C	N	O	S	37	0	0
			1557	979	279	293	6			

- Molecule 3 is a protein called 9-residue influenza virus hemagglutinin related peptide FVKQNA(MAA)AL

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	0	0	0
			71	47	12	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	PHE	TYR	engineered mutation	GB 3212739
C	6	ALA	THR	engineered mutation	GB 3212739
C	7	MAA	LEU	engineered mutation	GB 3212739
C	8	ALA	LYS	engineered mutation	GB 3212739

- Molecule 4 is a protein called Enterotoxin type C-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	230	Total	C	N	O	S	18	0	0
			1884	1196	306	372	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	43	SER	LYS	engineered mutation	UNP P23313
D	45	PHE	LEU	engineered mutation	UNP P23313
D	46	LYS	ALA	engineered mutation	UNP P23313
D	47	TRP	HIS	engineered mutation	UNP P23313


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	135	Total 135	O 135	0	0
5	B	113	Total 113	O 113	0	0
5	C	12	Total 12	O 12	0	0
5	D	154	Total 154	O 154	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain




- Molecule 2: HLA class II histocompatibility antigen, DR-1 beta chain



- Molecule 3: 9-residue influenza virus hemagglutinin related peptide FVKQNA(MAA)AL



- Molecule 4: Enterotoxin type C-3





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	172.51Å 172.51Å 121.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.10 20.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.00-2.10) 98.2 (20.00-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.08Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.235 0.208 , 0.235	Depositor DCC
R_{free} test set	7813 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5396	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, MAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1515	0.89	3/2065 (0.1%)
2	B	0.37	0/1597	0.87	4/2168 (0.2%)
3	C	0.37	0/62	0.87	1/81 (1.2%)
4	D	0.36	0/1926	0.84	4/2592 (0.2%)
All	All	0.37	0/5100	0.86	12/6906 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	GLU	N-CA-C	-6.78	97.68	108.73
2	B	120	SER	N-CA-C	5.87	117.89	109.14
1	A	31	ILE	N-CA-C	-5.79	106.07	111.45
4	D	7	MET	N-CA-C	-5.50	103.19	110.40
4	D	81	VAL	N-CA-C	-5.42	99.94	108.23
3	C	4	GLN	N-CA-C	-5.25	100.74	109.46
1	A	144	LEU	N-CA-C	-5.13	101.90	109.86
4	D	225	SER	N-CA-C	5.04	117.50	111.71
2	B	164	VAL	CA-C-N	5.01	124.92	119.76
2	B	164	VAL	C-N-CA	5.01	124.92	119.76
2	B	34	GLN	N-CA-C	-5.01	105.55	113.02
4	D	24	MET	N-CA-C	-5.01	104.83	112.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1470	0	1406	28	0
2	B	1557	0	1488	52	0
3	C	71	0	76	3	0
4	D	1884	0	1811	33	0
5	A	135	0	0	3	0
5	B	113	0	0	0	0
5	C	12	0	0	0	0
5	D	154	0	0	1	0
All	All	5396	0	4781	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:ASN:HD22	2:B:154:THR:HG22	1.19	1.05
2:B:105:LYS:HE3	2:B:105:LYS:H	1.12	1.04
4:D:117:LYS:HE2	4:D:119:GLU:HB3	1.53	0.89
2:B:65:LYS:H	2:B:65:LYS:HE3	1.36	0.87
2:B:105:LYS:H	2:B:105:LYS:CE	1.92	0.83
1:A:94:ASN:HD22	1:A:104:VAL:HB	1.47	0.80
1:A:147:LYS:HE3	1:A:149:HIS:HE1	1.47	0.80
4:D:88:ASN:H	4:D:88:ASN:HD22	1.29	0.80
2:B:150:ASN:ND2	2:B:154:THR:HG22	1.96	0.79
4:D:215:MET:O	4:D:218:ASN:HB2	1.82	0.79
4:D:121:ASN:HD21	4:D:153:SER:H	1.31	0.78
2:B:105:LYS:HE3	2:B:105:LYS:N	1.98	0.72
2:B:65:LYS:H	2:B:65:LYS:CE	2.03	0.71
2:B:8:LEU:O	2:B:32:TYR:O	2.12	0.68
2:B:2:ASP:OD1	2:B:4:ARG:HD3	1.94	0.67
4:D:59:LYS:HA	4:D:59:LYS:HE2	1.79	0.65
2:B:152:ASP:OD1	2:B:154:THR:HB	1.97	0.64
1:A:81:PRO:HB3	2:B:5:PRO:HB2	1.82	0.62
5:A:187:HOH:O	4:D:46:LYS:HG2	1.99	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:145:THR:HG22	2:B:158:LEU:H	1.65	0.62
1:A:16:PRO:HD2	2:B:6:ARG:HD3	1.81	0.62
2:B:150:ASN:HD22	2:B:154:THR:CG2	2.06	0.61
4:D:88:ASN:H	4:D:88:ASN:ND2	1.99	0.60
2:B:2:ASP:CG	2:B:6:ARG:HH22	2.10	0.60
2:B:94:ARG:HH11	2:B:94:ARG:HG3	1.66	0.59
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.86	0.58
1:A:95:SER:HB2	1:A:96:PRO:HD2	1.85	0.58
1:A:98:GLU:HB2	1:A:101:GLU:HB3	1.87	0.56
2:B:27:LEU:HG	2:B:29:ARG:HD2	1.86	0.56
2:B:65:LYS:HE3	2:B:65:LYS:N	2.15	0.56
4:D:83:ASP:OD1	4:D:118:HIS:HD2	1.89	0.55
2:B:114:LEU:HD21	2:B:160:MET:HB3	1.86	0.55
2:B:18:PHE:HB2	2:B:23:ARG:HB3	1.87	0.55
4:D:122:HIS:O	4:D:152:LYS:HE3	2.05	0.55
2:B:94:ARG:HG3	2:B:94:ARG:NH1	2.22	0.54
4:D:37:LYS:HG3	4:D:81:VAL:CG1	2.38	0.54
4:D:2:SER:HB2	4:D:195:THR:H	1.73	0.54
1:A:143:HIS:HD2	2:B:12:LYS:NZ	2.06	0.53
1:A:128:VAL:HG12	1:A:129:THR:N	2.23	0.53
4:D:58:LEU:N	4:D:58:LEU:HD22	2.23	0.53
1:A:38:LYS:O	4:D:212:LYS:HE2	2.09	0.52
1:A:76:ARG:HH22	2:B:57:ASP:CG	2.18	0.50
1:A:12:PHE:C	1:A:12:PHE:CD1	2.90	0.50
1:A:17:ASP:O	1:A:18:GLN:HB2	2.12	0.50
1:A:67:LYS:O	1:A:71:GLU:HG3	2.12	0.50
1:A:182:ALA:HA	2:B:105:LYS:HG2	1.93	0.49
1:A:94:ASN:ND2	1:A:104:VAL:HB	2.21	0.48
2:B:52:GLU:OE1	2:B:55:ARG:HD2	2.13	0.48
4:D:121:ASN:ND2	4:D:153:SER:H	2.05	0.47
2:B:103:PRO:HD3	2:B:188:TRP:CZ3	2.50	0.47
2:B:61:TRP:CZ2	3:C:9:LEU:HD23	2.50	0.47
2:B:116:VAL:HG22	2:B:160:MET:HG3	1.97	0.46
4:D:64:VAL:HG22	4:D:109:THR:HG22	1.97	0.46
2:B:23:ARG:HH11	2:B:23:ARG:HG2	1.80	0.46
1:A:76:ARG:NH2	2:B:57:ASP:OD2	2.46	0.45
2:B:2:ASP:OD1	2:B:6:ARG:NH2	2.49	0.45
1:A:70:LEU:HD13	2:B:9:TRP:HB2	1.98	0.45
4:D:231:GLU:OE1	4:D:233:HIS:HE1	1.98	0.45
4:D:6:PRO:HB3	4:D:197:TRP:CZ2	2.51	0.45
4:D:37:LYS:HG3	4:D:81:VAL:HG11	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:88:ASN:HD22	4:D:88:ASN:N	2.05	0.44
4:D:226:LYS:HG2	5:D:286:HOH:O	2.16	0.44
2:B:105:LYS:CE	2:B:105:LYS:N	2.69	0.44
1:A:110:ASP:OD1	1:A:146:ARG:HG2	2.18	0.44
4:D:2:SER:CB	4:D:195:THR:H	2.30	0.44
1:A:14:LEU:HD11	2:B:6:ARG:HB3	1.99	0.44
1:A:130:THR:HG23	5:A:292:HOH:O	2.16	0.44
2:B:134:ASN:ND2	2:B:170:VAL:H	2.15	0.44
4:D:129:GLN:NE2	4:D:226:LYS:HB3	2.33	0.43
4:D:216:MET:C	4:D:218:ASN:H	2.25	0.43
2:B:74:ALA:O	2:B:78:TYR:HB3	2.19	0.43
4:D:55:ASP:OD1	4:D:60:ASN:HB2	2.19	0.43
2:B:130:ARG:HD3	2:B:137:GLU:CD	2.43	0.43
3:C:6:ALA:HA	3:C:7:MAA:HM1	1.53	0.43
1:A:140:ARG:HG3	1:A:146:ARG:HG3	2.00	0.43
4:D:121:ASN:HD21	4:D:153:SER:N	2.08	0.42
2:B:23:ARG:HH11	2:B:23:ARG:CG	2.33	0.42
1:A:122:LEU:HD23	1:A:127:PRO:HA	2.01	0.42
1:A:36:MET:HE2	1:A:36:MET:HA	2.02	0.42
2:B:57:ASP:CG	3:C:9:LEU:HD22	2.45	0.42
2:B:143:VAL:HG13	2:B:160:MET:HB2	2.02	0.42
4:D:231:GLU:HB3	4:D:233:HIS:CE1	2.55	0.41
1:A:105:LEU:HG	1:A:153:PHE:CE1	2.54	0.41
2:B:2:ASP:OD1	2:B:4:ARG:CD	2.64	0.41
4:D:6:PRO:HB3	4:D:197:TRP:CH2	2.55	0.41
4:D:94:TYR:O	4:D:95:PHE:HB3	2.21	0.41
4:D:125:ASN:O	4:D:127:ASN:N	2.53	0.41
2:B:104:SER:HB3	2:B:107:GLN:HE21	1.86	0.41
1:A:17:ASP:CG	2:B:6:ARG:HH11	2.29	0.41
1:A:49:GLY:HA3	5:A:293:HOH:O	2.20	0.41
2:B:118:SER:HA	2:B:158:LEU:HD23	2.02	0.41
2:B:166:ARG:HB2	2:B:166:ARG:HH11	1.86	0.41
2:B:133:ARG:O	2:B:134:ASN:HB2	2.21	0.41
4:D:1:GLU:HG3	4:D:2:SER:N	2.36	0.41
1:A:82:ILE:HG13	2:B:33:ASN:HB3	2.02	0.40
2:B:87:GLU:HG2	2:B:92:GLN:NE2	2.35	0.40
2:B:104:SER:HB3	2:B:107:GLN:NE2	2.35	0.40
1:A:143:HIS:HD2	2:B:12:LYS:HZ1	1.68	0.40
2:B:166:ARG:CB	2:B:166:ARG:NH1	2.85	0.40
4:D:54:SER:HA	4:D:62:ASP:HA	2.02	0.40
4:D:64:VAL:HG22	4:D:109:THR:CG2	2.51	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:91:VAL:O	4:D:92:ASN:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/182 (97%)	173 (98%)	4 (2%)	0	100	100
2	B	188/190 (99%)	178 (95%)	6 (3%)	4 (2%)	5	2
3	C	7/10 (70%)	7 (100%)	0	0	100	100
4	D	226/239 (95%)	217 (96%)	7 (3%)	2 (1%)	14	10
All	All	598/621 (96%)	575 (96%)	17 (3%)	6 (1%)	12	9

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	108	PRO
2	B	110	GLN
2	B	111	HIS
2	B	32	TYR
4	D	57	LYS
4	D	126	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/166 (98%)	161 (99%)	2 (1%)	63	72
2	B	171/171 (100%)	162 (95%)	9 (5%)	20	19
3	C	6/6 (100%)	6 (100%)	0	100	100
4	D	212/220 (96%)	208 (98%)	4 (2%)	50	58
All	All	552/563 (98%)	537 (97%)	15 (3%)	39	45

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLU
1	A	101	GLU
2	B	29	ARG
2	B	53	LEU
2	B	65	LYS
2	B	68	LEU
2	B	105	LYS
2	B	109	LEU
2	B	110	GLN
2	B	114	LEU
2	B	154	THR
4	D	37	LYS
4	D	56	LYS
4	D	88	ASN
4	D	218	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	143	HIS
1	A	149	HIS
2	B	64	GLN
2	B	81	HIS
2	B	92	GLN
2	B	113	ASN
2	B	134	ASN
2	B	150	ASN
2	B	156	GLN
4	D	60	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	88	ASN
4	D	92	ASN
4	D	118	HIS
4	D	121	ASN
4	D	125	ASN
4	D	127	ASN
4	D	148	GLN
4	D	157	GLN
4	D	177	ASN
4	D	194	ASN
4	D	218	ASN
4	D	233	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MAA	C	7	3	4,5,6	2.17	1 (25%)	2,5,7	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAA	C	7	3	-	1/2/4/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	7	MAA	CM-N	-4.16	1.36	1.46

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	7	MAA	CB-CA-N-CM

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	7	MAA	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/182 (98%)	0.19	21 (11%) 9 9	15, 26, 61, 84	0
2	B	186/190 (97%)	0.28	12 (6%) 25 26	14, 30, 55, 79	0
3	C	8/10 (80%)	-0.21	1 (12%) 8 8	18, 21, 24, 37	0
4	D	228/239 (95%)	0.03	16 (7%) 22 24	17, 28, 53, 70	0
All	All	601/621 (96%)	0.15	50 (8%) 17 18	14, 28, 57, 84	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	ALA	7.9
4	D	239	GLY	5.2
1	A	180	PHE	5.1
2	B	106	THR	4.9
2	B	108	PRO	4.6
4	D	177	ASN	4.2
1	A	99	LEU	4.1
1	A	101	GLU	4.0
1	A	100	ARG	4.0
1	A	128	VAL	3.8
4	D	178	SER	3.8
1	A	181	ASP	3.6
2	B	145	THR	3.6
2	B	1	GLY	3.4
1	A	97	VAL	3.3
4	D	106	GLY	3.3
4	D	58	LEU	3.3
1	A	157	THR	3.3
4	D	125	ASN	3.3
1	A	158	GLU	3.2
4	D	7	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	55	ASP	3.1
1	A	160	VAL	3.0
3	C	9	LEU	3.0
1	A	96	PRO	3.0
4	D	123	PHE	3.0
1	A	94	ASN	2.8
2	B	164	VAL	2.8
1	A	159	ASP	2.7
2	B	113	ASN	2.7
2	B	136	GLN	2.7
2	B	162	GLU	2.7
1	A	130	THR	2.6
1	A	131	GLY	2.6
4	D	1	GLU	2.6
2	B	139	LYS	2.5
4	D	218	ASN	2.5
1	A	108	PHE	2.4
1	A	57	GLN	2.3
4	D	124	ASP	2.3
4	D	96	SER	2.2
4	D	176	PHE	2.2
1	A	154	LEU	2.2
2	B	143	VAL	2.2
4	D	238	ASN	2.2
1	A	102	PRO	2.1
2	B	105	LYS	2.1
1	A	98	GLU	2.1
2	B	19	ASN	2.1
4	D	236	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAA	C	7	6/7	0.97	0.05	17,19,19,20	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.