



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 08:37 PM UTC

PDB ID : 4PYS / pdb_00004pys
Title : The crystal structure of beta-N-acetylhexosaminidase from Bacteroides fragilis NCTC 9343
Authors : Tan, K.; Hatzos-Skintges, C.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2014-03-27
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	506	10% 88% 9% .
1	B	506	9% 86% 9% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	A	607	-	-	X	-
4	FMT	B	608	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-N-acetylhexosaminidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	492	Total	C	N	O	S	Se	0	1	0
			3972	2564	675	713	8	12			
1	B	483	Total	C	N	O	S	Se	0	4	0
			3909	2517	674	698	8	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	ASN	TYR	conflict	UNP Q5LAT3
A	15	ALA	SER	conflict	UNP Q5LAT3
B	14	ASN	TYR	conflict	UNP Q5LAT3
B	15	ALA	SER	conflict	UNP Q5LAT3

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	280	Total	O	0	0
			280	280		
5	B	324	Total	O	0	0
			324	324		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: beta-N-acetylhexosaminidase



- Molecule 1: beta-N-acetylhexosaminidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.82Å 96.82Å 315.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.03 – 1.82 34.03 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.6 (34.03-1.82) 99.6 (34.03-1.82)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 1.82Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.165 , 0.190 0.164 , 0.189	Depositor DCC
R_{free} test set	6740 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8553	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9169e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/4078	0.74	0/5512
1	B	0.40	0/4016	0.73	0/5423
All	All	0.40	0/8094	0.74	0/10935

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3972	0	3798	29	0
1	B	3909	0	3766	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	24	0	32	1	0
3	B	12	0	16	0	0
4	A	15	0	5	2	0
4	B	15	0	5	2	0
5	A	280	0	0	2	0
5	B	324	0	0	2	0
All	All	8553	0	7622	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:LEU:HD21	1:A:263:GLN:HA	1.69	0.74
1:B:388[A]:ARG:NH2	5:B:1022:HOH:O	2.23	0.71
1:A:21:GLN:HG3	1:A:22:ILE:H	1.61	0.64
1:B:203:VAL:HB	1:B:261:VAL:HG21	1.82	0.60
1:A:275:LYS:NZ	1:A:315:ASP:OD2	2.37	0.57
1:A:192:ARG:NH2	5:A:780:HOH:O	2.39	0.55
1:A:104:GLU:OE2	3:A:605:GOL:H12	2.06	0.54
1:A:44:ARG:HB3	1:A:71:CYS:SG	2.47	0.54
1:A:338:ARG:NH2	1:A:362:TYR:OH	2.40	0.54
1:B:192[A]:ARG:HD3	1:B:193:TYR:CZ	2.43	0.53
1:B:148:MSE:HG3	1:B:177:HIS:CG	2.44	0.53
1:A:20:GLY:N	1:A:121:TRP:O	2.41	0.52
1:A:148:MSE:HG3	1:A:177:HIS:CG	2.44	0.52
1:A:307:LYS:HD2	1:A:310:TRP:CE3	2.44	0.52
1:B:356:VAL:HG23	1:B:357:ILE:HG23	1.93	0.50
1:A:364:LEU:HB3	1:A:368:VAL:HG21	1.93	0.50
1:A:383:LEU:O	1:A:387:VAL:HG13	2.13	0.49
1:A:505:PRO:HD3	1:A:515:TYR:CZ	2.48	0.49
1:B:247:ASN:ND2	1:B:259:VAL:O	2.45	0.49
1:B:36:ARG:NH2	1:B:125:GLU:OE1	2.46	0.49
1:A:250:PRO:HB3	1:A:257:VAL:O	2.15	0.46
1:A:252:LEU:HD11	1:A:281:PHE:CZ	2.51	0.46
1:A:148:MSE:HE1	1:A:448:TRP:CD2	2.50	0.46
1:B:388[A]:ARG:NH1	5:B:920:HOH:O	2.47	0.46
1:A:252:LEU:HD21	1:A:281:PHE:CD2	2.51	0.46
1:B:444:SER:HB2	4:B:608:FMT:C	2.47	0.45
1:B:360:ASP:OD1	1:B:362:TYR:N	2.50	0.45
1:B:148:MSE:HE1	1:B:448:TRP:CD2	2.53	0.44
1:A:432:ARG:NH2	1:A:436:GLU:OE1	2.39	0.44
1:A:59:TYR:HA	1:A:63:VAL:O	2.17	0.44
1:A:307:LYS:HA	1:A:307:LYS:HD3	1.62	0.44
1:A:435:GLU:O	1:A:436:GLU:HB3	2.17	0.43
1:A:306:PRO:HD3	5:A:970:HOH:O	2.18	0.43
1:A:382:ALA:CB	4:A:607:FMT:H	2.49	0.43
1:B:177:HIS:ND1	4:B:608:FMT:O2	2.52	0.42
1:B:505:PRO:HD3	1:B:515:TYR:CZ	2.55	0.42
1:B:24:MSE:HA	1:B:25:PRO:C	2.45	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ILE:HD12	1:B:63:VAL:HG21	2.02	0.42
1:B:44:ARG:HB3	1:B:71:CYS:SG	2.60	0.41
1:B:59:TYR:HA	1:B:63:VAL:O	2.21	0.41
1:B:240:GLY:HA2	1:B:270:ILE:HG22	2.02	0.41
1:B:364:LEU:HB3	1:B:368:VAL:HG21	2.03	0.41
1:A:342:TYR:O	1:A:345:GLN:HG2	2.21	0.41
1:A:382:ALA:HB2	4:A:607:FMT:H	2.01	0.41
1:B:92:TYR:CE2	1:B:138:ASP:HB3	2.55	0.41
1:A:150:ASP:HA	1:A:179:HIS:HB3	2.03	0.41
1:A:192:ARG:NH2	1:B:503:PHE:H	2.19	0.41
1:A:255:PHE:HB2	1:A:257:VAL:HG13	2.02	0.41
1:B:82:LYS:O	1:B:104:GLU:HA	2.21	0.41
1:B:218:VAL:O	1:B:222:ILE:HG12	2.20	0.41
1:B:301:GLY:HA2	1:B:353:TRP:CD1	2.56	0.41
1:B:195:PHE:HB2	1:B:249:TYR:CE2	2.56	0.40
1:B:511:ALA:O	1:B:515:TYR:HB2	2.21	0.40
1:B:195:PHE:HB2	1:B:249:TYR:CZ	2.57	0.40
1:A:38:ARG:HH21	1:A:130:ILE:HD13	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	487/506 (96%)	471 (97%)	15 (3%)	1 (0%)	43	33
1	B	475/506 (94%)	460 (97%)	14 (3%)	1 (0%)	43	33
All	All	962/1012 (95%)	931 (97%)	29 (3%)	2 (0%)	43	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	435	GLU
1	B	314	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/425 (95%)	403 (100%)	2 (0%)	81	77
1	B	403/425 (95%)	399 (99%)	4 (1%)	68	60
All	All	808/850 (95%)	802 (99%)	6 (1%)	76	70

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	341	ASP
1	A	513	THR
1	B	75	VAL
1	B	228	ARG
1	B	333	LEU
1	B	363	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	389	HIS
1	A	478	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	602	-	5,5,5	0.37	0	5,5,5	0.29	0
4	FMT	A	607	-	2,2,2	0.81	0	1,1,1	0.27	0
4	FMT	B	608	-	2,2,2	0.63	0	1,1,1	0.13	0
4	FMT	A	608	-	2,2,2	0.70	0	1,1,1	0.26	0
4	FMT	B	604	-	2,2,2	0.80	0	1,1,1	0.27	0
4	FMT	B	605	-	2,2,2	0.87	0	1,1,1	0.21	0
4	FMT	A	610	-	2,2,2	0.68	0	1,1,1	0.13	0
4	FMT	A	609	-	2,2,2	0.68	0	1,1,1	0.15	0
4	FMT	B	607	-	2,2,2	0.69	0	1,1,1	0.23	0
4	FMT	A	606	-	2,2,2	0.66	0	1,1,1	0.29	0
4	FMT	B	606	-	2,2,2	0.67	0	1,1,1	0.14	0
3	GOL	A	603	-	5,5,5	0.37	0	5,5,5	0.62	0
3	GOL	A	605	-	5,5,5	0.33	0	5,5,5	0.31	0
3	GOL	B	603	-	5,5,5	0.40	0	5,5,5	0.29	0
3	GOL	A	604	-	5,5,5	0.33	0	5,5,5	0.50	0
3	GOL	B	602	-	5,5,5	0.39	0	5,5,5	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	602	-	-	0/4/4/4	-
3	GOL	A	603	-	-	0/4/4/4	-
3	GOL	A	605	-	-	2/4/4/4	-
3	GOL	B	603	-	-	0/4/4/4	-
3	GOL	A	604	-	-	0/4/4/4	-
3	GOL	B	602	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	605	GOL	O1-C1-C2-C3
3	A	605	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	607	FMT	2	0
4	B	608	FMT	2	0
3	A	605	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	480/506 (94%)	0.37	53 (11%) 10 12	16, 34, 89, 118	1 (0%)
1	B	471/506 (93%)	0.22	47 (9%) 12 14	16, 32, 81, 104	4 (0%)
All	All	951/1012 (93%)	0.29	100 (10%) 11 12	16, 33, 85, 118	5 (0%)

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	259	VAL	8.6
1	A	255	PHE	7.7
1	A	260	LYS	7.6
1	A	309	ASN	7.4
1	A	262	PRO	6.4
1	A	20	GLY	6.4
1	A	310	TRP	6.3
1	B	260	LYS	6.0
1	B	257	VAL	5.6
1	B	317	ARG	5.4
1	B	320	ILE	4.8
1	B	316	CYS	4.8
1	B	321	GLU	4.7
1	B	322	LYS	4.6
1	B	325	LEU	4.6
1	A	311	ASP	4.5
1	B	330	ASP	4.5
1	A	306	PRO	4.4
1	A	317	ARG	4.4
1	B	274	GLY	4.3
1	B	313	CYS	4.2
1	B	362	TYR	4.2
1	A	513	THR	4.1
1	A	435	GLU	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	314	PRO	4.1
1	B	261	VAL	4.0
1	A	261	VAL	4.0
1	B	258	ALA	4.0
1	A	316	CYS	3.7
1	A	270	ILE	3.7
1	B	259	VAL	3.7
1	B	254	CYS	3.6
1	B	127	ARG	3.5
1	B	329	HIS	3.5
1	A	356	VAL	3.5
1	A	254	CYS	3.5
1	B	479	PRO	3.5
1	B	256	ASN	3.4
1	A	308	GLY	3.4
1	A	258	ALA	3.4
1	A	273	ALA	3.4
1	B	363	SER	3.3
1	A	257	VAL	3.3
1	B	273	ALA	3.3
1	A	127	ARG	3.2
1	B	306	PRO	3.2
1	A	21	GLN	3.1
1	A	320	ILE	3.0
1	A	302	GLY	3.0
1	B	360	ASP	3.0
1	A	271	PHE	2.9
1	B	335	PHE	2.9
1	B	270	ILE	2.9
1	A	256	ASN	2.8
1	A	322	LYS	2.8
1	A	272	CYS	2.8
1	A	291	ARG	2.8
1	A	314	PRO	2.8
1	A	312	LYS	2.8
1	A	303	ASP	2.8
1	A	124	GLY	2.7
1	A	387	VAL	2.7
1	A	263	GLN	2.6
1	B	514	ASN	2.6
1	B	478	ASN	2.6
1	B	272	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	SER	2.5
1	A	34	ASP	2.5
1	B	315	ASP	2.5
1	A	514	ASN	2.4
1	B	78	HIS	2.4
1	B	40	GLN	2.4
1	B	334	TRP	2.4
1	B	55	ILE	2.3
1	B	389	HIS	2.3
1	A	274	GLY	2.3
1	B	328	SER	2.3
1	A	71	CYS	2.3
1	A	436	GLU	2.3
1	B	253	GLY	2.3
1	A	360	ASP	2.3
1	B	124	GLY	2.2
1	B	318	SER	2.2
1	B	275	LYS	2.2
1	A	434	ARG	2.2
1	B	125	GLU	2.2
1	A	361	GLY	2.2
1	A	269	ASN	2.2
1	A	328	SER	2.2
1	A	248	ALA	2.2
1	A	363	SER	2.2
1	A	251	ARG	2.2
1	B	53	SER	2.1
1	A	313	CYS	2.1
1	B	513	THR	2.1
1	A	195	PHE	2.1
1	B	512	GLY	2.1
1	B	252	LEU	2.1
1	B	269	ASN	2.0
1	A	55	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	605	6/6	0.71	0.20	44,51,62,64	0
4	FMT	A	610	3/3	0.83	0.15	32,32,48,50	0
4	FMT	A	608	3/3	0.86	0.12	47,47,53,66	0
3	GOL	B	602	6/6	0.86	0.16	29,46,52,52	0
4	FMT	B	606	3/3	0.87	0.18	43,43,44,52	0
3	GOL	A	603	6/6	0.88	0.14	32,42,46,49	0
4	FMT	A	609	3/3	0.89	0.17	39,39,41,46	0
4	FMT	B	608	3/3	0.90	0.32	40,40,44,45	0
4	FMT	B	607	3/3	0.91	0.14	38,38,44,52	0
2	ZN	A	601	1/1	0.93	0.13	66,66,66,66	1
2	ZN	B	601	1/1	0.93	0.08	91,91,91,91	1
3	GOL	A	602	6/6	0.93	0.08	28,29,33,35	0
4	FMT	B	605	3/3	0.96	0.07	22,22,30,35	0
3	GOL	B	603	6/6	0.96	0.09	29,33,36,36	0
3	GOL	A	604	6/6	0.96	0.08	26,31,33,34	0
4	FMT	B	604	3/3	0.96	0.08	24,24,31,33	0
4	FMT	A	606	3/3	0.97	0.08	22,22,33,34	0
4	FMT	A	607	3/3	0.97	0.07	21,21,35,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.