



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 12:29 AM UTC

PDB ID : 4PYV / pdb\_00004pyv  
Title : Crystal structure of renin in complex with compound4  
Authors : Ostermann, N.; Zink, F.  
Deposited on : 2014-03-28  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	 4% 88% 10% ..
1	B	340	 2% 86% 10% ..

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Renin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	Total 2578	C 1648	N 417	O 499	S 14	0	0	0
1	B	334	Total 2570	C 1642	N 416	O 498	S 14	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 3 is tert-butyl {(3S,5R)-5-[cyclopropyl(2,3-dichlorobenzyl)carbamoyl]piperidin-3-yl}carbamate (CCD ID: 2XF) (formula:  $C_{21}H_{29}Cl_2N_3O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
3	A	1	Total	C	Cl	N	O	0	0
			29	21	2	3	3		
3	B	1	Total	C	Cl	N	O	0	0
			29	21	2	3	3		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			S
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			S
4	B	1	4	2	1	1	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	61	61	61	0	0
6	B	77	77	77	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Renin



- Molecule 1: Renin



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.90Å 141.90Å 141.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.78 – 2.65 42.78 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.78-2.65) 99.6 (42.78-2.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 2.65Å)	Xtrriage
Refinement program	BUSTER 2.11.4	Depositor
R, $R_{free}$	0.201 , 0.241 0.201 , 0.242	Depositor DCC
$R_{free}$ test set	2781 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.9	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.023 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5375	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2XF, NAG, DMS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/2637	1.21	6/3575 (0.2%)
1	B	0.84	0/2629	1.23	6/3564 (0.2%)
All	All	0.84	0/5266	1.22	12/7139 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	32	ASP	CA-CB-CG	6.78	119.38	112.60
1	A	32	ASP	CA-CB-CG	6.60	119.20	112.60
1	B	291	ILE	CB-CA-C	5.92	116.45	110.94
1	A	265	LYS	N-CA-C	5.88	118.29	110.53
1	B	112	PHE	CA-CB-CG	-5.79	108.02	113.80
1	A	264	GLY	CA-C-N	5.58	131.13	120.97
1	A	264	GLY	C-N-CA	5.58	131.13	120.97
1	A	291	ILE	CB-CA-C	5.51	116.07	110.94
1	A	116	GLU	N-CA-C	-5.48	105.33	112.23
1	B	180	HIS	N-CA-C	-5.42	100.87	109.59
1	B	221	ILE	N-CA-C	-5.37	100.85	108.58
1	B	248	LYS	N-CA-C	-5.29	102.02	109.96

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2578	0	2510	8	0
1	B	2570	0	2500	15	0
2	A	14	0	13	0	0
3	A	29	0	29	1	0
3	B	29	0	29	1	0
4	A	4	0	6	0	0
4	B	8	0	12	0	0
5	A	5	0	0	0	0
6	A	61	0	0	0	0
6	B	77	0	0	0	0
All	All	5375	0	5099	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:LYS:HD3	1:B:261:HIS:HD2	1.47	0.80
1:B:195:LYS:HD3	1:B:261:HIS:CD2	2.20	0.76
1:B:277:GLN:HG2	1:B:280:TYR:CE2	2.21	0.75
1:B:4:VAL:HB	1:B:165:ILE:HG23	1.74	0.70
1:B:277:GLN:HG2	1:B:280:TYR:CZ	2.30	0.67
1:B:150:VAL:HG12	1:B:314:ASP:HA	1.82	0.61
1:A:6:LEU:HD21	1:A:165:ILE:HD12	1.90	0.53
1:A:275:VAL:HG22	1:A:284:LEU:HD22	1.92	0.50
1:B:195:LYS:HB2	1:B:261:HIS:CD2	2.46	0.49
1:B:0:THR:HB	1:B:145:VAL:O	2.12	0.49
1:A:0:THR:HB	1:A:145:VAL:O	2.14	0.47
1:A:88:ILE:HG23	1:A:95:THR:HG23	1.98	0.46
1:A:41:PRO:HB2	1:A:55:LEU:HD23	1.98	0.46
1:A:150:VAL:HG12	1:A:314:ASP:HA	1.98	0.45
1:B:88:ILE:HG23	1:B:95:THR:HG23	1.99	0.45
3:B:1000:2XF:C14	3:B:1000:2XF:H8	2.46	0.45
1:B:41:PRO:HB2	1:B:55:LEU:HD23	2.00	0.44
1:B:69:THR:HB	1:B:84:LEU:HD12	2.00	0.44
1:B:195:LYS:HB2	1:B:261:HIS:HD2	1.84	0.42
1:A:100:MET:HB3	1:A:133:VAL:CG1	2.49	0.42
1:B:221:ILE:HG13	1:B:304:THR:HB	2.01	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:ALA:HB1	1:B:107:MET:HE2	2.02	0.41
1:B:205:LEU:HD21	1:B:230:LYS:HB2	2.03	0.41
1:A:217:GLY:C	3:A:1001:2XF:H6	2.47	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/340 (97%)	325 (98%)	5 (2%)	1 (0%)	36	52
1	B	330/340 (97%)	325 (98%)	3 (1%)	2 (1%)	21	34
All	All	661/680 (97%)	650 (98%)	8 (1%)	3 (0%)	24	39

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	240	ARG
1	A	203	THR
1	B	68	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/290 (98%)	265 (94%)	18 (6%)	16	28
1	B	282/290 (97%)	264 (94%)	18 (6%)	16	28
All	All	565/580 (97%)	529 (94%)	36 (6%)	16	28

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	THR
1	A	17	GLU
1	A	36	SER
1	A	47	ARG
1	A	47(A)	LEU
1	A	47(B)	TYR
1	A	59	SER
1	A	90	THR
1	A	204	LEU
1	A	233	GLU
1	A	240	ARG
1	A	248	LYS
1	A	250	ASN
1	A	254	THR
1	A	269	LEU
1	A	286	ILE
1	A	291	ILE
1	A	308	LYS
1	B	0	THR
1	B	36	SER
1	B	47	ARG
1	B	47(A)	LEU
1	B	47(B)	TYR
1	B	59	SER
1	B	80	VAL
1	B	107	MET
1	B	165	ILE
1	B	195	LYS
1	B	204	LEU
1	B	233	GLU
1	B	250	ASN
1	B	269	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	277	GLN
1	B	284	LEU
1	B	289	MET
1	B	291	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	A	174	HIS
1	A	180	HIS
1	A	250	ASN
1	B	143	GLN
1	B	250	ASN
1	B	277	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DMS	A	1002	-	3,3,3	0.42	0	3,3,3	0.69	0
4	DMS	B	1002	-	3,3,3	0.14	0	3,3,3	0.43	0
2	NAG	A	1000	1	14,14,15	0.45	0	17,19,21	0.96	1 (5%)
3	2XF	A	1001	-	29,31,31	1.53	6 (20%)	40,45,45	2.29	11 (27%)
5	SO4	A	1003	-	4,4,4	0.28	0	6,6,6	0.28	0
4	DMS	B	1001	-	3,3,3	0.22	0	3,3,3	0.58	0
3	2XF	B	1000	-	29,31,31	1.33	4 (13%)	40,45,45	2.08	16 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2XF	B	1000	-	-	1/25/37/37	0/3/3/3
2	NAG	A	1000	1	-	1/6/23/26	0/1/1/1
3	2XF	A	1001	-	-	0/25/37/37	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	2XF	C13-C11	3.04	1.55	1.48
3	A	1001	2XF	C14-C11	2.91	1.55	1.48
3	B	1000	2XF	C14-C11	2.80	1.54	1.48
3	B	1000	2XF	C13-C11	2.72	1.54	1.48
3	A	1001	2XF	C20-C15	2.51	1.42	1.39
3	A	1001	2XF	C27-C26	2.36	1.57	1.51
3	A	1001	2XF	C28-C26	2.18	1.57	1.51
3	A	1001	2XF	C6-C1	2.08	1.56	1.53
3	B	1000	2XF	C27-C26	2.08	1.57	1.51
3	B	1000	2XF	C20-C15	2.07	1.41	1.39

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	2XF	C15-C12-N8	6.93	123.56	113.14
3	A	1001	2XF	C4-N3-C2	5.78	118.49	111.78
3	B	1000	2XF	O9-C7-C5	-5.57	112.67	120.86
3	A	1001	2XF	O9-C7-C5	-4.52	114.22	120.86
3	B	1000	2XF	C26-O25-C23	3.94	126.95	120.97
3	B	1000	2XF	C2-C1-N10	3.83	114.97	109.98

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	2XF	C6-C5-C4	-3.79	106.01	109.79
3	A	1001	2XF	C1-N10-C23	-3.66	116.59	122.23
3	A	1001	2XF	C4-C5-C7	3.60	116.83	110.07
3	B	1000	2XF	C4-C5-C7	3.43	116.51	110.07
3	B	1000	2XF	C4-N3-C2	3.25	115.55	111.78
3	B	1000	2XF	C13-C11-N8	3.17	122.40	118.46
2	A	1000	NAG	C1-O5-C5	3.15	116.40	112.19
3	A	1001	2XF	C12-C15-C20	-3.07	116.66	121.67
3	B	1000	2XF	C15-C20-CL1	2.97	122.19	119.16
3	B	1000	2XF	C16-C15-C20	2.71	119.46	117.33
3	A	1001	2XF	O25-C26-C27	2.53	117.18	107.21
3	B	1000	2XF	C15-C12-N8	2.52	116.93	113.14
3	A	1001	2XF	C6-C1-N10	2.47	114.68	110.62
3	B	1000	2XF	C6-C5-C7	-2.44	106.07	110.42
3	A	1001	2XF	O25-C23-N10	2.38	113.93	110.03
3	B	1000	2XF	C5-C6-C1	-2.30	108.70	110.82
3	B	1000	2XF	C20-C19-CL2	-2.25	118.44	120.51
3	B	1000	2XF	O25-C26-C27	2.23	116.00	107.21
3	B	1000	2XF	C1-N10-C23	-2.23	118.81	122.23
3	B	1000	2XF	C12-N8-C11	2.20	120.52	118.34
3	A	1001	2XF	C16-C15-C20	2.19	119.05	117.33
3	B	1000	2XF	C12-C15-C20	-2.13	118.19	121.67

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	NAG	O5-C5-C6-O6
3	B	1000	2XF	C27-C26-O25-C23


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	2XF	1	0
3	B	1000	2XF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/340 (98%)	0.39	15 (4%) 38 31	44, 66, 100, 121	0
1	B	334/340 (98%)	0.21	8 (2%) 59 53	41, 62, 91, 129	0
All	All	669/680 (98%)	0.30	23 (3%) 48 40	41, 63, 95, 129	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	159	SER	8.9
1	B	160(C)	GLN	6.5
1	A	159	SER	6.2
1	A	160(C)	GLN	4.9
1	B	47(A)	LEU	4.1
1	A	204	LEU	3.6
1	A	287	HIS	2.9
1	A	-3	LEU	2.8
1	B	47(B)	TYR	2.8
1	A	207	GLU	2.6
1	A	158	ASP	2.3
1	A	47(B)	TYR	2.3
1	A	295	THR	2.3
1	A	76	SER	2.2
1	A	47(A)	LEU	2.2
1	B	52	TYR	2.2
1	A	202	SER	2.2
1	A	211	LEU	2.1
1	B	47	ARG	2.1
1	B	-2	GLY	2.1
1	A	200	GLY	2.1
1	A	297	PRO	2.0
1	B	131	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	2XF	A	1001	29/29	0.77	0.21	69,88,99,102	0
2	NAG	A	1000	14/15	0.79	0.13	83,87,93,97	0
5	SO4	A	1003	5/5	0.91	0.20	101,103,104,105	0
3	2XF	B	1000	29/29	0.95	0.10	48,60,76,81	0
4	DMS	B	1002	4/4	0.96	0.12	72,75,77,78	0
4	DMS	A	1002	4/4	0.97	0.11	58,63,66,72	0
4	DMS	B	1001	4/4	0.98	0.09	51,54,54,55	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 2XF A 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 2XF B 1000:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.