



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 02:33 AM UTC

PDB ID : 6PYS / pdb_00006pys
Title : Human PI3Kalpha in complex with Compound 2-10 ((3S)-3-benzyl-3-methyl-5-[5-(2-methylpyrimidin-5-yl)pyrazolo[1,5-a]pyrimidin-3-yl]-1,3-dihydro-2H-indol-2-one)
Authors : Lesburg, C.A.; Augustin, M.A.
Deposited on : 2019-07-30
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	945	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	890	7346	4686	1263	1335	62	268	7	0

- Molecule 2 is (3S)-3-benzyl-3-methyl-5-[5-(2-methylpyrimidin-5-yl)pyrazolo[1,5-a]pyrimidin-3-yl]-1,3-dihydro-2H-indol-2-one (CCD ID: P5J) (formula: C₂₇H₂₂N₆O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	34	27	6	1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	1	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	162	Total	O	0	4
			166	166		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.44Å 133.91Å 141.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.33 – 2.19 97.33 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.2 (97.33-2.19) 99.2 (97.33-2.19)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.74 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.224 , 0.259 0.227 , 0.252	Depositor DCC
R_{free} test set	1498 reflections (2.58%)	wwPDB-VP
Wilson B-factor (Å ²)	47.9	Xtriage
Anisotropy	0.449	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7558	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, P5J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	1/7508 (0.0%)	0.80	2/10140 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	352	ASP	CB-CG	-5.69	1.37	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	352	ASP	CA-CB-CG	6.79	119.39	112.60
1	A	1012	GLU	CB-CG-CD	6.17	123.10	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7346	0	7304	15	0
2	A	34	0	0	0	0
3	A	12	0	16	0	0
4	A	166	0	0	0	0
All	All	7558	0	7320	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459[B]:ILE:CG2	1:A:641[B]:TYR:HE1	2.11	0.63
1:A:459[B]:ILE:HG22	1:A:459[B]:ILE:O	1.99	0.61
1:A:459[B]:ILE:CG2	1:A:641[B]:TYR:CE1	2.85	0.59
1:A:459[B]:ILE:HG23	1:A:641[B]:TYR:HE1	1.69	0.57
1:A:459[B]:ILE:HG23	1:A:641[B]:TYR:CE1	2.39	0.57
1:A:357:ARG:NH1	1:A:372:ASN:OD1	2.46	0.46
1:A:459[B]:ILE:HG22	1:A:641[B]:TYR:CE1	2.51	0.46
1:A:459[B]:ILE:CG2	1:A:459[B]:ILE:O	2.64	0.45
1:A:353:LYS:HG2	1:A:377:PRO:HA	1.98	0.45
1:A:376:VAL:HG13	1:A:377:PRO:HD2	1.99	0.45
1:A:641[B]:TYR:HE2	1:A:1007:GLY:HA3	1.82	0.44
1:A:376:VAL:CG1	1:A:380:ASN:O	2.67	0.43
1:A:459[B]:ILE:HG22	1:A:641[B]:TYR:HE1	1.80	0.42
1:A:376:VAL:CG1	1:A:377:PRO:HD2	2.49	0.42
1:A:376:VAL:HG13	1:A:377:PRO:CD	2.50	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	881/945 (93%)	856 (97%)	23 (3%)	2 (0%)	43 51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	352	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	956	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	821/859 (96%)	820 (100%)	1 (0%)	88 95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	937	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	384	ASN
1	A	530	GLN
1	A	630	GLN
1	A	670	HIS
1	A	676	HIS
1	A	885	ASN
1	A	1000	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	1103	-	5,5,5	0.52	0	5,5,5	0.55	0
3	GOL	A	1102	-	5,5,5	0.32	0	5,5,5	0.38	0
2	P5J	A	1101	-	39,39,39	0.76	0	55,58,58	1.96	18 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1103	-	-	0/4/4/4	-
3	GOL	A	1102	-	-	0/4/4/4	-
2	P5J	A	1101	-	-	5/13/29/29	0/6/6/6

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	P5J	C8-N9-N4	5.55	109.24	103.81
2	A	1101	P5J	C7-C5-N6	3.93	133.64	125.03
2	A	1101	P5J	C1-N6-C5	3.86	121.73	117.78
2	A	1101	P5J	C27-C16-C20	3.75	114.15	109.24
2	A	1101	P5J	C16-C14-C13	-3.38	106.25	109.26
2	A	1101	P5J	C33-N32-C31	3.17	121.27	116.07
2	A	1101	P5J	C28-C33-N32	-3.04	119.12	123.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	P5J	C29-N30-C31	3.03	121.04	116.07
2	A	1101	P5J	C28-C29-N30	-2.82	119.43	123.59
2	A	1101	P5J	C3-N4-N9	2.82	131.20	127.01
2	A	1101	P5J	C12-C13-C14	-2.77	119.33	121.92
2	A	1101	P5J	C7-C8-N9	-2.66	109.89	113.29
2	A	1101	P5J	C28-C1-N6	2.57	119.68	116.94
2	A	1101	P5J	C27-C16-C14	-2.57	107.13	111.42
2	A	1101	P5J	C15-C14-C16	2.47	133.52	130.58
2	A	1101	P5J	N4-C5-N6	-2.42	119.05	121.96
2	A	1101	P5J	N32-C31-N30	-2.34	121.66	125.46
2	A	1101	P5J	C14-C16-C17	2.29	103.24	101.83

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	P5J	C14-C16-C20-C21
2	A	1101	P5J	N6-C1-C28-C29
2	A	1101	P5J	N6-C1-C28-C33
2	A	1101	P5J	C2-C1-C28-C33
2	A	1101	P5J	C2-C1-C28-C29

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	890/945 (94%)	0.92	123 (13%) 6 5	20, 55, 90, 117	94 (10%)

All (123) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	TYR	5.9
1	A	459[A]	ILE	5.3
1	A	347	ASN	5.2
1	A	301	CYS	4.8
1	A	351	ILE	4.3
1	A	1051	TRP	4.1
1	A	377	PRO	3.9
1	A	299	MET	3.8
1	A	307	TYR	3.8
1	A	378	CYS	3.8
1	A	175	PRO	3.7
1	A	452	LEU	3.6
1	A	940	HIS	3.6
1	A	181	ILE	3.5
1	A	788	ILE	3.5
1	A	199	SER	3.5
1	A	182	TYR	3.4
1	A	198	VAL	3.4
1	A	178	PRO	3.4
1	A	409	VAL	3.3
1	A	213	HIS	3.3
1	A	247	GLN	3.2
1	A	176	GLU	3.1
1	A	338	ILE	3.1
1	A	295	SER	3.1
1	A	521	ASN	3.1
1	A	177	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	742	PRO	3.0
1	A	376	VAL	3.0
1	A	387	LEU	3.0
1	A	322	THR	3.0
1	A	773	SER	2.9
1	A	265	TYR	2.9
1	A	973	LYS	2.9
1	A	889	ILE	2.9
1	A	453	GLU	2.8
1	A	180	HIS	2.8
1	A	863	LYS	2.8
1	A	356	VAL	2.8
1	A	520	ASP	2.8
1	A	956	LEU	2.8
1	A	479	TRP	2.8
1	A	475	LEU	2.7
1	A	374	GLN	2.7
1	A	484	VAL	2.7
1	A	697	MET	2.7
1	A	229	THR	2.7
1	A	522	GLU	2.7
1	A	384	ASN	2.6
1	A	949	ARG	2.6
1	A	373	THR	2.6
1	A	972	THR	2.6
1	A	473	LEU	2.6
1	A	267	LEU	2.6
1	A	352	ASP	2.5
1	A	381	PRO	2.5
1	A	936	HIS	2.5
1	A	232	MET	2.5
1	A	383	TRP	2.5
1	A	220	VAL	2.4
1	A	721	GLN	2.4
1	A	723	LYS	2.4
1	A	309	ARG	2.4
1	A	183	ASN	2.4
1	A	346	VAL	2.4
1	A	898	THR	2.4
1	A	775	ALA	2.4
1	A	935	GLY	2.4
1	A	145	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	483	VAL	2.4
1	A	353	LYS	2.4
1	A	860	ILE	2.4
1	A	292	SER	2.4
1	A	185	LEU	2.3
1	A	281	ARG	2.3
1	A	375	ARG	2.3
1	A	330	ILE	2.3
1	A	344	VAL	2.3
1	A	448	VAL	2.3
1	A	331	ASN	2.3
1	A	324	THR	2.3
1	A	735	LEU	2.3
1	A	449	PRO	2.2
1	A	967	GLY	2.2
1	A	156	LEU	2.2
1	A	392	TYR	2.2
1	A	386	TRP	2.2
1	A	977	PHE	2.2
1	A	354	ILE	2.2
1	A	955	VAL	2.2
1	A	451	GLY	2.2
1	A	379	SER	2.2
1	A	577[A]	ARG	2.2
1	A	894	ILE	2.2
1	A	881	LEU	2.2
1	A	107	ASN	2.2
1	A	1048	HIS	2.1
1	A	341	ALA	2.1
1	A	215	CYS	2.1
1	A	306	SER	2.1
1	A	965	SER	2.1
1	A	214	ASP	2.1
1	A	355	TYR	2.1
1	A	964	ILE	2.1
1	A	979	ARG	2.1
1	A	297	LEU	2.1
1	A	939	ASP	2.1
1	A	1029	ASP	2.1
1	A	207	TYR	2.1
1	A	225	ILE	2.1
1	A	287	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	422	LEU	2.0
1	A	526	ASN	2.0
1	A	390	ASP	2.0
1	A	266	PRO	2.0
1	A	720	LYS	2.0
1	A	724	LYS	2.0
1	A	718	ILE	2.0
1	A	192	VAL	2.0
1	A	1036	LEU	2.0
1	A	852	ARG	2.0
1	A	407	CYS	2.0
1	A	248	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	1102	6/6	0.81	0.15	64,73,78,82	0
3	GOL	A	1103	6/6	0.85	0.23	22,34,34,38	6
2	P5J	A	1101	34/34	0.90	0.12	52,61,80,87	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around P5J A 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.