



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:10 PM UTC

PDB ID : 8PYM / pdb_00008pym
Title : Human IGF1R with inhibitor 54
Authors : Dreyer, M.K.; Wang, J.; Elkins, J.M.
Deposited on : 2023-07-25
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	321	9% 66% 19% • 13%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Insulin-like growth factor 1 receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	278	Total	C	N	O	S	0	0	0
			2213	1406	372	413	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	973	MET	-	initiating methionine	UNP P08069
AAA	1287	ALA	-	expression tag	UNP P08069
AAA	1288	GLU	-	expression tag	UNP P08069
AAA	1289	ASN	-	expression tag	UNP P08069
AAA	1290	LEU	-	expression tag	UNP P08069
AAA	1291	TYR	-	expression tag	UNP P08069
AAA	1292	PHE	-	expression tag	UNP P08069
AAA	1293	GLN	-	expression tag	UNP P08069

- Molecule 2 is 5,5-dimethyl-1-(1H-pyrrolo[2,3-b]pyridin-4-ylmethyl)-3-[4-(trifluoromethylsulfanyl)phenyl]imidazolidine-2,4-dione (CCD ID: IED) (formula: C₂₀H₁₇F₃N₄O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	AAA	1	Total	C	F	N	O	S	
			30	20	3	4	2	1	0

- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	1	Total	Ni		
			1	1	0	0

- Molecule 4 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	Cd		
			1	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	47	Total	O		
			47	47	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Insulin-like growth factor 1 receptor beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	69.35Å 69.35Å 142.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.39 – 2.65 62.39 – 2.65	Depositor EDS
% Data completeness (in resolution range)	96.2 (62.39-2.65) 96.2 (62.39-2.65)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.236 , 0.309 0.237 , 0.297	Depositor DCC
R_{free} test set	486 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2292	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, IED, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.07	0/2258	1.47	4/3044 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	AAA	1116	ALA	CA-C-N	5.53	126.09	120.00
1	AAA	1116	ALA	C-N-CA	5.53	126.09	120.00
1	AAA	1082	MET	CA-C-N	5.41	127.80	120.38
1	AAA	1082	MET	C-N-CA	5.41	127.80	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2213	0	2198	29	0
2	AAA	30	0	0	1	0
3	AAA	1	0	0	0	0
4	AAA	1	0	0	0	0
5	AAA	47	0	0	1	0
All	All	2292	0	2198	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:1066:LEU:HB2	1:AAA:1078:ILE:HG22	1.73	0.69
1:AAA:1113:ILE:HD11	1:AAA:1240:MET:HE1	1.73	0.69
1:AAA:1179:MET:HE2	1:AAA:1184:LEU:HD23	1.76	0.68
1:AAA:995:ALA:HB1	1:AAA:997:GLU:OE1	1.96	0.66
1:AAA:1074:PRO:O	1:AAA:1076:LEU:HD13	2.00	0.61
1:AAA:1069:VAL:HB	1:AAA:1076:LEU:HB2	1.83	0.59
1:AAA:1023:LYS:O	1:AAA:1024:ASP:HB2	2.05	0.56
1:AAA:985:ASP:OD2	1:AAA:985:ASP:N	2.39	0.55
1:AAA:1073:GLN:HB2	1:AAA:1074:PRO:HD3	1.88	0.53
1:AAA:1250:GLN:HB2	1:AAA:1256:ARG:HG3	1.93	0.51
1:AAA:1223:ARG:O	1:AAA:1227:GLU:HG3	2.10	0.51
1:AAA:1127:ASN:CG	1:AAA:1191:THR:HG22	2.36	0.51
1:AAA:1196:TRP:CE3	1:AAA:1249:TRP:HA	2.46	0.50
1:AAA:1114:GLN:HE21	1:AAA:1114:GLN:HB3	1.53	0.48
1:AAA:1247:MET:O	1:AAA:1250:GLN:HG2	2.12	0.48
1:AAA:1095:ARG:HB2	1:AAA:1096:PRO:HD3	1.96	0.47
1:AAA:1158:ARG:HB3	1:AAA:1160:ILE:HD12	1.97	0.46
1:AAA:1167:ARG:HG3	1:AAA:1173:LEU:HD23	1.97	0.46
1:AAA:1113:ILE:HD13	1:AAA:1113:ILE:HA	1.81	0.46
1:AAA:1058:ASN:CG	1:AAA:1058:ASN:O	2.59	0.45
1:AAA:1054:MET:HE1	1:AAA:1079:MET:HE1	2.00	0.44
1:AAA:1145:GLU:HA	1:AAA:1145:GLU:OE1	2.17	0.44
1:AAA:1177:ARG:HB2	5:AAA:1431:HOH:O	2.17	0.43
1:AAA:1133:HIS:O	1:AAA:1134:ARG:HG3	2.18	0.43
1:AAA:1073:GLN:H	1:AAA:1073:GLN:HG2	1.67	0.42
1:AAA:1146:ASP:O	1:AAA:1147:PHE:HB2	2.20	0.41
1:AAA:1054:MET:HE1	1:AAA:1079:MET:CE	2.49	0.41
1:AAA:1142:MET:HE2	2:AAA:1301:IED:C1	2.50	0.41
1:AAA:1065:LEU:HD12	1:AAA:1078:ILE:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	274/321 (85%)	246 (90%)	27 (10%)	1 (0%)	30	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	1169	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	242/280 (86%)	217 (90%)	25 (10%)	7	10

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	985	ASP
1	AAA	1000	THR
1	AAA	1001	MET
1	AAA	1002	SER
1	AAA	1005	LEU
1	AAA	1009	SER
1	AAA	1029	ARG
1	AAA	1033	LYS
1	AAA	1035	VAL
1	AAA	1036	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	AAA	1041	MET
1	AAA	1043	GLU
1	AAA	1071	GLN
1	AAA	1073	GLN
1	AAA	1089	SER
1	AAA	1111	LYS
1	AAA	1113	ILE
1	AAA	1114	GLN
1	AAA	1154	PHE
1	AAA	1168	LYS
1	AAA	1171	LYS
1	AAA	1214	GLN
1	AAA	1255	MET
1	AAA	1264	SER
1	AAA	1266	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IED	AAA	1301	-	32,33,33	1.88	10 (31%)	45,51,51	2.00	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IED	AAA	1301	-	-	0/13/37/37	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	1301	IED	C16-N14	-6.38	1.34	1.44
2	AAA	1301	IED	C13-N14	-3.30	1.33	1.40
2	AAA	1301	IED	C1-C9	-3.21	1.33	1.44
2	AAA	1301	IED	C19-S22	-3.14	1.73	1.78
2	AAA	1301	IED	C15-N14	-2.52	1.35	1.40
2	AAA	1301	IED	C8-N7	-2.28	1.33	1.38
2	AAA	1301	IED	C15-N11	-2.23	1.33	1.37
2	AAA	1301	IED	C6-N7	-2.11	1.33	1.36
2	AAA	1301	IED	C8-C9	-2.09	1.32	1.36
2	AAA	1301	IED	C1-C2	-2.01	1.39	1.42

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	1301	IED	C1-C6-N5	-6.45	121.35	126.40
2	AAA	1301	IED	C1-C6-N7	-4.94	105.52	108.33
2	AAA	1301	IED	N7-C6-N5	3.97	133.12	126.69
2	AAA	1301	IED	C29-C12-N11	-3.89	106.81	111.48
2	AAA	1301	IED	O28-C15-N11	-3.25	122.10	125.70
2	AAA	1301	IED	C6-C1-C9	3.15	107.96	106.40
2	AAA	1301	IED	C23-S22-C19	3.15	106.94	99.30
2	AAA	1301	IED	C29-C12-C13	3.10	113.79	107.89
2	AAA	1301	IED	C6-N7-C8	2.39	109.49	108.11
2	AAA	1301	IED	C10-N11-C15	2.38	125.80	121.80

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	1301	IED	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	AAA	278/321 (86%)	0.66	28 (10%)	12 10	18, 39, 82, 95	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	1096	PRO	5.5
1	AAA	1106	PRO	4.6
1	AAA	984	ALA	4.6
1	AAA	1107	PRO	3.7
1	AAA	986	VAL	3.6
1	AAA	1026	PRO	3.3
1	AAA	1024	ASP	3.0
1	AAA	1270	MET	2.9
1	AAA	1071	GLN	2.9
1	AAA	1023	LYS	2.8
1	AAA	1074	PRO	2.6
1	AAA	1040	SER	2.6
1	AAA	985	ASP	2.6
1	AAA	1022	VAL	2.5
1	AAA	1003	ARG	2.5
1	AAA	1035	VAL	2.5
1	AAA	995	ALA	2.5
1	AAA	1090	TYR	2.4
1	AAA	999	ILE	2.4
1	AAA	1021	VAL	2.3
1	AAA	1169	GLY	2.3
1	AAA	1240	MET	2.2
1	AAA	1038	ALA	2.2
1	AAA	1039	ALA	2.1
1	AAA	1045	ILE	2.1
1	AAA	1000	THR	2.1
1	AAA	1017	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	1160	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IED	AAA	1301	30/30	0.94	0.08	23,27,30,31	0
3	NI	AAA	1302	1/1	0.99	0.03	29,29,29,29	0
4	CD	AAA	1303	1/1	1.00	0.02	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IED AAA 1301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.