



Full wwPDB EM Validation Report ⓘ

Mar 5, 2026 – 07:53 PM UTC

PDB ID : 6PZ9 / pdb_00006pz9
EMDB ID : EMD-20528
Title : Cryo-EM structure of the pancreatic beta-cell SUR1 bound to ATP and repaglinide
Authors : Shyng, S.L.; Yoshioka, C.; Martin, G.M.; Sung, M.W.
Deposited on : 2019-07-31
Resolution : 3.65 Å(reported)
Based on initial model : 6BAA

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11564 (3.15 - 4.15)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	1582	
2	D	390	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7773 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-binding cassette sub-family C member 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	C	1177	7612	4844	1385	1357	26	0	0


- Molecule 2 is a protein called ATP-sensitive inward rectifier potassium channel 11.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	D	19	97	59	19	19	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	191	PRO	LEU	conflict	UNP P70673

- Molecule 3 is Repaglinide (CCD ID: BJX) (formula: $C_{27}H_{36}N_2O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
3	C	1	Total	C	N	O	0
			33	27	2	4	

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					AltConf
4	C	1	Total 31	C 10	N 5	O 13	P 3	0



ARG	TRP	LEU	PRO	ILE	THR
ARG	GLY	VAL	ARG	GLY	LEU
SER	GLN	ALA	HIS	PHE	LEU
VAL	ARG	PRO	GLY	GLY	ILE
VAL	PHE	LEU	ARG	ARG	PHE
ALA	VAL	ILE	CYS	MET	THR
LYS	PRO	ILE	LEU	ARG	MET
ALA	ILE	TYR	PHE	VAL	SER
LYS	VAL	HIS	MET	THR	PHE
PRO	ALA	VAL	LEU	GLU	LEU
LYS	GLU	ILE	ARG	GLU	LEU
PHE	GLU	ASP	VAL	CYS	CYS
SER	ASP	SER	GLY	PRO	SER
ILE	GLY	ASN	ASP	LEU	THR
SER	ARG	SER	LEU	LEU	GLU
PRO	TYR	PRO	ARG	ILE	PRO
ASP	SER	LEU	LYS	LEU	ARG
LEU	VAL	TYR	MET	ILE	ALA
SER	ASP	ASP	SER	LEU	THR
	TYR	LEU	ILE	ILE	ARG
	SER	ALA	ILE	VAL	THR
	LYS	PRO	SER	GLN	GLU
	PHE	SER	ALA	ASN	ARG
	GLY	ASP	THR	ILE	ALA
	ASN	LEU	ILE	VAL	ALA
	THR	HIS	HIS	GLY	ARG
	VAL	HIS	MET	LEU	PHE
	LYS	HIS	GLN	MET	VAL
	VAL	GLN	VAL	ILE	VAL
	PRO	ASP	VAL	ASN	LYS
	THR	LEU	ARG	ALA	GLY
	PRO	GLU	LYS	ILE	ASN
	LEU	ILE	THR	GLY	CYS
	CYS	ILE	THR	LEU	ASN
	THR	VAL	SER	GLY	VAL
	ALA	ILE	PRO	CYS	THR
	ARG	LEU	GLY	PHE	ASN
	GLN	GLY	GLU	VAL	HIS
	LEU	GLY	GLU	MET	VAL
	ASP	VAL	VAL	LYS	PRO
	GLU	VAL	VAL	THR	ASN
	ASP	GLU	PRO	VAL	ILE
	ARG	THR	LEU	ALA	ARG
	SER	THR	HIS	ALA	GLU
	LEU	GLY	GLN	HIS	ILE
	LEU	ILE	VAL	ARG	GLY
	ASP	THR	ASP	ARG	PHE
	ALA	THR	ILE	ALA	LEU
	LEU	GLN	PRO	THR	GLN
	THR	ARG	MET	THR	SER
	LEU	ARG	GLU	LEU	PHE
	ALA	THR	ASN	ILE	THR
	SER	SER	GLY	PHE	THR
	SER	TYR	VAL	SER	THR
	ARG	LEU	GLY	LYS	LEU
	GLY	ALA	GLY	HIS	VAL
	PRO	ASP	ASN	ALA	ASP
	LEU	GLU	ILE	VAL	LEU
	ARG	ILE	ILE	THR	LYS
	LYS	LEU	PHE	THR	TRP
					PRO

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	312000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.161	Depositor
Minimum map value	-0.108	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.0245	Depositor
Map size ($\text{\AA}$)	136.89499, 126.44499, 178.69499	wwPDB
Map dimensions	131, 121, 171	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.045, 1.045, 1.045	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BJX, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.39	0/7757	0.67	0/10666
2	D	0.36	0/96	0.98	1/132 (0.8%)
All	All	0.39	0/7853	0.67	1/10798 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	5	LYS	N-CA-C	-6.99	101.94	110.88

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1331	ALA	Peptide
1	C	1528	PHE	Peptide
1	C	280	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	7612	0	6135	48	0
2	D	97	0	52	1	0
3	C	33	0	0	0	0
4	C	31	0	12	0	0
All	All	7773	0	6199	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:ARG:NH2	1:C:437:ASN:OD1	2.31	0.64
1:C:768:GLY:O	1:C:1218:ARG:NH1	2.32	0.62
1:C:1217:PHE:HB2	1:C:1219:TYR:HB2	1.81	0.62
1:C:253:LEU:HD23	1:C:1236:ILE:HD13	1.81	0.62
1:C:263:TYR:OH	1:C:397:ASN:ND2	2.32	0.61
1:C:273:GLN:HE22	1:C:286:ALA:HA	1.72	0.55
1:C:234:ASN:OD1	1:C:1247:TRP:NE1	2.40	0.54
1:C:273:GLN:O	1:C:277:ASP:N	2.32	0.52
1:C:448:GLY:HA3	1:C:586:LEU:HD11	1.90	0.52
1:C:863:HIS:O	1:C:867:HIS:ND1	2.36	0.52
1:C:441:MET:HG2	1:C:442:PRO:HD3	1.90	0.52
1:C:781:ASN:N	1:C:1206:GLU:OE2	2.23	0.51
1:C:381:ILE:HG21	1:C:1242:THR:HG21	1.92	0.50
1:C:1139:THR:HG22	1:C:1305:MET:HE2	1.92	0.50
1:C:778:TRP:CD2	1:C:1210:GLY:HA3	2.49	0.47
1:C:1253:GLU:OE2	1:C:1293:ASN:ND2	2.47	0.47
1:C:586:LEU:HA	1:C:589:PRO:HD2	1.97	0.47
1:C:865:SER:O	1:C:869:MET:N	2.40	0.47
1:C:302:SER:O	1:C:376:SER:OG	2.25	0.47
1:C:431:PHE:HE1	1:C:600:THR:HG22	1.81	0.46
1:C:707:GLN:NE2	1:C:709:THR:OG1	2.49	0.45
1:C:311:LEU:HA	1:C:314:PHE:CD2	2.52	0.45
1:C:1522:LYS:O	1:C:1526:THR:N	2.40	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:263:TYR:HE1	1:C:394:LYS:HD3	1.81	0.45
1:C:893:TYR:HA	1:C:896:HIS:HD2	1.81	0.45
1:C:995:LYS:HA	1:C:1306:GLU:OE2	2.16	0.45
1:C:1145:ARG:NH2	2:D:2:LEU:O	2.50	0.44
1:C:1192:LEU:O	1:C:1196:THR:OG1	2.26	0.44
1:C:1161:THR:O	1:C:1161:THR:OG1	2.26	0.44
1:C:783:THR:O	1:C:787:ASN:N	2.41	0.44
1:C:371:THR:OG1	1:C:1254:TYR:OH	2.20	0.43
1:C:253:LEU:HA	1:C:254:PRO:HD3	1.88	0.43
1:C:1549:VAL:O	1:C:1561:ASP:N	2.46	0.42
1:C:396:TYR:HA	1:C:399:ILE:HD12	2.00	0.42
1:C:778:TRP:O	1:C:842:ARG:NH2	2.40	0.42
1:C:789:THR:O	1:C:792:SER:OG	2.32	0.42
1:C:1027:LEU:HA	1:C:1030:ILE:HG22	2.02	0.42
1:C:771:ALA:HB3	1:C:851:VAL:HA	2.02	0.42
1:C:374:GLN:NE2	1:C:1249:GLU:OE1	2.46	0.41
1:C:537:ALA:HA	1:C:540:THR:HG22	2.02	0.41
1:C:458:VAL:HG23	1:C:461:LEU:HD12	2.00	0.41
1:C:1169:LEU:HD23	1:C:1169:LEU:HA	1.92	0.41
1:C:324:VAL:HB	1:C:574:SER:HA	2.03	0.41
1:C:299:LEU:HD21	1:C:435:CYS:SG	2.61	0.41
1:C:1251:CYS:O	1:C:1255:ILE:HG12	2.21	0.41
1:C:685:PHE:HA	1:C:697:SER:HA	2.03	0.40
1:C:291:LEU:HD22	1:C:387:LEU:HD21	2.02	0.40
1:C:368:LEU:HD13	1:C:368:LEU:HA	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	1163/1582 (74%)	1083 (93%)	80 (7%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	17/390 (4%)	15 (88%)	2 (12%)	0	100	100
All	All	1180/1972 (60%)	1098 (93%)	82 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	508/1373 (37%)	508 (100%)	0	100	100
2	D	1/341 (0%)	1 (100%)	0	100	100
All	All	509/1714 (30%)	509 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	264	GLN
1	C	273	GLN
1	C	374	GLN
1	C	397	ASN
1	C	444	GLN
1	C	491	HIS
1	C	707	GLN
1	C	892	GLN
1	C	896	HIS
1	C	1225	GLN
1	C	1273	HIS
1	C	1316	HIS
1	C	1364	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ATP	C	1602	-	32,33,33	1.31	5 (15%)	48,52,52	1.71	12 (25%)
3	BJX	C	1601	-	35,35,35	1.38	3 (8%)	45,47,47	2.37	12 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	C	1602	-	-	6/22/38/38	0/3/3/3
3	BJX	C	1601	-	-	15/27/35/35	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1601	BJX	C16-N1	5.29	1.45	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1602	ATP	C5-C4	4.30	1.46	1.39
3	C	1601	BJX	C6-N	3.10	1.48	1.41
4	C	1602	ATP	C5-N7	-2.73	1.34	1.39
4	C	1602	ATP	C4-N9	-2.73	1.32	1.37
3	C	1601	BJX	O-C16	-2.36	1.18	1.23
4	C	1602	ATP	C5-C6	2.25	1.47	1.41
4	C	1602	ATP	C8-N7	2.18	1.35	1.31

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1601	BJX	C5-N1-C16	8.25	135.75	123.44
3	C	1601	BJX	C8-C5-N1	7.11	121.50	109.71
4	C	1602	ATP	C5-C4-N3	-4.76	120.16	126.72
3	C	1601	BJX	C7-C6-N	4.40	126.08	119.32
3	C	1601	BJX	C11-C7-C5	-4.14	110.76	120.46
4	C	1602	ATP	N3-C4-N9	3.77	133.58	127.17
3	C	1601	BJX	C7-C5-N1	-3.70	104.14	111.44
4	C	1602	ATP	N3-C2-N1	-3.49	123.30	128.58
4	C	1602	ATP	C2-N3-C4	3.38	120.09	111.83
3	C	1601	BJX	C10-C6-N	-3.02	117.69	122.42
3	C	1601	BJX	C10-C6-C7	-2.97	116.22	120.08
4	C	1602	ATP	C4-C5-N7	-2.91	107.25	110.58
3	C	1601	BJX	C2-C-N	-2.88	105.61	111.06
3	C	1601	BJX	C21-C23-C24	-2.71	118.84	123.73
4	C	1602	ATP	C4-N9-C8	2.69	108.56	105.74
3	C	1601	BJX	C11-C7-C6	2.60	121.29	118.36
3	C	1601	BJX	C-N-C1	-2.59	105.73	111.57
3	C	1601	BJX	C4-C3-C1	2.49	115.80	111.19
4	C	1602	ATP	C2'-C1'-N9	-2.43	107.25	113.30
4	C	1602	ATP	O3G-PG-O2G	2.40	116.80	107.80
4	C	1602	ATP	C3'-C2'-C1'	2.24	105.71	101.46
4	C	1602	ATP	C2-N1-C6	2.23	122.40	118.73
4	C	1602	ATP	C2'-C3'-C4'	2.13	106.72	102.61
4	C	1602	ATP	C6-C5-N7	2.13	136.19	132.09

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1601	BJX	C7-C5-C8-C9
3	C	1601	BJX	N1-C5-C7-C6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	1601	BJX	C8-C5-N1-C16
3	C	1601	BJX	C21-C23-C24-O2
4	C	1602	ATP	C5'-O5'-PA-O2A
4	C	1602	ATP	C5'-O5'-PA-O3A
4	C	1602	ATP	C4'-C5'-O5'-PA
3	C	1601	BJX	C21-C23-C24-O3
3	C	1601	BJX	N1-C5-C8-C9
3	C	1601	BJX	O-C16-C17-C18
4	C	1602	ATP	PB-O3A-PA-O1A
3	C	1601	BJX	C8-C5-C7-C11
3	C	1601	BJX	C8-C5-C7-C6
3	C	1601	BJX	N1-C5-C7-C11
3	C	1601	BJX	N1-C16-C17-C18
3	C	1601	BJX	C19-C21-O1-C25
3	C	1601	BJX	C22-C23-C24-O2
3	C	1601	BJX	C23-C21-O1-C25
3	C	1601	BJX	C22-C23-C24-O3
4	C	1602	ATP	PB-O3A-PA-O2A
4	C	1602	ATP	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20528. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 75



Y Index: 77




Z Index: 102


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0245. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 77 nm³; this corresponds to an approximate mass of 69 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.


8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-20528 and PDB model 6PZ9. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0245 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0245).

9.4 Atom inclusion [i](#)



At the recommended contour level, 89% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0245) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8120	0.4270
C	0.8150	0.4280
D	0.6390	0.4050

