



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:19 PM UTC

PDB ID : 3Q0Z / pdb_00003q0z
Title : Crystal structure of the hepatitis C virus NS5B RNA-dependent RNA polymerase complex with (2E)-3-(4-{{(1-{{(13-cyclohexyl-6-oxo-6,7-dihydro-5h-indolo[1,2-d][1,4]benzodiazepin-10-yl)carbonyl]amino}cyclopentyl)carbonyl]amino}phenyl)prop-2-enoic acid
Authors : Sheriff, S.
Deposited on : 2010-12-16
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ①](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	574	3% 87% 8% • •
1	B	574	4% 87% 8% 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	551	Total	C	N	O	S	0	0	0
			4274	2694	758	789	33			
1	B	545	Total	C	N	O	S	0	0	0
			4206	2648	744	782	32			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q9WMX2
B	0	MET	-	initiating methionine	UNP Q9WMX2

- Molecule 2 is (2E)-3-(4-((1-((13-cyclohexyl-6-oxo-6,7-dihydro-5H-indolo[1,2-d][1,4]benzodiazepin-10-yl)carbonyl]amino)cyclopentyl)carbonyl]amino)phenyl)prop-2-enoic acid (CCD ID: 23E) (formula: C₃₈H₃₈N₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			47	38	4	5		
2	B	1	Total	C	N	O	0	0
			47	38	4	5		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	254	Total 254	O 254	0	0
4	B	255	Total 255	O 255	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.30Å 91.40Å 231.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.08 – 2.29 23.08 – 2.29	Depositor EDS
% Data completeness (in resolution range)	97.9 (23.08-2.29) 97.8 (23.08-2.29)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.29Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.9.4, BUSTER 2.9.4	Depositor
R, R_{free}	0.200 , 0.224 0.198 , 0.224	Depositor DCC
R_{free} test set	919 reflections (1.46%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9133	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 23E, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/4369	1.24	13/5928 (0.2%)
1	B	0.86	0/4297	1.26	13/5834 (0.2%)
All	All	0.85	0/8666	1.25	26/11762 (0.2%)

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	550	TRP	N-CA-C	8.88	120.88	111.03
1	B	550	TRP	N-CA-C	8.27	120.21	111.03
1	A	318	ASP	CA-CB-CG	7.01	119.61	112.60
1	B	318	ASP	CA-CB-CG	6.65	119.25	112.60
1	A	40	THR	CB-CA-C	6.15	119.68	109.53
1	B	244	ASP	N-CA-C	-6.15	99.84	109.25
1	A	115	ALA	CA-C-N	5.92	128.57	120.46
1	A	115	ALA	C-N-CA	5.92	128.57	120.46
1	A	244	ASP	N-CA-C	-5.89	100.24	109.25
1	B	261	TYR	N-CA-C	5.79	117.27	111.07
1	A	550	TRP	CA-C-N	-5.72	115.31	123.20
1	A	550	TRP	C-N-CA	-5.72	115.31	123.20
1	B	115	ALA	CA-C-N	5.55	128.07	120.46
1	B	115	ALA	C-N-CA	5.55	128.07	120.46
1	A	375	ASP	CA-CB-CG	5.46	118.06	112.60
1	A	530	VAL	N-CA-C	5.37	116.96	109.55
1	A	11	ILE	N-CA-C	-5.35	99.78	107.37
1	B	550	TRP	CA-C-N	-5.34	115.83	123.20
1	B	550	TRP	C-N-CA	-5.34	115.83	123.20
1	B	1	SER	CA-C-N	5.22	128.28	120.87
1	B	1	SER	C-N-CA	5.22	128.28	120.87
1	A	55	ASP	CA-CB-CG	5.17	117.77	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	418	THR	CA-C-N	5.10	127.07	120.44
1	B	418	THR	C-N-CA	5.10	127.07	120.44
1	A	261	TYR	N-CA-C	5.06	116.79	111.28
1	B	175	LEU	N-CA-C	5.03	121.11	114.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4274	0	4253	21	0
1	B	4206	0	4179	20	0
2	A	47	0	37	2	0
2	B	47	0	37	1	0
3	A	25	0	0	0	0
3	B	25	0	0	0	0
4	A	254	0	0	1	0
4	B	255	0	0	1	0
All	All	9133	0	8506	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:THR:HG22	1:A:143:GLU:H	1.37	0.87
1:B:308:LEU:HB3	1:B:311:CYS:SG	2.31	0.71
1:A:405:VAL:HG11	1:A:446:GLN:HG2	1.77	0.66
1:B:321:VAL:CG2	1:B:365:SER:HB2	2.26	0.65
1:A:321:VAL:HG21	1:A:365:SER:CB	2.27	0.64
1:B:321:VAL:HG21	1:B:365:SER:CB	2.28	0.63
1:A:321:VAL:CG2	1:A:365:SER:HB2	2.29	0.62
1:B:321:VAL:CG2	1:B:365:SER:CB	2.78	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.82	0.61
1:A:321:VAL:CG2	1:A:365:SER:CB	2.79	0.60
2:A:601:23E:O6	2:A:601:23E:H7	2.01	0.60
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.86	0.58
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.43	0.57
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.85	0.57
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.89	0.54
1:A:40:THR:HG21	4:A:859:HOH:O	2.08	0.53
1:A:540:PRO:HB2	1:A:573:MET:HE2	1.91	0.52
1:A:197:PRO:O	1:A:201:VAL:HG23	2.09	0.51
1:B:197:PRO:O	1:B:201:VAL:HG23	2.11	0.51
1:A:40:THR:HG23	1:A:141:LYS:O	2.11	0.50
1:B:172:LYS:HE3	1:B:560:ILE:HD13	1.94	0.49
1:B:440:GLU:H	1:B:440:GLU:CD	2.21	0.48
1:A:440:GLU:H	1:A:440:GLU:CD	2.22	0.47
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.95	0.47
2:A:601:23E:O6	2:A:601:23E:C7	2.64	0.46
1:B:309:GLN:O	1:B:324:CYS:HB2	2.17	0.45
1:A:12:THR:HG21	1:A:270:LYS:HG3	1.99	0.44
1:B:236:GLU:OE1	1:B:280:ARG:NH2	2.39	0.43
1:A:321:VAL:CG2	1:A:365:SER:HB3	2.48	0.43
1:B:309:GLN:HB2	1:B:325:GLU:HB3	2.00	0.43
1:A:234:ARG:HG3	1:A:262:ILE:HD11	2.00	0.42
2:B:601:23E:H7	2:B:601:23E:O6	2.19	0.42
1:A:523:LYS:HE3	1:A:534:LEU:HD23	2.02	0.42
1:B:321:VAL:HG21	1:B:365:SER:HB2	1.95	0.41
1:B:466:LEU:HD22	1:B:551:PHE:HE2	1.86	0.41
1:B:321:VAL:CG2	1:B:365:SER:HB3	2.51	0.41
1:B:405:VAL:HG13	1:B:445:CYS:HA	2.02	0.41
1:A:83:LEU:HB2	1:A:173:MET:HA	2.03	0.41
1:A:572:PHE:CE2	1:A:573:MET:HE3	2.55	0.41
1:B:335:SER:HB3	4:B:773:HOH:O	2.20	0.41
1:B:445:CYS:SG	1:B:454:ILE:HD12	2.60	0.41
1:B:448:TYR:CE2	1:B:551:PHE:HD1	2.39	0.41
1:A:187:MET:HE2	1:A:293:LEU:HD23	2.03	0.40
1:B:12:THR:HG21	1:B:270:LYS:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	547/574 (95%)	539 (98%)	8 (2%)	0	100	100
1	B	541/574 (94%)	528 (98%)	13 (2%)	0	100	100
All	All	1088/1148 (95%)	1067 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	458/486 (94%)	443 (97%)	15 (3%)	33	50
1	B	451/486 (93%)	439 (97%)	12 (3%)	39	58
All	All	909/972 (94%)	882 (97%)	27 (3%)	36	53

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	THR
1	A	141	LYS
1	A	159	LEU
1	A	236	GLU
1	A	262	ILE
1	A	308	LEU
1	A	329	THR
1	A	336	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	ASP
1	A	372	VAL
1	A	380	ARG
1	A	476	SER
1	A	498	ARG
1	A	531	ARG
1	A	536	LEU
1	B	43	ARG
1	B	98	ARG
1	B	141	LYS
1	B	151	LYS
1	B	159	LEU
1	B	307	LYS
1	B	308	LEU
1	B	329	THR
1	B	352	ASP
1	B	379	LYS
1	B	399	THR
1	B	531	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	272	GLN
1	B	309	GLN
1	B	461	GLN
1	B	514	GLN
1	B	562	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SO4	A	655	-	4,4,4	0.20	0	6,6,6	0.07	0
3	SO4	B	651	-	4,4,4	0.50	0	6,6,6	0.22	0
3	SO4	A	651	-	4,4,4	0.57	0	6,6,6	0.10	0
3	SO4	A	654	-	4,4,4	0.19	0	6,6,6	0.09	0
3	SO4	A	652	-	4,4,4	0.42	0	6,6,6	0.19	0
2	23E	A	601	-	52,53,53	1.62	6 (11%)	68,77,77	1.26	7 (10%)
3	SO4	B	652	-	4,4,4	0.29	0	6,6,6	0.16	0
2	23E	B	601	-	52,53,53	1.64	6 (11%)	68,77,77	1.23	8 (11%)
3	SO4	B	657	-	4,4,4	0.52	0	6,6,6	0.25	0
3	SO4	B	653	-	4,4,4	0.17	0	6,6,6	0.13	0
3	SO4	B	656	-	4,4,4	0.42	0	6,6,6	0.39	0
3	SO4	A	653	-	4,4,4	0.45	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	23E	B	601	-	-	4/28/57/57	0/6/7/7
2	23E	A	601	-	-	1/28/57/57	0/6/7/7

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	23E	O15-C14	5.14	1.35	1.23
2	B	601	23E	O15-C14	4.96	1.35	1.23
2	B	601	23E	C32-C30	4.82	1.51	1.47
2	A	601	23E	C30-N26	-4.37	1.32	1.39
2	A	601	23E	C28-C29	4.19	1.51	1.44
2	B	601	23E	C30-N26	-4.19	1.32	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	23E	C32-C30	3.92	1.50	1.47
2	B	601	23E	C28-C29	3.90	1.50	1.44
2	B	601	23E	O20-C14	-2.71	1.23	1.30
2	A	601	23E	C27-N26	-2.66	1.34	1.39
2	A	601	23E	O20-C14	-2.60	1.23	1.30
2	B	601	23E	C27-N26	-2.30	1.35	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	23E	C27-N26-C30	4.00	111.57	108.37
2	B	601	23E	C27-N26-C30	3.59	111.24	108.37
2	B	601	23E	C27-C28-C29	-3.15	104.37	107.02
2	A	601	23E	C27-C28-C29	-2.86	104.61	107.02
2	B	601	23E	C24-C28-C27	2.81	122.48	119.24
2	B	601	23E	C17-C2-C16	-2.78	99.89	103.39
2	B	601	23E	C23-C24-C28	-2.57	116.62	120.86
2	A	601	23E	C9-C12-C13	-2.49	121.47	126.92
2	A	601	23E	C28-C29-C30	-2.42	104.94	107.28
2	A	601	23E	C16-C2-C3	2.34	116.22	110.75
2	A	601	23E	C17-C2-C16	-2.29	100.51	103.39
2	B	601	23E	C16-C2-C3	2.27	116.06	110.75
2	B	601	23E	O38-C21-C22	-2.17	116.60	120.90
2	A	601	23E	C5-N4-C3	-2.12	123.09	126.86
2	B	601	23E	O47-C46-C44	-2.04	117.68	121.40

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	23E	C12-C13-C14-O20
2	B	601	23E	C12-C13-C14-O15
2	B	601	23E	C13-C12-C9-C8
2	B	601	23E	C13-C12-C9-C10
2	A	601	23E	C13-C12-C9-C8

There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	23E	2	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	23E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	551/574 (95%)	0.08	19 (3%)	48 50	23, 35, 66, 121	0
1	B	545/574 (94%)	0.19	24 (4%)	39 41	23, 36, 68, 130	0
All	All	1096/1148 (95%)	0.13	43 (3%)	43 45	23, 36, 67, 130	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	567	ALA	7.2
1	B	149	PRO	5.2
1	A	149	PRO	4.6
1	B	151	LYS	4.2
1	B	566	ARG	4.1
1	A	152	GLY	4.0
1	B	37	VAL	3.9
1	A	532	THR	3.9
1	B	405	VAL	3.6
1	A	568	ARG	3.5
1	A	573	MET	3.5
1	A	147	VAL	3.5
1	A	153	GLY	3.4
1	A	151	LYS	3.3
1	A	148	GLN	3.3
1	B	542	ALA	3.3
1	A	14	CYS	3.3
1	B	147	VAL	3.2
1	A	150	GLU	3.1
1	B	402	HIS	3.1
1	B	564	LEU	3.1
1	B	541	ALA	3.0
1	B	14	CYS	2.9
1	B	148	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	569	PRO	2.8
1	A	37	VAL	2.7
1	A	531	ARG	2.6
1	B	152	GLY	2.5
1	A	533	LYS	2.5
1	A	329	THR	2.4
1	B	404	PRO	2.4
1	B	403	THR	2.4
1	B	150	GLU	2.4
1	A	327	ALA	2.3
1	B	544	GLN	2.3
1	B	153	GLY	2.3
1	B	399	THR	2.1
1	B	531	ARG	2.1
1	A	40	THR	2.0
1	A	154	ARG	2.0
1	B	550	TRP	2.0
1	B	400	ALA	2.0
1	B	488	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	657	5/5	0.72	0.15	94,98,99,100	0
3	SO4	B	651	5/5	0.76	0.12	77,81,82,83	0
3	SO4	B	652	5/5	0.79	0.28	80,85,86,86	0
3	SO4	B	656	5/5	0.83	0.16	75,79,81,81	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	655	5/5	0.83	0.17	111,115,116,117	0
3	SO4	B	653	5/5	0.84	0.12	87,91,92,93	0
3	SO4	A	652	5/5	0.87	0.17	70,75,75,75	0
3	SO4	A	654	5/5	0.91	0.12	70,74,75,76	0
3	SO4	A	651	5/5	0.93	0.07	58,62,63,64	0
2	23E	A	601	47/47	0.93	0.08	25,30,60,71	0
2	23E	B	601	47/47	0.93	0.08	29,34,45,55	0
3	SO4	A	653	5/5	0.96	0.07	47,52,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 23E A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 23E B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.