



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2026 – 01:12 PM EDT

PDB ID : 6Q0C / pdb_00006q0c
Title : MutY adenine glycosylase bound to DNA containing a transition state analog (1N) paired with undamaged dG
Authors : O'Shea Murray, V.L.; Russelburg, L.P.; Horvath, M.P.; David, S.S.
Deposited on : 2019-08-01
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	23% 88% 6% • 5%
2	B	11	100%
3	C	11	82% 9% 9%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6137 atoms, of which 2885 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called A/G-specific adenine glycosylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	349	Total	C	H	N	O	S	0	4	0
			5360	1764	2630	463	492	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	ASP	ASN	variant	UNP P83847

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*AP*GP*AP*CP*GP*TP*GP*GP*AP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	11	Total	C	H	N	O	P	0	1	0
			381	118	136	53	64	10			

- Molecule 3 is a DNA chain called DNA (5'-D(P*GP*TP*CP*CP*AP*(NR1)P*GP*TP*CP*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	10	Total	C	H	N	O	P	0	0	0
			306	92	113	31	60	10			

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	56	Total	O	0	0
			56	56		
7	B	7	Total	O	0	1
			8	8		
7	C	7	Total	O	0	0
			7	7		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: A/G-specific adenine glycosylase



- Molecule 2: DNA (5'-D(*AP*AP*GP*AP*CP*GP*TP*GP*GP*AP*C)-3')



There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(P*GP*TP*CP*CP*AP*(NR1)P*GP*TP*CP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	37.86Å 86.14Å 141.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.30 – 2.00 41.30 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.3 (41.30-2.00) 96.4 (41.30-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.15rc2_3428	Depositor
R, R_{free}	0.247 , 0.271 0.250 , 0.274	Depositor DCC
R_{free} test set	1498 reflections (2.90%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6137	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, NRI, EDO, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/2812	0.37	0/3835
2	B	0.22	0/277	0.52	0/427
3	C	0.19	0/201	0.49	0/305
All	All	0.15	0/3290	0.39	0/4567

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2730	2630	2631	13	1
2	B	245	136	136	0	0
3	C	193	113	112	0	1
4	A	8	0	0	0	0
5	A	1	0	0	0	0
6	A	4	6	6	0	0
7	A	56	0	0	0	0
7	B	8	0	0	0	0
7	C	7	0	0	0	0
All	All	3252	2885	2885	13	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:LEU:HD13	1:A:294:VAL:HG11	1.46	0.97
1:A:283:GLU:HG2	1:A:296:LEU:HD12	1.59	0.84
1:A:68:LEU:HG	1:A:99:VAL:HG13	1.78	0.65
1:A:292:LEU:CD2	1:A:321:LEU:HD11	2.28	0.63
1:A:292:LEU:HD22	1:A:321:LEU:HD11	1.82	0.61
1:A:273:THR:O	1:A:273:THR:HG22	2.06	0.56
1:A:271[B]:CYS:SG	1:A:282:LEU:HD23	2.54	0.48
1:A:81:LYS:HA	1:A:343:ALA:HB2	1.97	0.47
1:A:174:VAL:O	1:A:178:MET:N	2.49	0.46
1:A:254:ARG:NH2	1:A:270:SER:O	2.49	0.46
1:A:68:LEU:CG	1:A:99:VAL:HG13	2.45	0.46
1:A:72:ALA:HB2	1:A:99:VAL:HG12	2.01	0.41
1:A:234:VAL:HG13	1:A:310:LEU:HA	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59[B]:GLU:OE2	3:C:22:DT:O3'[1_455]	1.96	0.24

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/366 (95%)	341 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	276/313 (88%)	268 (97%)	8 (3%)	37 40

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	ASP
1	A	234	VAL
1	A	243	VAL
1	A	284	GLN
1	A	292	LEU
1	A	294	VAL
1	A	313	GLN
1	A	322	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NRI	C	18	3	7,11,12	0.56	0	3,14,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NRI	C	18	3	-	0/3/15/16	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	A	401	1	0,12,12	-	-	-		
6	EDO	A	403	-	3,3,3	0.41	0	2,2,2	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	401	1	-	-	0/6/5/5
6	EDO	A	403	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/366 (95%)	1.37	84 (24%) 2 2	23, 62, 98, 109	4 (1%)
2	B	11/11 (100%)	0.44	0 100 100	26, 47, 76, 77	1 (9%)
3	C	9/11 (81%)	0.26	0 100 100	33, 44, 70, 79	0
All	All	369/388 (95%)	1.31	84 (22%) 2 2	23, 62, 98, 109	5 (1%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	ALA	5.7
1	A	355	TYR	5.4
1	A	296	LEU	4.7
1	A	287	GLY	4.5
1	A	325	GLY	4.5
1	A	279	LYS	4.3
1	A	273	THR	4.3
1	A	213	TYR	4.2
1	A	282	LEU	4.2
1	A	217	PHE	3.9
1	A	241	VAL	3.8
1	A	242	ALA	3.8
1	A	240	ALA	3.8
1	A	278	GLY	3.8
1	A	252	LEU	3.6
1	A	274	ASP	3.6
1	A	218	ALA	3.4
1	A	358	TRP	3.4
1	A	105	ARG	3.3
1	A	294	VAL	3.3
1	A	326	PRO	3.3
1	A	340	LYS	3.2
1	A	106	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	324	GLY	3.1
1	A	243	VAL	3.0
1	A	233	ALA	3.0
1	A	317	PHE	2.9
1	A	316	VAL	2.9
1	A	221	VAL	2.9
1	A	301	VAL	2.9
1	A	319	GLY	2.9
1	A	321	LEU	2.8
1	A	315	THR	2.8
1	A	266	TRP	2.8
1	A	156	LEU	2.8
1	A	235	LYS	2.8
1	A	239	LEU	2.7
1	A	331	TYR	2.7
1	A	323	HIS	2.7
1	A	108	GLY	2.7
1	A	251	VAL	2.7
1	A	322	VAL	2.7
1	A	320	ARG	2.6
1	A	258	SER	2.6
1	A	15	ARG	2.6
1	A	311	VAL	2.6
1	A	237	VAL	2.6
1	A	244	LEU	2.6
1	A	227	VAL	2.5
1	A	214	CYS	2.5
1	A	74	ALA	2.5
1	A	216	ALA	2.5
1	A	238	PRO	2.5
1	A	314	LEU	2.5
1	A	138	VAL	2.4
1	A	272	GLU	2.4
1	A	222	ALA	2.4
1	A	99	VAL	2.4
1	A	248	GLU	2.4
1	A	180	TYR	2.3
1	A	357	GLU	2.3
1	A	225	LEU	2.3
1	A	174	VAL	2.3
1	A	286	VAL	2.3
1	A	293	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	333	LEU	2.3
1	A	157	VAL	2.3
1	A	209	PRO	2.3
1	A	10	ALA	2.3
1	A	12	GLU	2.2
1	A	176	GLU	2.2
1	A	206	LEU	2.2
1	A	68	LEU	2.2
1	A	277	ASP	2.2
1	A	297	THR	2.2
1	A	179	ALA	2.1
1	A	269	PRO	2.1
1	A	352	TRP	2.1
1	A	268	PHE	2.1
1	A	292	LEU	2.1
1	A	139	PRO	2.1
1	A	302	SER	2.1
1	A	77[A]	ASP	2.1
1	A	318	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NRI	C	18	11/12	0.98	0.05	21,28,32,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	A	403	4/4	0.89	0.16	34,42,55,66	0
5	CA	A	402	1/1	0.96	0.07	47,47,47,47	0
4	SF4	A	401	8/8	0.97	0.06	40,45,49,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.