



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:00 AM UTC

PDB ID : 2Q1L / pdb_00002q1l
Title : Design and Synthesis of Pyrrole-based, Hepatoselective HMG-CoA Reductase Inhibitors
Authors : Pavlovsky, A.; Pfefferkorn, J.A.; Harris, M.S.; Finzel, B.C.
Deposited on : 2007-05-24
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	10% 81% 7% 11%
1	B	441	15% 84% 7% 9%
1	C	441	11% 86% 7% 6%
1	D	441	7% 84% 5% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxy-3-methylglutaryl-coenzyme A reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			2903	1806	511	557	29			
1	B	401	Total	C	N	O	S	0	0	0
			2979	1859	524	567	29			
1	C	416	Total	C	N	O	S	0	0	0
			3087	1923	540	594	30			
1	D	394	Total	C	N	O	S	0	0	0
			2920	1818	514	559	29			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	435	HIS	-	expression tag	UNP P04035
A	436	HIS	-	expression tag	UNP P04035
A	437	HIS	-	expression tag	UNP P04035
A	438	HIS	-	expression tag	UNP P04035
A	439	HIS	-	expression tag	UNP P04035
A	440	HIS	-	expression tag	UNP P04035
A	485	ILE	MET	engineered mutation	UNP P04035
B	435	HIS	-	expression tag	UNP P04035
B	436	HIS	-	expression tag	UNP P04035
B	437	HIS	-	expression tag	UNP P04035
B	438	HIS	-	expression tag	UNP P04035
B	439	HIS	-	expression tag	UNP P04035
B	440	HIS	-	expression tag	UNP P04035
B	485	ILE	MET	engineered mutation	UNP P04035
C	435	HIS	-	expression tag	UNP P04035
C	436	HIS	-	expression tag	UNP P04035
C	437	HIS	-	expression tag	UNP P04035
C	438	HIS	-	expression tag	UNP P04035
C	439	HIS	-	expression tag	UNP P04035
C	440	HIS	-	expression tag	UNP P04035
C	485	ILE	MET	engineered mutation	UNP P04035

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	435	HIS	-	expression tag	UNP P04035
D	436	HIS	-	expression tag	UNP P04035
D	437	HIS	-	expression tag	UNP P04035
D	438	HIS	-	expression tag	UNP P04035
D	439	HIS	-	expression tag	UNP P04035
D	440	HIS	-	expression tag	UNP P04035
D	485	ILE	MET	engineered mutation	UNP P04035

- Molecule 2 is (3R,5R)-7-[5-(ANILINOCARBONYL)-3,4-BIS(4-FLUOROPHENYL)-1-ISOPROPYL-1H-PYRROL-2-YL]-3,5-DIHYDROXYHEPTANOIC ACID (CCD ID: 882) (formula: C₃₃H₃₄F₂N₂O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			42	33	2	2	5		
2	A	1	Total	C	F	N	O	0	0
			42	33	2	2	5		
2	C	1	Total	C	F	N	O	0	0
			42	33	2	2	5		
2	D	1	Total	C	F	N	O	0	0
			42	33	2	2	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	132	Total	O	0	0
			132	132		





Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	112	Total 112	O 112	0	0
3	C	110	Total 110	O 110	0	0
3	D	131	Total 131	O 131	0	0



● Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.73Å 176.96Å 76.73Å 90.00° 118.83° 90.00°	Depositor
Resolution (Å)	29.49 – 2.05 29.49 – 2.05	Depositor EDS
% Data completeness (in resolution range)	79.7 (29.49-2.05) 65.7 (29.49-2.05)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.04Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.223 , 0.263 0.238 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.294	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.026 for -h-l,k,h 0.026 for l,k,-h-l 0.040 for h,-k,-h-l 0.045 for -h-l,-k,l 0.038 for l,-k,h	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	12542	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
882

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2943	0.80	1/3977 (0.0%)
1	B	0.51	1/3022 (0.0%)	0.79	0/4084
1	C	0.52	0/3130	0.79	0/4230
1	D	0.52	0/2960	0.80	0/3999
All	All	0.52	1/12055 (0.0%)	0.80	1/16290 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	555	MET	SD-CE	-5.47	1.65	1.79

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	802	ILE	N-CA-C	5.59	115.69	107.75

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2903	0	2933	14	0
1	B	2979	0	3020	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3087	0	3121	21	0
1	D	2920	0	2957	9	0
2	A	84	0	66	1	0
2	C	42	0	33	1	0
2	D	42	0	33	1	0
3	A	132	0	0	0	0
3	B	112	0	0	0	0
3	C	110	0	0	0	0
3	D	131	0	0	0	0
All	All	12542	0	12163	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:655:MET:SD	1:A:657:MET:HG2	2.27	0.74
1:A:817:CYS:HA	1:A:820:MET:HE3	1.68	0.74
1:A:771:ASN:OD1	1:A:775:SER:OG	2.10	0.69
1:C:681:LEU:HD22	1:D:731:VAL:HG22	1.73	0.69
1:C:655:MET:SD	1:C:657:MET:HG2	2.35	0.67
1:C:445:ASN:HD22	1:C:469:GLN:HE21	1.44	0.65
1:C:523:MET:HE1	1:C:530:VAL:CG2	2.29	0.63
1:C:771:ASN:OD1	1:C:775:SER:OG	2.17	0.62
1:C:456:GLU:C	1:C:458:GLY:N	2.60	0.59
1:B:655:MET:SD	1:B:657:MET:HG2	2.43	0.57
1:C:655:MET:HE1	1:C:657:MET:HG3	1.90	0.54
1:D:629:ALA:HB2	1:D:659:MET:HE3	1.88	0.54
1:C:523:MET:HE1	1:C:530:VAL:HG23	1.89	0.53
1:C:774:SER:HA	1:C:799:SER:O	2.08	0.53
1:D:781:MET:HE2	1:D:793:ILE:HD12	1.92	0.51
1:C:487:THR:HG23	1:C:490:ARG:HB2	1.92	0.51
1:B:555:MET:CE	1:B:563:VAL:HA	2.41	0.51
1:A:590:ARG:NH2	1:A:657:MET:HE3	2.27	0.49
1:B:655:MET:HE1	1:B:657:MET:HG3	1.95	0.49
1:D:606:LYS:NZ	1:D:610:GLU:OE2	2.43	0.49
2:C:876:882:O1	2:C:876:882:H133	2.12	0.49
1:C:731:VAL:HG12	1:C:854:MET:HE1	1.95	0.48
1:B:606:LYS:NZ	1:B:610:GLU:OE2	2.47	0.47
1:A:796:THR:HG21	1:D:638:ILE:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:533:TYR:CD1	1:B:535:PRO:HB3	2.49	0.47
1:B:700:GLU:OE2	1:C:700:GLU:OE2	2.33	0.46
1:A:596:LEU:HB3	1:A:601:ASP:HB2	1.98	0.46
1:B:592:PRO:HD2	1:B:645:ILE:O	2.15	0.46
1:B:555:MET:HE1	1:B:563:VAL:HA	1.97	0.46
1:C:470:LEU:O	1:C:473:ALA:O	2.34	0.45
1:A:541:ALA:HA	1:A:567:ASN:HD22	1.82	0.45
1:A:542:GLY:H	1:A:567:ASN:ND2	2.13	0.45
1:C:719:GLU:O	1:C:722:LYS:NZ	2.48	0.45
1:C:651:SER:HA	1:C:659:MET:HE3	1.98	0.44
1:C:820:MET:HE3	1:D:499:LEU:HD21	1.98	0.44
1:B:774:SER:HA	1:B:799:SER:O	2.18	0.43
1:B:702:ARG:O	1:B:799:SER:HA	2.17	0.43
1:D:588:MET:HB3	1:D:660:ILE:HD13	2.00	0.43
1:B:817:CYS:HA	1:B:820:MET:HE3	2.01	0.43
1:C:754:ALA:HB1	1:C:771:ASN:HD21	1.84	0.42
1:A:590:ARG:NH2	1:A:657:MET:CE	2.82	0.42
1:A:808:GLY:O	1:A:814:GLN:HG3	2.18	0.42
1:C:531:ILE:HD13	1:D:540:VAL:CG2	2.50	0.42
2:D:876:882:H133	2:D:876:882:O1	2.20	0.41
1:A:637:SER:HB2	1:A:687:TYR:OH	2.20	0.41
1:C:560:GLY:O	1:C:561:CYS:HB2	2.21	0.41
1:D:702:ARG:O	1:D:799:SER:HA	2.20	0.41
1:A:702:ARG:O	1:A:799:SER:HA	2.21	0.41
1:C:593:VAL:HG13	1:C:681:LEU:HB3	2.03	0.41
1:A:655:MET:HE1	1:A:657:MET:HG3	2.03	0.41
2:A:877:882:H133	2:A:877:882:O1	2.22	0.40
1:B:838:LEU:O	1:B:842:VAL:HG23	2.22	0.40
1:C:756:ILE:HD12	1:C:756:ILE:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/441 (88%)	374 (96%)	13 (3%)	1 (0%)	36	30
1	B	397/441 (90%)	381 (96%)	15 (4%)	1 (0%)	36	30
1	C	412/441 (93%)	391 (95%)	20 (5%)	1 (0%)	43	37
1	D	390/441 (88%)	375 (96%)	15 (4%)	0	100	100
All	All	1587/1764 (90%)	1521 (96%)	63 (4%)	3 (0%)	43	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	445	ASN
1	B	525	ALA
1	A	860	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/355 (87%)	297 (96%)	13 (4%)	26	20
1	B	318/355 (90%)	305 (96%)	13 (4%)	27	21
1	C	330/355 (93%)	317 (96%)	13 (4%)	28	23
1	D	312/355 (88%)	301 (96%)	11 (4%)	32	27
All	All	1270/1420 (89%)	1220 (96%)	50 (4%)	28	23

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	463	SER
1	A	485	ILE
1	A	486	GLU
1	A	498	LEU
1	A	512	LEU
1	A	540	VAL
1	A	630	ARG
1	A	634	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	704	LYS
1	A	788	ASN
1	A	802	ILE
1	A	814	GLN
1	A	851	LEU
1	B	470	LEU
1	B	482	GLU
1	B	505	GLU
1	B	523	MET
1	B	540	VAL
1	B	613	GLU
1	B	634	LEU
1	B	662	LYS
1	B	669	SER
1	B	746	ILE
1	B	756	ILE
1	B	771	ASN
1	B	828	LYS
1	C	445	ASN
1	C	452	LEU
1	C	460	LYS
1	C	470	LEU
1	C	476	ILE
1	C	487	THR
1	C	505	GLU
1	C	523	MET
1	C	657	MET
1	C	660	ILE
1	C	662	LYS
1	C	681	LEU
1	C	725	THR
1	D	470	LEU
1	D	501	LYS
1	D	613	GLU
1	D	657	MET
1	D	659	MET
1	D	660	ILE
1	D	688	CYS
1	D	788	ASN
1	D	791	LEU
1	D	802	ILE
1	D	851	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	518	ASN
1	A	567	ASN
1	A	635	HIS
1	A	732	ASN
1	A	736	ASN
1	A	788	ASN
1	A	861	HIS
1	B	472	ASN
1	B	510	GLN
1	B	529	ASN
1	B	632	GLN
1	C	469	GLN
1	C	472	ASN
1	C	488	HIS
1	C	497	GLN
1	C	679	GLN
1	C	819	GLN
1	D	472	ASN
1	D	488	HIS
1	D	518	ASN
1	D	632	GLN
1	D	770	GLN
1	D	788	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	882	A	877	-	45,45,45	0.85	2 (4%)	53,63,63	1.63	8 (15%)
2	882	C	876	-	45,45,45	0.82	2 (4%)	53,63,63	1.73	8 (15%)
2	882	A	876	-	45,45,45	0.82	2 (4%)	53,63,63	1.84	9 (16%)
2	882	D	876	-	45,45,45	0.88	2 (4%)	53,63,63	1.85	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	882	A	877	-	-	5/33/33/33	0/4/4/4
2	882	C	876	-	-	5/33/33/33	0/4/4/4
2	882	A	876	-	-	6/33/33/33	0/4/4/4
2	882	D	876	-	-	3/33/33/33	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	876	882	C1-C5	-2.33	1.36	1.46
2	D	876	882	C1-C5	-2.26	1.36	1.46
2	A	877	882	C1-C5	-2.19	1.36	1.46
2	C	876	882	C1-C5	-2.14	1.36	1.46
2	D	876	882	C2-N1	-2.08	1.36	1.41
2	A	876	882	C2-N1	-2.08	1.36	1.41
2	A	877	882	C2-N1	-2.03	1.36	1.41
2	C	876	882	C2-N1	-2.01	1.36	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	876	882	C3-C2-N1	6.87	129.91	122.33
2	D	876	882	C3-C2-N1	6.60	129.62	122.33
2	A	877	882	C3-C2-N1	6.10	129.06	122.33
2	C	876	882	C3-C2-N1	5.90	128.84	122.33
2	D	876	882	C2-C3-N2	5.05	122.14	115.32
2	A	876	882	C2-C3-N2	4.92	121.96	115.32
2	C	876	882	C2-C3-N2	4.61	121.54	115.32
2	A	877	882	C2-C3-N2	4.23	121.03	115.32
2	D	876	882	C21-C27-C5	-4.17	114.23	120.79
2	A	876	882	O1-C3-C2	-3.77	114.62	121.96
2	C	876	882	C32-N2-C3	-3.74	120.81	127.45
2	D	876	882	O1-C3-C2	-3.53	115.09	121.96
2	C	876	882	O1-C3-C2	-3.31	115.52	121.96
2	C	876	882	C21-C27-C5	-3.26	115.66	120.79
2	A	877	882	C21-C27-C5	-3.25	115.68	120.79
2	A	876	882	C32-N2-C3	-3.24	121.70	127.45
2	D	876	882	C18-C27-C5	3.17	125.78	120.79
2	A	876	882	C21-C27-C5	-3.15	115.83	120.79
2	A	877	882	O1-C3-C2	-3.03	116.05	121.96
2	D	876	882	C9-C10-C11	-2.87	109.74	114.28
2	A	876	882	C7-C8-C9	-2.83	108.23	113.58
2	D	876	882	C7-C8-C9	-2.59	108.69	113.58
2	A	877	882	C32-N2-C3	-2.59	122.86	127.45
2	A	876	882	C14-C6-N1	-2.46	108.25	111.69
2	D	876	882	C32-N2-C3	-2.45	123.11	127.45
2	A	876	882	C9-C10-C11	-2.40	110.48	114.28
2	C	876	882	C18-C27-C5	2.38	124.53	120.79
2	C	876	882	C7-C8-C9	-2.37	109.10	113.58
2	A	877	882	C19-C28-C22	-2.34	119.73	122.80
2	C	876	882	C19-C28-C22	-2.17	119.95	122.80
2	A	877	882	C25-C22-C28	2.14	120.58	118.38
2	A	876	882	C18-C15-C30	2.08	120.51	118.38
2	A	877	882	C18-C27-C5	2.06	124.02	120.79

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	877	882	C7-C8-C9-O4
2	C	876	882	C7-C8-C9-O4
2	A	877	882	C7-C8-C9-C10
2	C	876	882	C7-C8-C9-C10
2	D	876	882	C7-C8-C9-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	876	882	C7-C8-C9-O4
2	A	876	882	N1-C2-C3-O1
2	A	876	882	N1-C2-C3-N2
2	A	877	882	N1-C2-C3-N2
2	C	876	882	N1-C2-C3-N2
2	D	876	882	N1-C2-C3-N2
2	A	876	882	C7-C8-C9-C10
2	D	876	882	C7-C8-C9-C10
2	C	876	882	C26-C32-N2-C3
2	C	876	882	C17-C32-N2-C3
2	A	876	882	C1-C2-C3-N2
2	A	877	882	C14-C6-N1-C4
2	A	877	882	C26-C32-N2-C3
2	A	876	882	C4-C7-C8-C9


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	877	882	1	0
2	C	876	882	1	0
2	D	876	882	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 882 A 877



Ligand 882 C 876





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/441 (88%)	0.85	46 (11%) 9 8	22, 32, 55, 70	0
1	B	401/441 (90%)	1.09	66 (16%) 4 4	22, 37, 71, 89	0
1	C	416/441 (94%)	0.73	47 (11%) 10 9	19, 31, 60, 85	0
1	D	394/441 (89%)	0.71	33 (8%) 17 17	19, 31, 53, 90	0
All	All	1603/1764 (90%)	0.85	192 (11%) 9 8	19, 33, 58, 90	0

All (192) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	473	ALA	7.9
1	A	859	ALA	7.1
1	B	481	LEU	6.2
1	A	860	GLY	6.2
1	C	458	GLY	6.0
1	B	478	ALA	5.6
1	B	861	HIS	5.6
1	B	485	ILE	5.6
1	B	468	ILE	5.4
1	B	477	PRO	5.4
1	B	476	ILE	5.3
1	D	859	ALA	5.1
1	A	470	LEU	5.0
1	B	479	TYR	5.0
1	B	484	LEU	4.8
1	D	461	PHE	4.6
1	D	462	LEU	4.6
1	B	524	GLY	4.6
1	D	564	ALA	4.5
1	A	525	ALA	4.4
1	A	461	PHE	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	470	LEU	4.3
1	D	679	GLN	4.2
1	A	473	ALA	4.1
1	C	444	PRO	4.1
1	D	849	GLY	4.1
1	C	860	GLY	4.0
1	B	462	LEU	4.0
1	B	461	PHE	3.9
1	A	861	HIS	3.8
1	C	451	ILE	3.8
1	A	463	SER	3.7
1	C	455	ALA	3.7
1	A	486	GLU	3.7
1	B	483	THR	3.7
1	D	471	VAL	3.6
1	C	829	ASP	3.6
1	C	481	LEU	3.6
1	B	548	GLU	3.6
1	C	473	ALA	3.6
1	A	462	LEU	3.5
1	B	663	GLY	3.5
1	C	449	LEU	3.5
1	B	522	VAL	3.5
1	D	475	HIS	3.4
1	D	504	SER	3.4
1	C	859	ALA	3.4
1	A	469	GLN	3.3
1	A	465	ALA	3.3
1	C	598	ARG	3.3
1	C	456	GLU	3.2
1	A	468	ILE	3.2
1	A	504	SER	3.2
1	D	505	GLU	3.2
1	C	679	GLN	3.2
1	D	468	ILE	3.1
1	B	787	THR	3.1
1	B	459	ALA	3.1
1	D	473	ALA	3.1
1	C	478	ALA	3.0
1	A	485	ILE	3.0
1	B	786	PRO	3.0
1	B	517	TYR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	465	ALA	2.9
1	A	830	ASN	2.9
1	C	445	ASN	2.9
1	C	485	ILE	2.9
1	C	450	GLN	2.9
1	B	470	LEU	2.9
1	B	591	GLY	2.9
1	D	484	LEU	2.9
1	B	661	SER	2.9
1	B	466	GLU	2.8
1	B	488	HIS	2.8
1	B	491	GLY	2.8
1	D	486	GLU	2.8
1	A	487	THR	2.8
1	C	487	THR	2.8
1	B	514	TYR	2.8
1	B	525	ALA	2.8
1	C	677	GLU	2.8
1	B	471	VAL	2.8
1	B	860	GLY	2.7
1	D	628	PHE	2.7
1	B	474	LYS	2.7
1	B	831	PRO	2.7
1	B	512	LEU	2.7
1	C	483	THR	2.7
1	D	472	ASN	2.7
1	B	506	PRO	2.7
1	C	475	HIS	2.7
1	A	717	VAL	2.7
1	B	829	ASP	2.7
1	D	467	ILE	2.7
1	B	577	GLY	2.7
1	D	510	GLN	2.6
1	C	669	SER	2.6
1	A	727	ALA	2.6
1	C	657	MET	2.6
1	C	520	SER	2.6
1	D	577	GLY	2.6
1	B	480	LYS	2.6
1	B	625	THR	2.6
1	A	832	GLY	2.5
1	A	501	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	826	ALA	2.5
1	C	461	PHE	2.5
1	B	467	ILE	2.5
1	B	574	GLY	2.5
1	A	516	ASP	2.5
1	B	503	LEU	2.5
1	B	628	PHE	2.5
1	A	524	GLY	2.5
1	C	660	ILE	2.5
1	B	472	ASN	2.5
1	C	479	TYR	2.4
1	D	858	ALA	2.4
1	B	518	ASN	2.4
1	C	448	CYS	2.4
1	A	714	ALA	2.4
1	D	856	ALA	2.4
1	C	628	PHE	2.4
1	C	617	VAL	2.4
1	B	519	TYR	2.4
1	C	587	GLY	2.4
1	A	720	VAL	2.4
1	D	662	LYS	2.4
1	B	475	HIS	2.4
1	C	611	THR	2.4
1	B	719	GLU	2.3
1	A	512	LEU	2.3
1	A	507	SER	2.3
1	A	466	GLU	2.3
1	A	856	ALA	2.3
1	D	507	SER	2.3
1	D	485	ILE	2.3
1	B	492	VAL	2.3
1	D	772	VAL	2.3
1	C	462	LEU	2.3
1	C	484	LEU	2.3
1	C	822	GLY	2.3
1	A	715	LYS	2.3
1	C	474	LYS	2.3
1	C	525	ALA	2.3
1	B	721	LEU	2.3
1	B	665	GLU	2.2
1	A	467	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	829	ASP	2.2
1	A	508	SER	2.2
1	D	718	ARG	2.2
1	A	662	LYS	2.2
1	B	809	THR	2.2
1	C	524	GLY	2.2
1	C	446	GLU	2.2
1	C	494	ILE	2.2
1	B	482	GLU	2.2
1	B	832	GLY	2.2
1	C	856	ALA	2.2
1	A	511	TYR	2.2
1	B	677	GLU	2.1
1	C	675	PHE	2.1
1	A	630	ARG	2.1
1	B	533	TYR	2.1
1	C	447	GLU	2.1
1	A	506	PRO	2.1
1	A	713	PRO	2.1
1	B	494	ILE	2.1
1	A	492	VAL	2.1
1	C	488	HIS	2.1
1	D	598	ARG	2.1
1	B	463	SER	2.1
1	B	624	SER	2.1
1	C	508	SER	2.1
1	D	596	LEU	2.1
1	D	574	GLY	2.1
1	D	715	LYS	2.1
1	B	631	LEU	2.1
1	B	545	CYS	2.1
1	A	787	THR	2.1
1	D	487	THR	2.1
1	A	822	GLY	2.1
1	B	660	ILE	2.1
1	C	633	LYS	2.1
1	A	821	LEU	2.0
1	A	464	ASP	2.0
1	B	515	ARG	2.0
1	A	460	LYS	2.0
1	B	716	VAL	2.0
1	B	657	MET	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	503	LEU	2.0
1	C	516	ASP	2.0
1	A	519	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	882	A	876	42/42	0.89	0.10	24,35,41,44	0
2	882	C	876	42/42	0.90	0.10	27,35,37,38	0
2	882	A	877	42/42	0.91	0.09	25,31,35,39	0
2	882	D	876	42/42	0.95	0.07	21,27,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 882 A 876:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 882 C 876:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 882 A 877:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 882 D 876:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.