



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:57 AM UTC

PDB ID : 2Q1U / pdb_00002q1u
Title : Crystal structure of the Bordetella bronchiseptica enzyme WbmF in complex with NAD⁺ and UDP
Authors : Harmer, N.J.; King, J.D.; Palmer, C.M.; Maskell, D.; Blundell, T.L.
Deposited on : 2007-05-25
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	11% 85% • 11%
1	B	377	17% 84% 5% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative nucleotide sugar epimerase/ dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	7	0
			2585	1645	441	493	6			
1	B	335	Total	C	N	O	S	0	6	0
			2569	1629	442	492	6			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O87989
A	-18	GLY	-	expression tag	UNP O87989
A	-17	SER	-	expression tag	UNP O87989
A	-16	SER	-	expression tag	UNP O87989
A	-15	HIS	-	expression tag	UNP O87989
A	-14	HIS	-	expression tag	UNP O87989
A	-13	HIS	-	expression tag	UNP O87989
A	-12	HIS	-	expression tag	UNP O87989
A	-11	HIS	-	expression tag	UNP O87989
A	-10	HIS	-	expression tag	UNP O87989
A	-9	SER	-	expression tag	UNP O87989
A	-8	SER	-	expression tag	UNP O87989
A	-7	GLY	-	expression tag	UNP O87989
A	-6	LEU	-	expression tag	UNP O87989
A	-5	VAL	-	expression tag	UNP O87989
A	-4	PRO	-	expression tag	UNP O87989
A	-3	ARG	-	expression tag	UNP O87989
A	-2	GLY	-	expression tag	UNP O87989
A	-1	SER	-	expression tag	UNP O87989
A	0	HIS	-	expression tag	UNP O87989
B	-19	MET	-	expression tag	UNP O87989
B	-18	GLY	-	expression tag	UNP O87989
B	-17	SER	-	expression tag	UNP O87989
B	-16	SER	-	expression tag	UNP O87989
B	-15	HIS	-	expression tag	UNP O87989

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP O87989
B	-13	HIS	-	expression tag	UNP O87989
B	-12	HIS	-	expression tag	UNP O87989
B	-11	HIS	-	expression tag	UNP O87989
B	-10	HIS	-	expression tag	UNP O87989
B	-9	SER	-	expression tag	UNP O87989
B	-8	SER	-	expression tag	UNP O87989
B	-7	GLY	-	expression tag	UNP O87989
B	-6	LEU	-	expression tag	UNP O87989
B	-5	VAL	-	expression tag	UNP O87989
B	-4	PRO	-	expression tag	UNP O87989
B	-3	ARG	-	expression tag	UNP O87989
B	-2	GLY	-	expression tag	UNP O87989
B	-1	SER	-	expression tag	UNP O87989
B	0	HIS	-	expression tag	UNP O87989

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 21	C 9	N 2	O 9	P 1	0	0
3	B	1	Total 21	C 9	N 2	O 9	P 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	260	Total 260	O 260	0	0
5	B	235	Total 235	O 235	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative nucleotide sugar epimerase/ dehydratase



- Molecule 1: Putative nucleotide sugar epimerase/ dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.85Å 79.68Å 84.95Å 90.00° 113.10° 90.00°	Depositor
Resolution (Å)	50.00 – 1.70 50.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (50.00-1.70) 98.5 (50.00-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.177 , 0.211 0.176 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5791	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 86.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0565e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, UDP, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/2664	0.89	1/3625 (0.0%)
1	B	0.79	0/2645	0.88	3/3602 (0.1%)
All	All	0.84	0/5309	0.89	4/7227 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	159	SER	N-CA-C	7.12	114.63	108.22
1	A	299	ASP	N-CA-C	6.68	118.64	111.36
1	B	291[A]	ASP	N-CA-C	5.75	118.53	108.52
1	B	291[B]	ASP	N-CA-C	5.75	118.53	108.52

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	56[A]	ASP	Mainchain
1	A	56[B]	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2585	0	2505	13	0
1	B	2569	0	2479	11	0
2	A	44	0	26	2	0
2	B	44	0	26	2	0
3	A	21	0	11	0	0
3	B	21	0	11	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	260	0	0	3	1
5	B	235	0	0	0	0
All	All	5791	0	5074	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:PRO:HD3	5:A:1061:HOH:O	0.97	1.12
1:A:202:ARG:HD3	5:A:1058:HOH:O	1.73	0.89
1:B:157:ASN:OD1	1:B:166:ILE:HD12	2.02	0.59
1:B:16[B]:MET:HE2	1:B:43:VAL:HG21	1.87	0.57
1:A:57:HIS:ND1	1:A:58:PRO:HD2	2.21	0.56
1:B:344:ILE:O	1:B:347:LYS:HG2	2.07	0.54
1:A:119:HIS:H	1:A:119:HIS:CD2	2.24	0.53
1:A:57:HIS:CG	1:A:58:PRO:HD2	2.44	0.53
1:A:188:PHE:O	2:A:601:NAD:H5N	2.10	0.51
1:A:193:GLY:HA2	1:A:330:THR:OG1	2.15	0.47
1:A:103:HIS:CE1	1:B:103:HIS:CE1	3.06	0.44
1:A:189:GLN:HA	2:A:601:NAD:H4N	2.00	0.44
1:A:57:HIS:ND1	1:A:58:PRO:CD	2.81	0.43
1:A:103:HIS:NE2	1:B:103:HIS:NE2	2.62	0.43
1:B:188:PHE:O	2:B:602:NAD:H5N	2.19	0.42
1:A:55:PRO:CD	5:A:1061:HOH:O	1.86	0.42
1:A:151:ILE:CG2	1:B:151:ILE:CG2	2.98	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:TRP:CE2	1:B:351:ALA:HB1	2.55	0.41
1:B:189:GLN:HA	2:B:602:NAD:H4N	2.03	0.41
1:B:26[B]:SER:OG	1:B:53:ASN:HB3	2.20	0.40
1:B:93:GLN:HE21	1:B:206:THR:CB	2.33	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:990:HOH:O	5:A:1059:HOH:O[2_556]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/377 (90%)	335 (98%)	5 (2%)	0	100	100
1	B	338/377 (90%)	334 (99%)	4 (1%)	0	100	100
All	All	678/754 (90%)	669 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/315 (85%)	265 (99%)	2 (1%)	76	69
1	B	267/315 (85%)	266 (100%)	1 (0%)	84	80
All	All	534/630 (85%)	531 (99%)	3 (1%)	78	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	THR
1	A	119	HIS
1	B	35	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	119	HIS
1	A	180	GLN
1	B	38	ASN
1	B	92	ASN
1	B	93	GLN
1	B	213	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	802	-	5,5,5	0.43	0	5,5,5	0.81	0
2	NAD	B	602	-	46,48,48	1.62	5 (10%)	64,73,73	1.51	9 (14%)
3	UDP	B	702	-	22,22,26	0.94	1 (4%)	32,33,40	1.58	4 (12%)
3	UDP	A	701	-	22,22,26	0.93	1 (4%)	32,33,40	1.71	4 (12%)
4	GOL	A	801	-	5,5,5	0.49	0	5,5,5	0.76	0
2	NAD	A	601	-	46,48,48	1.61	3 (6%)	64,73,73	1.60	15 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	802	-	-	0/4/4/4	-
2	NAD	B	602	-	-	7/30/62/62	0/5/5/5
3	UDP	B	702	-	-	0/10/26/32	0/2/2/2
3	UDP	A	701	-	-	0/10/26/32	0/2/2/2
4	GOL	A	801	-	-	0/4/4/4	-
2	NAD	A	601	-	-	6/30/62/62	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	602	NAD	O7N-C7N	7.74	1.38	1.24
2	A	601	NAD	O7N-C7N	7.49	1.38	1.24
2	A	601	NAD	C2N-N1N	3.68	1.39	1.35
2	B	602	NAD	PN-O3	-3.24	1.56	1.59
2	B	602	NAD	C2A-N3A	3.09	1.39	1.33
2	B	602	NAD	C2A-N1A	2.25	1.37	1.33
2	A	601	NAD	C4A-N3A	2.21	1.38	1.34
2	B	602	NAD	PA-O2A	-2.14	1.45	1.55
3	B	702	UDP	C6-C5	2.11	1.40	1.35
3	A	701	UDP	C6-C5	2.06	1.39	1.35

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	UDP	C4-N3-C2	-5.30	120.03	126.61
2	B	602	NAD	C3N-C7N-N7N	5.05	123.97	117.74
3	B	702	UDP	C4-N3-C2	-5.00	120.40	126.61
3	A	701	UDP	C5-C4-N3	4.24	120.73	114.80
3	B	702	UDP	C5-C4-N3	4.23	120.72	114.80
2	A	601	NAD	C6N-N1N-C2N	-3.92	118.55	121.88
3	A	701	UDP	N3-C2-N1	3.72	119.73	114.89
2	A	601	NAD	N3A-C2A-N1A	-3.57	123.17	128.58
2	A	601	NAD	C3N-C7N-N7N	3.57	122.14	117.74
2	B	602	NAD	N3A-C2A-N1A	-3.52	123.26	128.58
2	B	602	NAD	O7N-C7N-C3N	-3.50	115.31	119.60
3	A	701	UDP	O2-C2-N3	-3.30	115.41	121.49
2	A	601	NAD	C4D-O4D-C1D	-3.21	106.98	109.92
2	A	601	NAD	C5A-N7A-C8A	3.18	108.45	103.45
2	A	601	NAD	N9A-C8A-N7A	-3.18	109.43	113.94
2	B	602	NAD	C5A-C4A-N3A	-3.07	122.49	126.72
2	A	601	NAD	C4A-C5A-N7A	-3.04	107.11	110.58
2	B	602	NAD	C4A-C5A-N7A	-2.92	107.24	110.58
2	B	602	NAD	C5A-N7A-C8A	2.91	108.03	103.45
3	B	702	UDP	N3-C2-N1	2.82	118.56	114.89
2	A	601	NAD	O7N-C7N-C3N	-2.81	116.17	119.60
2	B	602	NAD	C6A-C5A-C4A	2.71	120.88	117.18
2	A	601	NAD	C5A-C4A-N3A	-2.71	122.99	126.72
2	B	602	NAD	N9A-C8A-N7A	-2.65	110.18	113.94
3	B	702	UDP	O4-C4-C5	-2.63	120.64	125.16
2	A	601	NAD	C5N-C4N-C3N	-2.49	117.92	120.36
2	A	601	NAD	C6N-C5N-C4N	2.33	122.81	119.45
2	B	602	NAD	C5A-C4A-N9A	2.31	108.33	105.81
2	A	601	NAD	C6A-C5A-C4A	2.29	120.30	117.18
2	A	601	NAD	C2A-N1A-C6A	2.24	122.42	118.73
2	A	601	NAD	O2N-PN-O3	2.11	112.98	107.27
2	A	601	NAD	C5A-C4A-N9A	2.01	108.00	105.81

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	NAD	C5D-O5D-PN-O3
2	A	601	NAD	C5D-O5D-PN-O1N
2	A	601	NAD	C5D-O5D-PN-O2N
2	B	602	NAD	C5D-O5D-PN-O3

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	B	602	NAD	C5D-O5D-PN-O1N
2	B	602	NAD	C5D-O5D-PN-O2N
2	A	601	NAD	PN-O3-PA-O5B
2	B	602	NAD	PN-O3-PA-O5B
2	A	601	NAD	PA-O3-PN-O1N
2	B	602	NAD	PA-O3-PN-O2N
2	A	601	NAD	PA-O3-PN-O2N
2	B	602	NAD	PA-O3-PN-O1N
2	B	602	NAD	O4B-C4B-C5B-O5B



There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	602	NAD	2	0
2	A	601	NAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	336/377 (89%)	0.86	42 (12%)	8 8	5, 18, 28, 52	7 (2%)
1	B	335/377 (88%)	1.07	64 (19%)	3 3	5, 20, 33, 46	6 (1%)
All	All	671/754 (88%)	0.97	106 (15%)	5 4	5, 20, 31, 52	13 (1%)

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	ASN	6.3
1	A	7	ALA	5.4
1	B	58	PRO	5.4
1	B	7	ALA	5.1
1	B	353	ALA	5.0
1	B	156	ASN	5.0
1	B	132	GLY	4.9
1	A	119	HIS	4.7
1	B	326	GLY	4.4
1	A	350	SER	4.3
1	A	57	HIS	4.3
1	A	59	ALA	4.2
1	B	340	VAL	4.1
1	A	156	ASN	4.1
1	B	290	LEU	4.1
1	A	302	GLY	4.0
1	B	339	ALA	3.9
1	B	254	ALA	3.9
1	B	355	TYR	3.7
1	B	52	ILE	3.7
1	A	351	ALA	3.7
1	A	353	ALA	3.7
1	B	8	SER	3.5
1	B	37	VAL	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	59	ALA	3.3
1	A	301	SER	3.2
1	B	47	LEU	3.2
1	B	298	TRP	3.1
1	A	268	GLY	3.1
1	B	305	PHE	3.1
1	A	58	PRO	3.0
1	B	260	GLY	3.0
1	A	255	ALA	3.0
1	B	261	GLY	3.0
1	B	307	SER	2.9
1	B	11	ALA	2.9
1	A	348	HIS	2.9
1	B	33	LEU	2.8
1	B	12	ASN	2.8
1	A	243	VAL	2.8
1	A	307	SER	2.8
1	A	309[A]	GLU	2.8
1	A	311	ALA	2.8
1	A	60	VAL	2.8
1	A	312	ARG	2.8
1	B	243	VAL	2.8
1	B	256	ASP	2.7
1	A	62	PHE	2.6
1	A	242	PHE	2.6
1	A	288	THR	2.6
1	B	36	GLY	2.6
1	B	56	ASP	2.6
1	B	291[A]	ASP	2.6
1	B	13	THR	2.6
1	B	57	HIS	2.6
1	B	77	GLN	2.5
1	A	352	LEU	2.5
1	A	355	TYR	2.5
1	B	38	ASN	2.5
1	B	55	PRO	2.5
1	B	131	ALA	2.5
1	B	229	PRO	2.5
1	B	81	ASP	2.5
1	B	247	ALA	2.5
1	B	119	HIS	2.5
1	B	321	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	344	ILE	2.5
1	B	34	GLU	2.5
1	A	298	TRP	2.4
1	B	309	GLU	2.4
1	B	351	ALA	2.4
1	B	232	ASN	2.4
1	B	333	TRP	2.4
1	B	320	ASP	2.3
1	A	340	VAL	2.3
1	B	35	LEU	2.3
1	B	352	LEU	2.3
1	A	305	PHE	2.3
1	A	8	SER	2.3
1	B	285	GLY	2.3
1	B	230	LEU	2.3
1	A	308	PRO	2.2
1	B	21	ALA	2.2
1	A	24	VAL	2.2
1	A	229	PRO	2.2
1	B	258	THR	2.2
1	B	322	SER	2.2
1	B	72	LEU	2.2
1	A	339	ALA	2.2
1	B	10	LEU	2.2
1	B	315	LEU	2.2
1	B	350	SER	2.2
1	A	230	LEU	2.1
1	A	67	ILE	2.1
1	B	257	GLY	2.1
1	B	302	GLY	2.1
1	A	232	ASN	2.1
1	A	344	ILE	2.1
1	B	300	ASN	2.1
1	B	327	LEU	2.1
1	A	317	PHE	2.1
1	B	80	TYR	2.1
1	B	311	ALA	2.0
1	A	9	LYS	2.0
1	A	12	ASN	2.0
1	A	320	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UDP	A	701	21/25	0.82	0.15	36,39,55,56	0
3	UDP	B	702	21/25	0.82	0.14	42,44,57,57	0
4	GOL	B	802	6/6	0.93	0.11	26,31,33,34	0
4	GOL	A	801	6/6	0.94	0.11	23,28,32,33	0
2	NAD	A	601	44/44	0.95	0.10	14,17,22,23	0
2	NAD	B	602	44/44	0.95	0.10	16,19,23,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around UDP A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UDP B 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.