



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 09:31 AM UTC

PDB ID : 7Q13 / pdb_00007q13
EMDB ID : EMD-13753
Title : Human GYS1-GYG1 complex activated state bound to glucose-6-phosphate, uridine diphosphate, and glucose
Authors : McCorvie, T.J.; Shrestha, L.; Froese, D.S.; Ferreira, I.M.; Yue, W.W.
Deposited on : 2021-10-17
Resolution : 3.00 Å (reported)
Based on initial models : 4QLB, 3NB0

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	737	6% 60% 20% 20%
1	B	737	7% 60% 20% 20%
1	C	737	6% 59% 21% 20%
1	D	737	6% 60% 20% 20%
2	E	350	7% 7% 91%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	350	 6% 8% 91%
2	G	350	 7% 8% 91%
2	H	350	 7% 7% 91%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G6P	A	801	X	-	-	-
3	G6P	B	801	X	-	-	-
3	G6P	C	801	X	-	-	-
3	G6P	D	801	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycogen [starch] synthase, muscle.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	590	Total	C	N	O	S	1	0
			4767	3061	825	855	26		
1	B	590	Total	C	N	O	S	1	0
			4767	3061	825	855	26		
1	C	590	Total	C	N	O	S	1	0
			4767	3061	825	855	26		
1	D	590	Total	C	N	O	S	1	0
			4767	3061	825	855	26		

- Molecule 2 is a protein called Glycogenin-1.


Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	32	Total	C	N	O	S	0	0
			272	167	48	56	1		
2	F	32	Total	C	N	O	S	0	0
			272	167	48	56	1		
2	G	32	Total	C	N	O	S	0	0
			272	167	48	56	1		
2	H	32	Total	C	N	O	S	0	0
			272	167	48	56	1		

- Molecule 3 is 6-O-phosphono-alpha-D-glucopyranose (CCD ID: G6P) (formula: C₆H₁₃O₉P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	O	P	0
			16	6	9	1	
3	B	1	Total	C	O	P	0
			16	6	9	1	
3	C	1	Total	C	O	P	0
			16	6	9	1	
3	D	1	Total	C	O	P	0
			16	6	9	1	

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	N	O	P	0
			25	9	2	12	2	
4	B	1	Total	C	N	O	P	0
			25	9	2	12	2	
4	C	1	Total	C	N	O	P	0
			25	9	2	12	2	
4	D	1	Total	C	N	O	P	0
			25	9	2	12	2	

- Molecule 5 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
5	A	1	Total	C	O	0
			12	6	6	
5	B	1	Total	C	O	0
			12	6	6	
5	C	1	Total	C	O	0
			12	6	6	
5	D	1	Total	C	O	0
			12	6	6	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen [starch] synthase, muscle





- Molecule 1: Glycogen [starch] synthase, muscle






• Molecule 1: Glycogen [starch] synthase, muscle







• Molecule 2: Glycogenin-1



• Molecule 2: Glycogenin-1



• Molecule 2: Glycogenin-1



SER	THR	LYS	SER	VAL
HIS	HIS	HIS	ALA	VAL
LEU	PRO	LEU	ALA	VAL
LEU	GLU	PRO	PRO	LEU
LEU	PHE	PHE	ASP	ASP
GLY	LEU	ILE	PRO	SER
GLU	ILE	TYR	GLY	GLY
ILE	LEU	ASN	TRP	ASP
PRO	TRP	LEU	PRO	SER
ALA	TRP	SER	ASP	ALA
MET	ASN	SER	CYS	HIS
ALA	ILE	ILE	PHE	LEU
GLN	PHE	SER	ASN	THR
PRO	THR	ILE	SER	LEU
PHE	THR	TYR	GLY	MET
VAL	ASN	SER	LYS	LYS
SER	VAL	TYR	PHE	ARG
S318	LEU	LEU	VAL	PRO
E319	PRO	PRO	TYR	GLU
E320	LEU	ALA	GLN	LEU
R321	LEU	PHE	PRO	GLY
K322	GLN	LYS	SER	VAL
E323	PHE	PHE	VAL	THR
R324	GLY	GLY	GLU	LEU
W325	LEU	ALA	TYR	LEU
E326	LYS	ALA	GLN	LYS
Q327	ASP	LYS	LEU	CYS
Q328	THR	VAL	LEU	HIS
Q329	CYS	VAL	SER	SER
A330	TYR	PHE	LEU	LEU
D331	VAL	LEU	SER	THR
Y332	ASN	GLY	GLU	TYR
M333	VAL	ARG	GLN	SER
G334	LEU	VAL	LYS	LYS
A335	ASP	PRO	SER	CYS
D336	LEU	TRP	PHE	VAL
S337	VAL	ASN	ASP	PHE
F338	TYR	TYR	GLY	MET
D339	THR	TYR	GLY	ASP
M340	LEU	THR	ASP	ALA
I341	ALA	TYR	GLN	ASP
I342	PHE	ASP	GLY	THR
K342	SER	PRO	ILE	LEU
R343	SER	LYS	LEU	VAL
D346	CYS	THR	ASN	LEU
T347	GLY	LYS	THR	LEU
Y348	PHE	SER	PHE	ALA
I349	CYS	VAL	ILE	ASP
GLN	ARG	LYS	SER	PHE
	GLU	SER	SER	LEU
	ASP	GLU	TRP	LEU
	VAL	ALA	ALA	PHE
	THR	HIS	THR	ASP
	SER	ASP	THR	ARG
	GLY	PRO	ASP	GLU
	ILE	ASN	ILE	GLU
	THR	MET	ASP	LEU

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D2	Depositor
Number of particles used	35604	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2300	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.363	Depositor
Minimum map value	-0.190	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.03	Depositor
Map size (Å)	339.19998, 339.19998, 339.19998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, GLC, G6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/4896	0.36	0/6635
1	B	0.17	0/4896	0.36	0/6635
1	C	0.17	0/4896	0.36	0/6635
1	D	0.17	0/4896	0.36	0/6635
2	E	0.09	0/276	0.24	0/367
2	F	0.09	0/276	0.23	0/367
2	G	0.09	0/276	0.23	0/367
2	H	0.09	0/276	0.23	0/367
All	All	0.17	0/20688	0.36	0/28008

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4767	0	4677	111	0
1	B	4767	0	4677	110	0
1	C	4767	0	4677	111	0
1	D	4767	0	4677	107	0
2	E	272	0	251	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	272	0	251	5	0
2	G	272	0	251	6	0
2	H	272	0	251	6	0
3	A	16	0	11	1	0
3	B	16	0	11	1	0
3	C	16	0	11	1	0
3	D	16	0	11	1	0
4	A	25	0	11	1	0
4	B	25	0	11	1	0
4	C	25	0	11	1	0
4	D	25	0	11	1	0
5	A	12	0	12	1	0
5	B	12	0	12	1	0
5	C	12	0	12	1	0
5	D	12	0	12	1	0
All	All	20368	0	19848	427	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 427 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:LYS:HG3	1:A:102:TYR:HE1	1.47	0.79
1:C:100:LYS:HG3	1:C:102:TYR:HE1	1.47	0.79
1:B:100:LYS:HG3	1:B:102:TYR:HE1	1.48	0.79
1:D:100:LYS:HG3	1:D:102:TYR:HE1	1.47	0.79
1:B:209:LEU:HD22	1:B:269:LEU:HD11	1.65	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/737 (80%)	556 (94%)	33 (6%)	0	100	100
1	B	589/737 (80%)	556 (94%)	33 (6%)	0	100	100
1	C	589/737 (80%)	557 (95%)	32 (5%)	0	100	100
1	D	589/737 (80%)	557 (95%)	32 (5%)	0	100	100
2	E	30/350 (9%)	29 (97%)	1 (3%)	0	100	100
2	F	30/350 (9%)	29 (97%)	1 (3%)	0	100	100
2	G	30/350 (9%)	29 (97%)	1 (3%)	0	100	100
2	H	30/350 (9%)	29 (97%)	1 (3%)	0	100	100
All	All	2476/4348 (57%)	2342 (95%)	134 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	UDP	B	802	-	25,26,26	1.24	3 (12%)	38,40,40	1.66	5 (13%)
4	UDP	A	802	-	25,26,26	1.25	3 (12%)	38,40,40	1.64	5 (13%)
3	G6P	A	801	-	16,16,16	1.16	2 (12%)	23,24,24	0.78	0
5	GLC	D	803	-	12,12,12	0.47	0	17,17,17	0.83	0
5	GLC	A	803	-	12,12,12	0.46	0	17,17,17	0.83	0
4	UDP	D	802	-	25,26,26	1.25	3 (12%)	38,40,40	1.64	5 (13%)
5	GLC	C	803	-	12,12,12	0.46	0	17,17,17	0.84	0
3	G6P	C	801	-	16,16,16	1.16	2 (12%)	23,24,24	0.79	0
5	GLC	B	803	-	12,12,12	0.46	0	17,17,17	0.84	0
4	UDP	C	802	-	25,26,26	1.26	3 (12%)	38,40,40	1.65	5 (13%)
3	G6P	D	801	-	16,16,16	1.16	2 (12%)	23,24,24	0.78	0
3	G6P	B	801	-	16,16,16	1.16	2 (12%)	23,24,24	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UDP	B	802	-	-	4/16/32/32	0/2/2/2
4	UDP	A	802	-	-	3/16/32/32	0/2/2/2
3	G6P	A	801	-	4/4/6/6	4/6/26/26	0/1/1/1
5	GLC	D	803	-	-	2/2/22/22	0/1/1/1
5	GLC	A	803	-	-	2/2/22/22	0/1/1/1
4	UDP	D	802	-	-	3/16/32/32	0/2/2/2
5	GLC	C	803	-	-	2/2/22/22	0/1/1/1
3	G6P	C	801	-	4/4/6/6	3/6/26/26	0/1/1/1
5	GLC	B	803	-	-	2/2/22/22	0/1/1/1
4	UDP	C	802	-	-	4/16/32/32	0/2/2/2
3	G6P	D	801	-	4/4/6/6	4/6/26/26	0/1/1/1
3	G6P	B	801	-	4/4/6/6	3/6/26/26	0/1/1/1

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	802	UDP	C4-N3	-2.94	1.33	1.38
4	A	802	UDP	C4-N3	-2.93	1.33	1.38
4	D	802	UDP	C4-N3	-2.91	1.33	1.38
4	B	802	UDP	C4-N3	-2.88	1.33	1.38
4	C	802	UDP	C2-N3	-2.67	1.33	1.38

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	802	UDP	C4-N3-C2	-5.21	120.14	126.61
4	B	802	UDP	C4-N3-C2	-5.20	120.16	126.61
4	C	802	UDP	C4-N3-C2	-5.17	120.19	126.61
4	A	802	UDP	C4-N3-C2	-5.17	120.20	126.61
4	B	802	UDP	N3-C2-N1	4.37	120.58	114.89

5 of 16 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	801	G6P	C5
3	A	801	G6P	C4
3	A	801	G6P	C1
3	A	801	G6P	C3
3	B	801	G6P	C5

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	G6P	C4-C5-C6-O6
3	A	801	G6P	O5-C5-C6-O6
3	A	801	G6P	C6-O6-P-O3P
3	B	801	G6P	C4-C5-C6-O6
3	B	801	G6P	O5-C5-C6-O6

There are no ring outliers.

12 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	802	UDP	1	0
4	A	802	UDP	1	0
3	A	801	G6P	1	0
5	D	803	GLC	1	0
5	A	803	GLC	1	0



Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	802	UDP	1	0
5	C	803	GLC	1	0
3	C	801	G6P	1	0
5	B	803	GLC	1	0
4	C	802	UDP	1	0
3	D	801	G6P	1	0
3	B	801	G6P	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13753. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 132




Y Index: 145




Z Index: 159


6.3.2 Raw map



X Index: 132



Y Index: 145




Z Index: 159


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_13753_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 162 nm³; this corresponds to an approximate mass of 147 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	3.00	3.43	3.06
Unmasked-calculated*	3.46	5.89	3.59

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.46 differs from the reported value 3.0 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-13753 and PDB model 7Q13. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7890	0.4350
A	0.8190	0.4490
B	0.8140	0.4470
C	0.8170	0.4500
D	0.8190	0.4510
E	0.2930	0.1820
F	0.2550	0.1710
G	0.3000	0.1930
H	0.2590	0.1630

1.0

0.0

<0.0