



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 11:18 AM UTC

PDB ID : 8Q17 / pdb_00008q17
Title : Identification and optimisation of novel inhibitors of the Polyketide synthetase
13 thioesterase domain with antitubercular activity
Authors : Eadsforth, T.C.; Puneekar, A.S.; Green, S.R.; Baragana, B.
Deposited on : 2023-07-30
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	8% 88% 8% ..
1	B	286	12% 90% 6% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Polyketide synthase Pks13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	1	0
			2175	1381	382	406	6			
1	B	276	Total	C	N	O	S	0	3	0
			2176	1378	383	409	6			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1448	SER	-	expression tag	UNP I6X8D2
A	1449	ASN	-	expression tag	UNP I6X8D2
A	1450	ALA	-	expression tag	UNP I6X8D2
B	1448	SER	-	expression tag	UNP I6X8D2
B	1449	ASN	-	expression tag	UNP I6X8D2
B	1450	ALA	-	expression tag	UNP I6X8D2

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is {N}-[2-[4-(azetidin-1-ylcarbonyl)phenyl]ethyl]-3-(3,4-dimethoxyphenyl)-1,2,4-oxadiazole-5-carboxamide (CCD ID: IL8) (formula: C₂₃H₂₄N₄O₅) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 32 23 4 5	0	0
3	B	1	Total C N O 32 23 4 5	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (2R)-2-{[(2R)-2-{[(2S)-2-{[(2R)-2-hydroxypropyl]oxy}propyl]oxy}propyl]oxy}propan-1-ol (CCD ID: 2Q5) (formula: C₁₂H₂₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			17	12	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	179	Total 179	O 179	0	0
6	B	161	Total 161	O 161	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polyketide synthase Pks13



• Molecule 1: Polyketide synthase Pks13



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.45Å 109.85Å 57.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.45 – 1.71 68.89 – 1.71	Depositor EDS
% Data completeness (in resolution range)	91.3 (88.45-1.71) 91.3 (68.89-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.195 , 0.239 0.207 , 0.247	Depositor DCC
R_{free} test set	2738 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4809	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 2Q5, SO4, IL8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	5/2228 (0.2%)	1.17	5/3025 (0.2%)
1	B	1.30	8/2227 (0.4%)	1.15	4/3022 (0.1%)
All	All	1.28	13/4455 (0.3%)	1.16	9/6047 (0.1%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1458	THR	CA-CB	6.85	1.61	1.53
1	B	1510	ARG	C-O	6.32	1.31	1.24
1	B	1552	ASP	CA-C	5.68	1.60	1.53
1	B	1500	PHE	C-O	5.62	1.30	1.24
1	B	1592	VAL	N-CA	5.47	1.53	1.46
1	A	1463	PRO	N-CA	5.35	1.53	1.47
1	B	1667	ALA	N-CA	5.32	1.52	1.46
1	B	1515	VAL	CA-C	5.29	1.57	1.52
1	B	1694	PRO	C-O	-5.19	1.17	1.23
1	B	1638	LEU	N-CA	-5.13	1.39	1.46
1	A	1689	ASP	CA-C	5.10	1.58	1.53
1	A	1703	ILE	CA-C	5.08	1.58	1.52
1	A	1519	ILE	CA-C	5.01	1.59	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1502	ARG	CG-CD-NE	-6.25	98.24	112.00
1	A	1641	ARG	N-CA-C	5.39	118.08	111.82
1	B	1525	GLY	N-CA-C	-5.32	101.49	112.34
1	A	1495	THR	CA-C-N	-5.25	115.16	120.31
1	A	1495	THR	C-N-CA	-5.25	115.16	120.31
1	B	1524	ASP	N-CA-C	5.21	117.86	111.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1521	MET	N-CA-C	5.08	118.63	112.23
1	A	1563	ARG	CG-CD-NE	5.03	123.05	112.00
1	B	1554	ARG	CG-CD-NE	-5.00	100.99	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2175	0	2127	7	0
1	B	2176	0	2122	5	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0
3	A	32	0	0	0	0
3	B	32	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	B	17	0	26	0	0
6	A	179	0	0	2	0
6	B	161	0	0	0	0
All	All	4809	0	4291	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1591:ASN:O	1:B:1592:VAL:HG22	1.99	0.62
1:A:1581:ARG:HD2	6:A:2066:HOH:O	2.08	0.53
1:A:1611:VAL:HG11	1:A:1634:ARG:HG3	1.94	0.49
1:B:1587:GLU:O	1:B:1591:ASN:HA	2.13	0.49
1:A:1615:LEU:HD22	1:A:1630:ILE:HD12	1.96	0.47
1:A:1659:MET:HE1	1:A:1679:PRO:O	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1552:ASP:OD1	1:B:1554:ARG:NH1	2.49	0.45
1:B:1726[B]:ARG:HH11	1:B:1726[B]:ARG:HG2	1.81	0.45
1:A:1524:ASP:HA	1:A:1551:LYS:HD2	2.00	0.43
1:B:1588:LYS:O	1:B:1591:ASN:N	2.54	0.41
1:A:1581:ARG:HD2	6:A:2012:HOH:O	2.21	0.41
1:A:1526:PRO:HB3	1:A:1554:ARG:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/286 (96%)	270 (98%)	6 (2%)	0	100	100
1	B	275/286 (96%)	266 (97%)	7 (2%)	2 (1%)	18	7
All	All	551/572 (96%)	536 (97%)	13 (2%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1592	VAL
1	B	1451	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/231 (96%)	215 (97%)	7 (3%)	34	11
1	B	222/231 (96%)	218 (98%)	4 (2%)	51	29
All	All	444/462 (96%)	433 (98%)	11 (2%)	42	18

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1452	ILE
1	A	1533	SER
1	A	1563	ARG
1	A	1621	SER
1	A	1624	GLN
1	A	1659	MET
1	A	1669	MET
1	B	1554	ARG
1	B	1597	ILE
1	B	1602	LEU
1	B	1713	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1512	GLN
1	B	1624	GLN
1	B	1647	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SO4	A	1803	-	4,4,4	0.57	0	6,6,6	0.37	0
3	IL8	A	1804	-	35,35,35	2.08	10 (28%)	45,48,48	2.45	13 (28%)
5	2Q5	B	1805	-	16,16,16	1.22	2 (12%)	19,19,19	1.91	7 (36%)
3	IL8	B	1803	-	35,35,35	2.21	11 (31%)	45,48,48	2.38	14 (31%)
2	SO4	B	1802	-	4,4,4	0.41	0	6,6,6	0.78	0
4	GOL	B	1804	-	5,5,5	0.14	0	5,5,5	0.62	0
2	SO4	A	1802	-	4,4,4	0.66	0	6,6,6	0.44	0
2	SO4	A	1801	-	4,4,4	0.56	0	6,6,6	0.28	0
2	SO4	B	1801	-	4,4,4	0.40	0	6,6,6	0.23	0
4	GOL	A	1805	-	5,5,5	0.74	0	5,5,5	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IL8	A	1804	-	-	3/26/32/32	0/4/4/4
5	2Q5	B	1805	-	-	9/17/17/17	-
3	IL8	B	1803	-	-	2/26/32/32	0/4/4/4
4	GOL	B	1804	-	-	2/4/4/4	-
4	GOL	A	1805	-	-	0/4/4/4	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1804	IL8	C06-C11	-6.35	1.37	1.47
3	B	1803	IL8	C06-C11	-5.63	1.38	1.47
3	B	1803	IL8	C16-N18	5.02	1.42	1.33
3	A	1804	IL8	C16-N18	4.80	1.41	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1803	IL8	O02-C03	4.58	1.44	1.37
3	B	1803	IL8	C27-N29	4.36	1.43	1.34
3	B	1803	IL8	O13-N12	-3.76	1.35	1.42
3	A	1804	IL8	O13-N12	-3.49	1.36	1.42
3	B	1803	IL8	C08-C03	-3.32	1.33	1.40
3	A	1804	IL8	O09-C08	3.23	1.42	1.37
3	A	1804	IL8	C27-N29	3.17	1.41	1.34
3	A	1804	IL8	O13-C14	-3.15	1.30	1.34
3	A	1804	IL8	O17-C16	-2.79	1.18	1.23
3	B	1803	IL8	O09-C08	2.71	1.41	1.37
5	B	1805	2Q5	C4-C5	2.66	1.56	1.50
3	A	1804	IL8	C08-C03	-2.60	1.35	1.40
3	B	1803	IL8	C30-N29	2.59	1.49	1.47
3	B	1803	IL8	O17-C16	-2.40	1.19	1.23
3	A	1804	IL8	C31-C32	-2.37	1.48	1.54
5	B	1805	2Q5	C8-C7	2.13	1.55	1.50
3	B	1803	IL8	C24-C27	2.05	1.53	1.50
3	A	1804	IL8	C32-N29	2.04	1.49	1.47
3	B	1803	IL8	C32-N29	2.03	1.49	1.47

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1803	IL8	O13-N12-C11	7.75	107.82	103.28
3	A	1804	IL8	N15-C11-N12	-6.10	110.02	114.47
3	B	1803	IL8	N15-C11-N12	-5.84	110.21	114.47
3	A	1804	IL8	C06-C11-N12	-5.76	117.31	122.22
3	A	1804	IL8	O13-N12-C11	5.56	106.54	103.28
3	A	1804	IL8	C06-C11-N15	5.54	132.61	122.92
3	A	1804	IL8	O13-C14-N15	-4.80	109.02	114.12
3	A	1804	IL8	C24-C27-N29	4.72	124.55	118.66
3	B	1803	IL8	C06-C11-N15	4.59	130.96	122.92
5	B	1805	2Q5	C8-O4-C11	4.32	121.42	115.05
3	B	1803	IL8	O17-C16-C14	-4.25	115.64	120.72
3	A	1804	IL8	C11-N15-C14	4.15	105.69	102.61
5	B	1805	2Q5	C3-O2-C5	4.11	121.12	115.05
3	B	1803	IL8	C06-C11-N12	-3.99	118.82	122.22
3	B	1803	IL8	C31-C30-N29	-3.80	85.83	88.21
3	A	1804	IL8	C10-O09-C08	-3.79	111.95	117.51
3	B	1803	IL8	O09-C08-C03	-3.35	110.86	115.40
3	B	1803	IL8	C31-C32-N29	-3.00	86.33	88.21
3	B	1803	IL8	O28-C27-N29	-2.83	117.89	122.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1803	IL8	C05-C06-C07	-2.80	116.00	119.25
5	B	1805	2Q5	C4-O3-C7	2.65	118.95	115.05
3	B	1803	IL8	C07-C06-C11	2.50	123.62	120.27
5	B	1805	2Q5	O2-C5-C4	2.50	116.95	108.90
3	A	1804	IL8	O28-C27-N29	-2.47	118.46	122.35
3	B	1803	IL8	C10-O09-C08	2.45	121.11	117.51
3	B	1803	IL8	C01-O02-C03	-2.41	113.98	117.51
5	B	1805	2Q5	O2-C3-C2	2.35	113.52	108.94
3	A	1804	IL8	C14-O13-N12	2.31	108.68	106.08
3	A	1804	IL8	C25-C24-C23	2.23	121.41	118.57
3	A	1804	IL8	C05-C06-C11	-2.22	117.98	120.78
5	B	1805	2Q5	O3-C4-C5	2.20	116.18	110.82
3	B	1803	IL8	C11-N15-C14	2.12	104.18	102.61
5	B	1805	2Q5	C6-C5-C4	-2.07	106.86	112.69
3	A	1804	IL8	O28-C27-C24	-2.05	116.23	120.29

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1804	GOL	C1-C2-C3-O3
5	B	1805	2Q5	O3-C7-C8-O4
5	B	1805	2Q5	C9-C7-C8-O4
5	B	1805	2Q5	C12-C11-O4-C8
4	B	1804	GOL	O2-C2-C3-O3
5	B	1805	2Q5	C1-C2-C3-O2
5	B	1805	2Q5	C5-C4-O3-C7
3	A	1804	IL8	C07-C06-C11-N12
5	B	1805	2Q5	O5-C10-C11-O4
5	B	1805	2Q5	O5-C10-C11-C12
3	A	1804	IL8	C07-C06-C11-N15
5	B	1805	2Q5	C2-C3-O2-C5
5	B	1805	2Q5	C7-C8-O4-C11
3	A	1804	IL8	C05-C06-C11-N12
3	B	1803	IL8	C05-C06-C11-N12
3	B	1803	IL8	C05-C06-C11-N15


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/286 (96%)	0.48	23 (8%) 17 22	16, 28, 59, 96	1 (0%)
1	B	276/286 (96%)	0.68	33 (11%) 9 13	11, 29, 71, 91	3 (1%)
All	All	553/572 (96%)	0.58	56 (10%) 12 17	11, 29, 63, 96	4 (0%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1450	ALA	5.1
1	B	1594	ILE	4.8
1	A	1623	VAL	4.8
1	B	1597	ILE	4.3
1	A	1592	VAL	4.3
1	B	1592	VAL	4.1
1	B	1727	THR	3.9
1	B	1586	ALA	3.8
1	A	1452	ILE	3.8
1	B	1623	VAL	3.7
1	B	1599	TYR	3.7
1	B	1585	PHE	3.7
1	A	1727	THR	3.6
1	A	1590	PHE	3.4
1	B	1452	ILE	3.4
1	A	1612	ARG	3.3
1	B	1726[A]	ARG	3.3
1	B	1593	THR	3.2
1	B	1598	PRO	3.2
1	A	1621	SER	3.1
1	A	1622	GLY	3.1
1	B	1596	ALA	3.1
1	B	1724	ALA	3.1
1	A	1618	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1625	ILE	2.8
1	B	1595	PRO	2.8
1	A	1594	ILE	2.8
1	B	1591	ASN	2.7
1	B	1465	ALA	2.6
1	B	1588	LYS	2.6
1	B	1466	GLY	2.6
1	B	1463	PRO	2.5
1	A	1466	GLY	2.5
1	A	1630	ILE	2.5
1	B	1602	LEU	2.5
1	A	1602	LEU	2.4
1	B	1722	ILE	2.4
1	A	1669	MET	2.4
1	A	1465	ALA	2.3
1	B	1584	ALA	2.3
1	B	1464	GLU	2.3
1	A	1613	PHE	2.3
1	B	1618	VAL	2.3
1	B	1601	GLN	2.2
1	B	1624	GLN	2.2
1	A	1595	PRO	2.2
1	B	1451	GLN	2.2
1	B	1573	GLU	2.2
1	B	1580	ASP	2.1
1	B	1669	MET	2.1
1	A	1624	GLN	2.1
1	A	1586	ALA	2.1
1	A	1593	THR	2.1
1	A	1604	GLU	2.0
1	A	1617	ALA	2.0
1	A	1614	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	B	1802	5/5	0.65	0.14	77,82,86,94	0
2	SO4	A	1803	5/5	0.66	0.16	71,89,93,95	0
2	SO4	A	1802	5/5	0.66	0.13	70,71,90,92	0
4	GOL	B	1804	6/6	0.73	0.17	52,57,62,68	0
2	SO4	A	1801	5/5	0.76	0.11	83,85,94,94	0
5	2Q5	B	1805	17/17	0.76	0.24	58,66,82,86	0
4	GOL	A	1805	6/6	0.81	0.14	51,57,59,61	0
2	SO4	B	1801	5/5	0.92	0.10	34,35,40,41	5
3	IL8	B	1803	32/32	0.93	0.08	24,28,34,35	0
3	IL8	A	1804	32/32	0.93	0.09	27,32,44,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 2Q5 B 1805:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IL8 B 1803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around IL8 A 1804:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.