



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:58 AM UTC

PDB ID : 4Q27 / pdb_00004q27
Title : Galectin-1 in Complex with a Click-Activated N-Acetylactosamine
Authors : Grimm, C.; Bertleff-Zieschang, N.
Deposited on : 2014-04-07
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.20-1.20)
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	% 81% 16% ..
1	B	136	4% 82% 12% ..
2	C	2	100%
2	D	2	100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4925 atoms, of which 2310 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galectin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	133	Total	C	H	N	O	S	0	16	0
			2257	715	1129	193	210	10			
1	B	133	Total	C	H	N	O	S	0	15	0
			2247	711	1119	195	212	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P09382
B	-1	GLY	-	expression tag	UNP P09382

- Molecule 2 is an oligosaccharide called 3-O-prop-2-yn-1-yl-beta-D-galactopyranose-(1-4)-pro p-2-en-1-yl 2-(acetylamino)-2-deoxy-beta-D-glucopyranoside.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	2	Total	C	H	N	O	0	0	0
			63	20	31	1	11			
2	D	2	Total	C	H	N	O	0	0	0
			63	20	31	1	11			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	129	Total	O	0	0
			129	129		
4	B	161	Total	O	0	0
			161	161		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.47Å 58.13Å 110.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.47 – 1.20 40.47 – 1.20	Depositor EDS
% Data completeness (in resolution range)	96.3 (40.47-1.20) 96.3 (40.47-1.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 1.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.151 , 0.180 0.150 , 0.177	Depositor DCC
R_{free} test set	2000 reflections (1.96%)	wwPDB-VP
Wilson B-factor (Å ²)	11.5	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4925	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CME, TVV, TVS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	0/1158	0.97	1/1564 (0.1%)
1	B	1.19	1/1145 (0.1%)	0.97	0/1543
All	All	1.17	1/2303 (0.0%)	0.97	1/3107 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	35	GLY	N-CA	-5.17	1.40	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	78	PRO	CA-C-O	-5.82	116.90	121.38

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	25	PRO	Peptide,Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1128	1129	1084	26	1
1	B	1128	1119	1085	26	0
2	C	32	31	0	0	0
2	D	32	31	0	0	0
3	B	5	0	0	0	0
4	A	129	0	0	3	0
4	B	161	0	0	8	1
All	All	2615	2310	2169	49	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100[B]:LEU:HD12	1:A:104:TYR:HD2	1.33	0.92
1:B:26[A]:ASP:OD1	4:B:406:HOH:O	1.91	0.88
1:A:63:LYS:NZ	4:A:423:HOH:O	2.07	0.86
1:A:100[B]:LEU:HD12	1:A:104:TYR:CD2	2.16	0.81
1:A:16:CME:SD	1:A:90[B]:THR:HG23	2.25	0.77
1:B:20[B]:ARG:NE	1:B:130:CME:OH	2.21	0.73
1:B:74:GLU:OE2	4:B:433:HOH:O	2.09	0.70
1:A:99[B]:LYS:CA	1:A:100[B]:LEU:N	2.57	0.67
1:A:76[A]:VAL:HG22	1:A:104:TYR:CE2	2.31	0.66
1:A:76[A]:VAL:HG21	1:A:104:TYR:CG	2.31	0.66
1:B:2:CYS:O	4:B:421:HOH:O	2.13	0.65
1:A:98:VAL:C	1:A:99[B]:LYS:CA	2.69	0.65
1:B:28:LYS:HD2	4:B:430:HOH:O	1.98	0.62
1:B:2:CYS:N	4:B:432:HOH:O	2.33	0.62
1:B:32[B]:LEU:HD11	1:B:120:MET:HG3	1.81	0.62
1:A:9:LEU:HD22	1:A:133:PHE:CE1	2.35	0.61
1:A:76[A]:VAL:HG21	1:A:104:TYR:CD2	2.36	0.61
1:B:93[B]:GLN:H	1:B:93[B]:GLN:CD	2.07	0.61
1:A:76[A]:VAL:CG2	1:A:104:TYR:CE2	2.85	0.60
1:A:76[A]:VAL:CG2	1:A:104:TYR:CD2	2.85	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86:GLU:OE2	1:B:130:CME:OH	2.20	0.60
1:A:131[B]:VAL:CG2	1:B:131[B]:VAL:HG22	2.31	0.59
1:A:131[B]:VAL:HG22	1:B:131[B]:VAL:HG22	1.85	0.58
1:B:20[B]:ARG:HG3	1:B:130:CME:HE3	1.85	0.58
1:A:76[A]:VAL:HG22	1:A:104:TYR:CZ	2.40	0.56
1:B:4:LEU:C	1:B:4:LEU:HD23	2.31	0.56
1:A:12:LYS:NZ	4:A:389:HOH:O	2.23	0.53
1:B:24:ALA:C	1:B:26[A]:ASP:H	2.16	0.53
1:B:9[A]:LEU:HD21	1:B:120:MET:HE2	1.91	0.52
1:A:131[B]:VAL:HG22	1:B:131[B]:VAL:CG2	2.39	0.52
1:A:4[B]:LEU:C	1:A:4[B]:LEU:HD23	2.35	0.52
1:B:32[B]:LEU:HG	1:B:34[B]:LEU:HD11	1.93	0.51
1:B:33:ASN:C	1:B:34[B]:LEU:HD12	2.36	0.51
1:A:8:ASN:HB3	4:B:421:HOH:O	2.11	0.49
1:A:99[C]:LYS:HG2	1:A:105:GLU:HG3	1.95	0.48
1:A:100[A]:LEU:HB3	1:A:101:PRO:CD	2.46	0.46
1:B:20[C]:ARG:HG3	1:B:130:CME:HE3	1.98	0.46
1:A:16:CME:HZ2	1:A:99[C]:LYS:HD2	1.98	0.46
1:B:130:CME:SD	1:B:132:ALA:HB2	2.55	0.45
1:B:74:GLU:HG2	4:B:433:HOH:O	2.17	0.45
1:B:134:ASP:HB2	4:B:422:HOH:O	2.17	0.45
1:A:126:PHE:O	4:A:351:HOH:O	2.20	0.44
1:A:16:CME:HZ2	1:A:99[A]:LYS:HD2	2.01	0.43
1:A:100[A]:LEU:HB3	1:A:101:PRO:HD2	2.00	0.43
1:A:16:CME:HA	1:A:90[B]:THR:HG22	2.01	0.42
1:B:32[B]:LEU:HG	1:B:34[B]:LEU:CD1	2.50	0.42
1:B:32[B]:LEU:HD11	1:B:120:MET:CG	2.49	0.41
1:B:33:ASN:O	1:B:120:MET:HA	2.20	0.41
1:B:4:LEU:HD23	1:B:5:VAL:N	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:395:HOH:O	4:B:398:HOH:O[4_545]	2.16	0.04
1:A:20:ARG:HH12	1:A:26:ASP:OD2[4_555]	1.60	0.00

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/136 (107%)	142 (98%)	3 (2%)	0	100	100
1	B	144/136 (106%)	141 (98%)	3 (2%)	0	100	100
All	All	289/272 (106%)	283 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/107 (114%)	119 (98%)	3 (2%)	42	8
1	B	119/107 (111%)	115 (97%)	4 (3%)	32	4
All	All	241/214 (113%)	234 (97%)	7 (3%)	43	6

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	MET
1	A	127[A]	LYS
1	A	127[B]	LYS
1	B	4	LEU
1	B	26[A]	ASP
1	B	26[B]	ASP
1	B	134	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	95	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	16	1	8,9,10	1.44	1 (12%)	6,9,11	1.33	1 (16%)
1	CME	B	88	1	8,9,10	1.17	0	6,9,11	0.82	0
1	CME	B	16	1	8,9,10	1.16	0	6,9,11	1.03	0
1	CME	A	130	1	8,9,10	1.60	2 (25%)	6,9,11	2.97	2 (33%)
1	CME	B	130	1	8,9,10	1.16	1 (12%)	6,9,11	2.56	3 (50%)
1	CME	A	88	1	8,9,10	1.10	0	6,9,11	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	16	1	-	0/5/8/10	-
1	CME	B	88	1	-	2/5/8/10	-
1	CME	B	16	1	-	0/5/8/10	-
1	CME	A	130	1	-	1/5/8/10	-
1	CME	B	130	1	-	2/5/8/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	88	1	-	1/5/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	16	CME	CA-N	-3.11	1.39	1.48
1	A	130	CME	CE-SD	-2.57	1.71	1.82
1	A	130	CME	CB-CA	2.56	1.59	1.53
1	B	130	CME	CB-SG	-2.04	1.74	1.81

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	CME	CE-SD-SG	6.59	132.38	103.46
1	B	130	CME	CE-SD-SG	5.16	126.10	103.46
1	A	130	CME	CB-SG-SD	-2.83	96.53	103.86
1	B	130	CME	OH-CZ-CE	2.55	120.79	110.82
1	A	16	CME	CZ-CE-SD	2.17	120.64	113.39
1	B	130	CME	CB-SG-SD	2.08	109.23	103.86

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	130	CME	CE-SD-SG-CB
1	A	130	CME	SD-CE-CZ-OH
1	B	130	CME	CZ-CE-SD-SG
1	B	88	CME	CA-CB-SG-SD
1	B	88	CME	SD-CE-CZ-OH
1	A	88	CME	CA-CB-SG-SD

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	16	CME	4	0
1	B	130	CME	5	0

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TVS	C	1	2	18,18,18	1.81	3 (16%)	23,24,24	2.69	5 (21%)
2	TVV	C	2	2	14,14,15	1.87	3 (21%)	18,18,20	1.31	3 (16%)
2	TVS	D	1	2	18,18,18	1.92	3 (16%)	23,24,24	1.73	6 (26%)
2	TVV	D	2	2	14,14,15	1.48	2 (14%)	18,18,20	1.66	4 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TVS	C	1	2	-	2/10/30/30	0/1/1/1
2	TVV	C	2	2	-	3/6/23/26	0/1/1/1
2	TVS	D	1	2	-	1/10/30/30	0/1/1/1
2	TVV	D	2	2	-	2/6/23/26	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	TVS	O3-C3	4.92	1.55	1.43
2	D	2	TVV	O2-C2	4.53	1.52	1.43
2	D	1	TVS	CAA-CAB	4.42	1.57	1.29
2	C	1	TVS	O1-C1	4.39	1.47	1.40
2	C	2	TVV	CAD-CAC	-4.22	1.05	1.18
2	C	1	TVS	CAA-CAB	4.13	1.55	1.29
2	C	2	TVV	CAQ-CAD	-3.61	1.32	1.46
2	D	1	TVS	O1-C1	3.28	1.45	1.40
2	C	1	TVS	C3-C2	2.58	1.57	1.53
2	D	2	TVV	C1-C2	-2.18	1.47	1.52
2	C	2	TVV	O5-C5	2.11	1.47	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	TVS	CAK-O1-C1	9.08	123.09	114.09
2	C	1	TVS	O5-C5-C6	5.63	120.38	106.44
2	C	1	TVS	O5-C1-O1	4.69	121.11	110.04
2	D	2	TVV	CAQ-O3-C3	4.66	119.56	114.56
2	D	1	TVS	CAK-O1-C1	3.88	117.93	114.09
2	D	1	TVS	O3-C3-C2	-3.80	102.03	109.58
2	D	1	TVS	O3-C3-C4	-3.24	102.74	110.38
2	D	2	TVV	O2-C2-C3	-3.21	102.48	109.99
2	C	1	TVS	C2-N2-C7	3.03	130.19	123.11
2	C	2	TVV	O5-C5-C4	-2.58	104.54	110.83
2	D	1	TVS	O5-C1-O1	2.43	115.79	110.04
2	D	1	TVS	C1-O5-C5	-2.27	109.29	113.72
2	D	2	TVV	O2-C2-C1	-2.18	104.23	109.22
2	C	2	TVV	O4-C4-C5	2.15	114.61	109.32
2	C	1	TVS	O6-C6-C5	-2.14	104.04	111.33
2	D	1	TVS	C8-C7-N2	-2.12	112.60	116.12
2	D	2	TVV	C2-C3-C4	-2.10	109.09	110.91
2	C	2	TVV	CAQ-O3-C3	2.07	116.78	114.56

There are no chirality outliers.


All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	TVV	CAC-CAD-CAQ-O3
2	D	1	TVS	CAA-CAB-CAK-O1
2	C	1	TVS	CAA-CAB-CAK-O1
2	C	1	TVS	CAB-CAK-O1-C1
2	D	2	TVV	C4-C3-O3-CAQ
2	D	2	TVV	C2-C3-O3-CAQ
2	C	2	TVV	C4-C3-O3-CAQ
2	C	2	TVV	C2-C3-O3-CAQ

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	202	-	4,4,4	0.28	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	130/136 (95%)	-0.01	2 (1%) 72 73	5, 17, 35, 43	12 (9%)
1	B	130/136 (95%)	-0.06	6 (4%) 37 39	6, 16, 30, 53	11 (8%)
All	All	260/272 (95%)	-0.03	8 (3%) 51 51	5, 16, 33, 53	23 (8%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	PHE	4.8
1	B	2	CYS	4.7
1	B	26[A]	ASP	3.3
1	B	80	GLN	3.2
1	B	93[A]	GLN	3.0
1	B	134	ASP	2.6
1	A	50	ASN	2.1
1	B	20[A]	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	130	10/11	0.95	0.12	10,24,51,61	0
1	CME	A	16	10/11	0.96	0.10	16,23,49,54	0
1	CME	A	88	10/11	0.97	0.10	10,15,37,39	0
1	CME	B	16	10/11	0.97	0.09	11,19,43,50	0
1	CME	B	88	10/11	0.97	0.09	9,19,38,39	0
1	CME	B	130	10/11	0.97	0.14	7,14,55,66	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TVS	C	1	18/18	0.93	0.13	20,38,55,60	10
2	TVS	D	1	18/18	0.96	0.09	17,25,50,55	0
2	TVV	C	2	14/15	0.97	0.07	14,20,34,48	5
2	TVV	D	2	14/15	0.98	0.06	13,18,34,45	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	B	202	5/5	0.98	0.08	19,20,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.