



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:07 PM UTC

PDB ID : 7Q20 / pdb_00007q20
Title : Ruminococcus gnavus ATC29149 endo-beta-1,4-galactosidase (RgGH98) in complex with blood group A trisaccharide
Authors : Owen, C.D.; Wu, H.; Crost, E.H.; van Bakel, W.; Gascuena, A.M.; Latousakis, D.; Hicks, T.; Walpole, S.; Urbanowicz, P.A.; Ndeh, D.; Monaco, S.; Salom, L.S.; Griffiths, R.; Colvile, A.; Spencer, D.I.R.; Walsh, M.A.; Angulo, J.; Juge, N.
Deposited on : 2021-10-22
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	846	3% 95% .
1	G	846	% 95% 5%
2	E	3	67% 33%
2	H	3	100%

Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 14264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ruminococcus gnavus endogalactosidase GH98.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	845	Total	C	N	O	S	0	1	0
			6679	4220	1101	1333	25			
1	G	845	Total	C	N	O	S	0	0	0
			6672	4216	1100	1331	25			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	3	Total	C	N	O	0	0	0
			36	20	1	15			
2	E	3	Total	C	N	O	0	0	0
			36	20	1	15			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		
3	G	1	Total	Ca	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	3	Total	Mg	0	0
			3	3		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	C	O	0	0
			10	6	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	368	Total	O	0	0
			368	368		
6	G	455	Total	O	0	0
			455	455		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: *Ruminococcus gnavus* endogalactosidase GH98




- Molecule 1: *Ruminococcus gnavus* endogalactosidase GH98



- Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose



- Molecule 2: alpha-L-fucopyranose-(1-2)-[2-acetamido-2-deoxy-alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	107.59Å 86.95Å 110.20Å 90.00° 100.33° 90.00°	Depositor
Resolution (Å)	108.41 – 1.95 108.41 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (108.41-1.95) 98.8 (108.41-1.95)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.185 , 0.219 0.192 , 0.225	Depositor DCC
R_{free} test set	7403 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14264	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, A2G, FUC, PGE, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	2/6841 (0.0%)	0.89	0/9292
1	G	0.71	0/6833	0.88	0/9279
All	All	0.69	2/13674 (0.0%)	0.88	0/18571

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	889	ILE	CA-C	6.07	1.58	1.53
1	A	842	ASP	CB-CG	5.17	1.65	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6679	0	6351	23	0
1	G	6672	0	6350	31	0
2	E	36	0	32	0	0
2	H	36	0	32	0	0
3	A	3	0	0	0	0
3	G	1	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	3	0	0	0	0
5	G	10	0	14	0	0
6	A	368	0	0	1	0
6	G	455	0	0	0	0
All	All	14264	0	12779	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:824:LYS:HG3	1:A:859:GLY:HA3	1.54	0.88
1:G:373:ALA:H	1:G:412:ASN:HD22	1.22	0.87
1:G:775:THR:OG1	1:G:777:ASP:OD1	2.05	0.74
1:G:475:ASN:HB3	1:G:623:GLN:HE21	1.53	0.74
1:G:736:HIS:HD2	1:G:758:ARG:HH11	1.36	0.73
1:G:708:VAL:O	1:G:736:HIS:HE1	1.81	0.63
1:A:445:HIS:HD2	1:A:482:ASP:OD2	1.84	0.61
1:A:824:LYS:CG	1:A:859:GLY:HA3	2.30	0.58
1:G:82:LYS:HZ2	1:G:134:ASN:HD21	1.53	0.57
1:G:370:LEU:HD21	1:G:420:PRO:HA	1.87	0.57
1:G:308:ASN:ND2	1:G:567:ASN:H	2.01	0.57
1:G:82:LYS:NZ	1:G:134:ASN:HD21	2.04	0.55
1:A:729:LEU:HD23	1:A:741:MET:HE2	1.92	0.52
1:A:512:ASP:HA	1:A:530:ARG:HB3	1.92	0.52
1:A:824:LYS:HG3	1:A:859:GLY:CA	2.33	0.50
1:A:319:ASP:OD1	1:A:322:LYS:HE2	2.11	0.50
1:G:512:ASP:HA	1:G:530:ARG:HB3	1.94	0.50
1:G:754:LEU:HD12	1:G:754:LEU:C	2.37	0.50
1:G:308:ASN:HD22	1:G:567:ASN:H	1.61	0.49
1:G:736:HIS:CD2	1:G:758:ARG:HH11	2.23	0.49
1:G:285:ILE:HB	1:G:555:THR:HB	1.95	0.48
1:G:373:ALA:H	1:G:412:ASN:ND2	2.01	0.48
1:G:409:ASN:OD1	1:G:409:ASN:C	2.57	0.48
1:A:754:LEU:C	1:A:754:LEU:HD12	2.38	0.47
1:A:409:ASN:OD1	1:A:409:ASN:C	2.57	0.46
1:A:314:TRP:CH2	1:A:364:PRO:HD2	2.50	0.46
1:A:729:LEU:CD2	1:A:741:MET:HE2	2.45	0.46
1:A:761:LYS:NZ	6:A:1011:HOH:O	2.48	0.46
1:A:285:ILE:HB	1:A:555:THR:HB	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:889:ILE:HD12	1:A:889:ILE:N	2.30	0.46
1:A:488:SER:HB3	1:A:635:VAL:HG23	1.98	0.45
1:G:127:SER:OG	1:G:174:ASN:ND2	2.50	0.45
1:G:672:SER:O	1:G:673:GLY:C	2.60	0.45
1:G:708:VAL:O	1:G:736:HIS:CE1	2.68	0.45
1:G:729:LEU:HD23	1:G:741:MET:HE2	1.97	0.45
1:A:483:ASP:HB2	1:A:623:GLN:CD	2.42	0.45
1:A:677:GLN:HE21	1:A:677:GLN:HB3	1.65	0.45
1:A:505:LEU:C	1:A:505:LEU:HD23	2.42	0.44
1:G:314:TRP:CH2	1:G:364:PRO:HD2	2.52	0.44
1:G:505:LEU:C	1:G:505:LEU:HD23	2.43	0.44
1:G:729:LEU:CD2	1:G:741:MET:HE2	2.48	0.44
1:G:710:THR:HG22	1:G:712:GLU:HG3	1.99	0.43
1:G:373:ALA:N	1:G:412:ASN:HD22	2.02	0.43
1:A:543:MET:HG3	1:A:556:PHE:HZ	1.82	0.43
1:A:754:LEU:O	1:A:870:TRP:HA	2.19	0.43
1:G:211:GLY:CA	1:G:344:GLN:HE22	2.32	0.43
1:G:473:TYR:CZ	1:G:504:MET:HB2	2.54	0.42
1:G:460:LYS:HE3	1:G:460:LYS:HA	2.01	0.42
1:G:308:ASN:ND2	1:G:313:ARG:HH11	2.17	0.42
1:A:473:TYR:CZ	1:A:504:MET:HB2	2.55	0.42
1:G:483:ASP:HB2	1:G:623:GLN:CD	2.45	0.41
1:G:754:LEU:O	1:G:870:TRP:HA	2.20	0.41
1:A:855:ASN:O	1:A:858:THR:O	2.39	0.40
1:A:543:MET:CG	1:A:556:PHE:HZ	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	844/846 (100%)	826 (98%)	18 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	843/846 (100%)	825 (98%)	18 (2%)	0	100	100
All	All	1687/1692 (100%)	1651 (98%)	36 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	713/713 (100%)	705 (99%)	8 (1%)	65	64
1	G	712/713 (100%)	706 (99%)	6 (1%)	73	74
All	All	1425/1426 (100%)	1411 (99%)	14 (1%)	70	67

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	GLU
1	A	143[A]	GLN
1	A	143[B]	GLN
1	A	340	ASP
1	A	370	LEU
1	A	409	ASN
1	A	453	VAL
1	A	669	LYS
1	G	94	GLU
1	G	340	ASP
1	G	370	LEU
1	G	409	ASN
1	G	460	LYS
1	G	835	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	86	GLN
1	A	87	HIS
1	A	198	GLN
1	A	268	GLN
1	A	315	ASN
1	A	378	ASN
1	A	445	HIS
1	A	491	GLN
1	A	575	HIS
1	A	677	GLN
1	A	802	ASN
1	G	68	ASN
1	G	100	ASN
1	G	134	ASN
1	G	174	ASN
1	G	305	GLN
1	G	308	ASN
1	G	315	ASN
1	G	333	ASN
1	G	344	GLN
1	G	378	ASN
1	G	412	ASN
1	G	480	GLN
1	G	520	GLN
1	G	586	ASN
1	G	623	GLN
1	G	677	GLN
1	G	736	HIS
1	G	742	ASN
1	G	786	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GAL	E	1	2	12,12,12	0.56	0	17,17,17	0.74	0
2	FUC	E	2	2	10,10,11	0.52	0	14,14,16	0.56	0
2	A2G	E	3	2	14,14,15	0.38	0	17,19,21	0.68	1 (5%)
2	GAL	H	1	2	12,12,12	0.47	0	17,17,17	0.60	0
2	FUC	H	2	2	10,10,11	0.39	0	14,14,16	0.47	0
2	A2G	H	3	2	14,14,15	0.43	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	E	1	2	-	0/2/22/22	0/1/1/1
2	FUC	E	2	2	-	-	0/1/1/1
2	A2G	E	3	2	-	0/6/23/26	0/1/1/1
2	GAL	H	1	2	-	0/2/22/22	0/1/1/1
2	FUC	H	2	2	-	-	0/1/1/1
2	A2G	H	3	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	3	A2G	O5-C1-C2	-2.28	107.76	111.29


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PGE	G	901	-	9,9,9	0.21	0	8,8,8	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PGE	G	901	-	-	5/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	901	PGE	O2-C3-C4-O3
5	G	901	PGE	O1-C1-C2-O2
5	G	901	PGE	C3-C4-O3-C5
5	G	901	PGE	C4-C3-O2-C2
5	G	901	PGE	O3-C5-C6-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	845/846 (99%)	0.30	24 (2%) 55 62	25, 43, 74, 105	1 (0%)
1	G	845/846 (99%)	0.08	7 (0%) 82 86	22, 37, 63, 85	0
All	All	1690/1692 (99%)	0.19	31 (1%) 67 75	22, 40, 70, 105	1 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	826	ALA	4.1
1	A	673	GLY	3.7
1	A	854	PHE	3.0
1	A	833	VAL	2.9
1	A	781	GLY	2.7
1	A	858	THR	2.7
1	A	859	GLY	2.7
1	G	781	GLY	2.6
1	A	49	ALA	2.6
1	A	875	ILE	2.5
1	G	675	ALA	2.4
1	A	893	PRO	2.4
1	A	831	VAL	2.4
1	A	195	GLY	2.4
1	A	881	VAL	2.3
1	A	876	THR	2.3
1	A	672	SER	2.2
1	G	682	ASP	2.2
1	A	857	GLU	2.2
1	A	822	LEU	2.1
1	A	886	ALA	2.1
1	A	745	GLY	2.1
1	A	880	PHE	2.1
1	G	881	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	861	ALA	2.1
1	A	744	SER	2.0
1	G	882	TYR	2.0
1	G	825	LEU	2.0
1	A	827	LYS	2.0
1	G	195	GLY	2.0
1	A	852	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GAL	H	1	12/12	-	-	40,43,51,53	0
2	FUC	H	2	10/11	-	-	35,35,36,38	0
2	A2G	H	3	14/15	-	-	32,41,45,47	0
2	GAL	E	1	12/12	-	-	33,38,41,41	0
2	FUC	E	2	10/11	-	-	30,33,36,37	0
2	A2G	E	3	14/15	-	-	28,34,39,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PGE	G	901	10/10	0.68	0.25	70,76,79,81	0
4	MG	G	905	1/1	0.90	0.16	38,38,38,38	0
4	MG	A	904	1/1	0.90	0.07	54,54,54,54	0
3	CA	A	903	1/1	0.91	0.17	58,58,58,58	0
3	CA	A	901	1/1	0.92	0.08	56,56,56,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	902	1/1	0.96	0.06	32,32,32,32	0
4	MG	G	904	1/1	0.96	0.11	30,30,30,30	0
4	MG	G	903	1/1	0.97	0.04	31,31,31,31	0
3	CA	G	902	1/1	0.97	0.04	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.