



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:12 PM UTC

PDB ID : 8Q24 / pdb_00008q24
Title : HsNMT1 in complex with both MyrCoA and (DAB)SFSKPR inhibitor peptide
Authors : Dian, C.; Giglione, C.; Meinnel, T.
Deposited on : 2023-08-01
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	5% 87% 9%
1	B	402	3% 89% 8%
2	C	7	57% 86% 14%
2	E	7	71% 14% 14%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	3	0
			3145	2038	529	562	16			
1	B	392	Total	C	N	O	S	0	6	0
			3223	2087	540	579	17			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	GLY	-	expression tag	UNP P30419
A	96	GLY	-	expression tag	UNP P30419
A	97	SER	-	expression tag	UNP P30419
A	98	GLU	-	expression tag	UNP P30419
B	95	GLY	-	expression tag	UNP P30419
B	96	GLY	-	expression tag	UNP P30419
B	97	SER	-	expression tag	UNP P30419
B	98	GLU	-	expression tag	UNP P30419

- Molecule 2 is a protein called DAB-SER-PHE-SER-LYS-PRO-ARG.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	7	Total	C	N	O	0	0	0
			58	36	12	10			
2	C	7	Total	C	N	O	0	0	0
			58	36	12	10			

- Molecule 3 is TETRADECANOYL-COA (CCD ID: MYA) (formula: C₃₅H₆₂N₇O₁₇P₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		
3	B	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	238	Total	O	0	0
			238	238		
6	B	246	Total	O	0	0
			246	246		
6	E	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1




- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1



- Molecule 2: DAB-SER-PHE-SER-LYS-PRO-ARG



- Molecule 2: DAB-SER-PHE-SER-LYS-PRO-ARG



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.74Å 177.99Å 58.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.75 – 1.90 48.75 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.75-1.90) 99.9 (48.75-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.17_3644	Depositor
R, R_{free}	0.172 , 0.215 0.173 , 0.216	Depositor DCC
R_{free} test set	3244 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7129	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MYA, CL, DAB, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/3230	0.48	0/4389
1	B	0.28	0/3314	0.47	0/4507
2	C	0.23	0/52	0.49	0/66
2	E	0.25	0/52	0.63	0/66
All	All	0.28	0/6648	0.48	0/9028

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3145	0	3110	22	0
1	B	3223	0	3176	21	1
2	C	58	0	61	2	0
2	E	58	0	61	2	0
3	A	63	0	58	1	0
3	B	63	0	58	1	0
4	A	12	0	16	1	0
4	B	18	0	24	3	0
5	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	238	0	0	2	0
6	B	246	0	0	1	0
6	E	3	0	0	0	0
All	All	7129	0	6564	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:414:LYS:H	4:B:504:GOL:H32	1.62	0.64
1:A:173:TYR:CE2	1:A:194:PRO:HG3	2.34	0.63
1:A:361:HIS:CD2	1:A:490:LYS:HE2	2.35	0.62
1:B:410:HIS:HD2	1:B:412:THR:H	1.46	0.62
2:E:2:DAB:HD1	2:E:2:DAB:H2	1.48	0.61
1:B:410:HIS:CD2	1:B:413:HIS:H	2.20	0.59
1:A:179:ASN:O	1:A:255:ARG:NH2	2.37	0.58
1:A:170:LYS:HA	1:A:173:TYR:CE2	2.39	0.57
1:A:108:THR:HG23	1:A:111:GLU:H	1.69	0.57
1:B:304:ARG:O	1:B:308:GLU:HG3	2.04	0.57
1:B:258:ARG:NH2	3:B:501:MYA:O9A	2.37	0.57
1:B:108:THR:HG23	1:B:111:GLU:H	1.71	0.56
1:B:334:LYS:NZ	6:B:603:HOH:O	2.40	0.55
1:A:282:THR:HG23	1:A:495:LEU:HD12	1.90	0.54
1:B:302:ASN:OD1	1:B:443:LYS:NZ	2.27	0.53
1:A:195:GLU:HB3	1:A:381:ILE:HD11	1.90	0.53
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.44	0.53
1:A:307:ILE:HD13	1:A:314:LEU:HD13	1.92	0.52
1:A:185:ASP:HB2	2:C:6:LYS:HE3	1.92	0.52
1:A:496:GLN:HE22	4:A:503:GOL:H31	1.75	0.51
1:B:259:VAL:O	1:B:262[B]:VAL:HG22	2.11	0.51
1:B:303:PRO:O	1:B:307:ILE:HG12	2.11	0.51
1:B:458:ASN:HA	1:B:461:PHE:CE2	2.46	0.51
1:B:301:LEU:O	1:B:443:LYS:NZ	2.40	0.49
1:A:130:GLU:O	1:A:289:LYS:HE3	2.14	0.48
1:A:183:ASP:HB2	2:C:6:LYS:NZ	2.29	0.47
1:A:395:THR:HB	1:A:428[B]:GLN:HB2	1.96	0.47
1:A:358:LYS:NZ	6:A:612:HOH:O	2.48	0.47
1:B:414:LYS:N	4:B:504:GOL:H32	2.29	0.46
1:A:318:MET:HE1	1:A:326:LEU:HD22	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:295:ARG:NH1	1:B:473:ASN:HD21	2.13	0.46
1:B:405:SER:OG	2:E:4:PHE:HE2	1.99	0.46
1:A:128:LEU:HD23	1:A:128:LEU:HA	1.86	0.45
1:A:238:THR:HG22	1:A:240:LYS:HG2	2.00	0.44
1:B:251:HIS:ND1	1:B:253:LYS:HG2	2.33	0.44
1:A:177:ASN:O	1:A:189:ARG:HD2	2.18	0.43
1:A:248:LEU:HG	3:A:501:MYA:H14	2.01	0.43
1:B:366:MET:HE2	4:B:503:GOL:H12	2.02	0.42
1:A:285:VAL:HG21	6:A:625:HOH:O	2.21	0.41
1:A:321:GLN:H	1:A:321:GLN:NE2	2.19	0.41
1:B:282:THR:HB	1:B:474:LEU:HD11	2.03	0.40
1:B:410:HIS:CD2	1:B:412:THR:H	2.31	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:LYS:NZ	1:B:331:GLU:OE1[1_556]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/402 (96%)	378 (98%)	7 (2%)	0	100	100
1	B	397/402 (99%)	386 (97%)	11 (3%)	0	100	100
2	C	5/7 (71%)	5 (100%)	0	0	100	100
2	E	5/7 (71%)	5 (100%)	0	0	100	100
All	All	792/818 (97%)	774 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/362 (94%)	337 (98%)	5 (2%)	57	56
1	B	353/362 (98%)	349 (99%)	4 (1%)	65	67
2	C	6/6 (100%)	6 (100%)	0	100	100
2	E	6/6 (100%)	6 (100%)	0	100	100
All	All	707/736 (96%)	698 (99%)	9 (1%)	61	61

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	LYS
1	A	254	LEU
1	A	285	VAL
1	A	318	MET
1	A	455	LEU
1	B	134	THR
1	B	239	GLU
1	B	304	ARG
1	B	455	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	GLN
1	A	154	GLN
1	A	234	HIS
1	A	321	GLN
1	A	458	ASN
1	B	118	GLN
1	B	380	ASN
1	B	410	HIS
1	B	458	ASN
1	B	473	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DAB	C	2	2	5,6,7	0.93	0	1,6,8	0.38	0
2	DAB	E	2	2	5,6,7	0.94	1 (20%)	1,6,8	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DAB	C	2	2	-	0/4/5/7	-
2	DAB	E	2	2	-	0/4/5/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	DAB	O-C	2.01	1.27	1.20

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2	DAB	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	502	-	5,5,5	1.07	0	5,5,5	0.89	0
4	GOL	B	504	-	5,5,5	1.06	0	5,5,5	0.99	0
4	GOL	A	503	-	5,5,5	0.76	0	5,5,5	1.23	0
3	MYA	A	501	-	63,65,65	1.45	10 (15%)	85,91,91	1.66	14 (16%)
4	GOL	B	502	-	5,5,5	1.14	0	5,5,5	1.03	0
3	MYA	B	501	-	63,65,65	1.66	10 (15%)	85,91,91	1.94	17 (20%)
4	GOL	B	503	-	5,5,5	1.10	0	5,5,5	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	502	-	-	0/4/4/4	-
4	GOL	B	504	-	-	4/4/4/4	-
4	GOL	A	503	-	-	4/4/4/4	-
3	MYA	A	501	-	-	6/64/80/80	0/3/3/3
4	GOL	B	502	-	-	0/4/4/4	-
3	MYA	B	501	-	-	6/64/80/80	0/3/3/3
4	GOL	B	503	-	-	4/4/4/4	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	MYA	P1A-O3A	7.40	1.67	1.59
3	B	501	MYA	P2A-O3A	4.26	1.64	1.59
3	A	501	MYA	C2A-N3A	3.81	1.40	1.33
3	B	501	MYA	C2A-N3A	3.59	1.40	1.33
3	A	501	MYA	P1A-O3A	3.53	1.63	1.59
3	A	501	MYA	C3M-C2M	3.51	1.54	1.50
3	A	501	MYA	P2A-O3A	3.44	1.63	1.59
3	B	501	MYA	C2A-N1A	3.32	1.39	1.33
3	A	501	MYA	C2A-N1A	3.28	1.39	1.33
3	B	501	MYA	C3M-C2M	2.81	1.53	1.50
3	A	501	MYA	C5A-N7A	-2.65	1.34	1.39
3	B	501	MYA	C5A-N7A	-2.55	1.34	1.39
3	B	501	MYA	C1X-N9A	2.45	1.53	1.46
3	A	501	MYA	P3X-O3X	2.44	1.63	1.59
3	A	501	MYA	C5A-C4A	-2.32	1.35	1.39
3	A	501	MYA	C5A-C6A	-2.30	1.34	1.41
3	A	501	MYA	C1X-N9A	2.23	1.52	1.46
3	B	501	MYA	C5A-C6A	-2.13	1.35	1.41
3	B	501	MYA	C5A-C4A	-2.12	1.35	1.39
3	B	501	MYA	P3X-O8A	-2.01	1.47	1.54

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	MYA	O3A-P2A-O4A	-8.44	85.31	110.70
3	A	501	MYA	N3A-C2A-N1A	-6.98	118.02	128.58
3	B	501	MYA	N3A-C2A-N1A	-6.86	118.19	128.58
3	B	501	MYA	O5A-P2A-O3A	-5.45	92.53	107.27
3	A	501	MYA	C2X-C3X-C4X	-4.42	95.50	103.24
3	B	501	MYA	C5A-C4A-N3A	-4.36	120.71	126.72
3	A	501	MYA	C5A-C4A-N3A	-4.32	120.76	126.72
3	A	501	MYA	N9A-C8A-N7A	-3.86	108.45	113.94
3	B	501	MYA	N9A-C8A-N7A	-3.71	108.68	113.94
3	A	501	MYA	C2A-N3A-C4A	3.29	119.87	111.83
3	B	501	MYA	C2A-N3A-C4A	3.28	119.85	111.83
3	B	501	MYA	C5A-C4A-N9A	3.16	109.25	105.81
3	B	501	MYA	C4A-C5A-N7A	-3.13	107.00	110.58
3	A	501	MYA	O4X-C1X-N9A	3.08	114.01	108.09
3	A	501	MYA	C5A-N7A-C8A	2.99	108.15	103.45
3	B	501	MYA	C5A-N7A-C8A	2.93	108.06	103.45
3	B	501	MYA	O5A-P2A-O4A	2.89	125.87	112.44
3	A	501	MYA	C4A-C5A-N7A	-2.83	107.34	110.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	MYA	C2X-C3X-C4X	-2.80	98.34	103.24
3	A	501	MYA	C5A-C4A-N9A	2.64	108.69	105.81
3	B	501	MYA	P3X-O3X-C3X	-2.64	116.38	123.43
3	B	501	MYA	O2M-C2M-C3M	-2.40	121.21	123.98
3	A	501	MYA	O2M-C2M-C3M	-2.37	121.25	123.98
3	A	501	MYA	C5A-C6A-N6A	-2.35	117.46	123.29
3	A	501	MYA	C6A-C5A-C4A	2.34	120.37	117.18
3	B	501	MYA	O6A-P2A-O4A	2.29	118.00	108.94
3	B	501	MYA	O4X-C1X-C2X	-2.24	101.82	106.62
3	B	501	MYA	C6A-C5A-C4A	2.24	120.23	117.18
3	A	501	MYA	C4M-C3M-C2M	-2.20	107.41	112.27
3	A	501	MYA	C3M-C2M-S1	2.11	115.92	113.40
3	B	501	MYA	C5A-C6A-N6A	-2.10	118.10	123.29

There are no chirality outliers.

All (24) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	501	MYA	C12-O6A-P2A-O4A
4	A	503	GOL	O1-C1-C2-C3
4	A	503	GOL	O2-C2-C3-O3
4	B	503	GOL	O1-C1-C2-C3
4	B	504	GOL	O2-C2-C3-O3
4	A	503	GOL	C1-C2-C3-O3
4	B	503	GOL	C1-C2-C3-O3
4	B	504	GOL	O1-C1-C2-C3
4	B	504	GOL	C1-C2-C3-O3
4	A	503	GOL	O1-C1-C2-O2
4	B	503	GOL	O1-C1-C2-O2
4	B	503	GOL	O2-C2-C3-O3
4	B	504	GOL	O1-C1-C2-O2
3	A	501	MYA	C3X-C4X-C5X-O5X
3	B	501	MYA	P1A-O3A-P2A-O4A
3	A	501	MYA	O4X-C4X-C5X-O5X
3	A	501	MYA	C2X-C1X-N9A-C8A
3	A	501	MYA	C5X-O5X-P1A-O2A
3	B	501	MYA	P1A-O3A-P2A-O5A
3	A	501	MYA	O4X-C1X-N9A-C8A
3	B	501	MYA	C2X-C1X-N9A-C8A
3	B	501	MYA	C11-C12-O6A-P2A
3	A	501	MYA	C2X-C1X-N9A-C4A
3	B	501	MYA	C8M-C9M-CAM-CBM


There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	504	GOL	2	0
4	A	503	GOL	1	0
3	A	501	MYA	1	0
3	B	501	MYA	1	0
4	B	503	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/402 (96%)	0.26	21 (5%) 31 33	14, 33, 69, 84	3 (0%)
1	B	392/402 (97%)	0.20	14 (3%) 46 49	11, 34, 65, 94	6 (1%)
2	C	6/7 (85%)	2.55	4 (66%) 0 0	68, 71, 75, 76	0
2	E	6/7 (85%)	0.98	0 100 100	46, 49, 52, 73	0
All	All	790/818 (96%)	0.25	39 (4%) 35 37	11, 34, 69, 94	9 (1%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	314	LEU	4.1
1	A	106	ALA	3.6
1	A	472	GLY	3.6
1	B	313[A]	HIS	3.6
1	B	393[A]	GLU	3.5
2	C	7	PRO	3.4
1	B	106	ALA	3.2
2	C	3	SER	3.1
1	A	133	ASN	3.1
2	C	4	PHE	3.0
1	A	181	VAL	3.0
1	B	131	VAL	3.0
1	B	315	SER	3.0
1	B	113	SER	2.9
1	B	114	LYS	2.9
1	A	131	VAL	2.9
1	A	317	ASN	2.8
1	B	105	PRO	2.8
1	A	314	LEU	2.7
1	A	108	THR	2.7
1	B	323	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	ASP	2.6
1	A	116	SER	2.6
1	A	192	TYR	2.5
1	A	254	LEU	2.5
1	A	408	MET	2.5
1	A	132	VAL	2.4
1	A	107	LYS	2.4
1	A	187	MET	2.2
1	B	184	ASP	2.2
1	A	414	LYS	2.2
1	A	182	GLU	2.2
1	B	132	VAL	2.1
1	B	318	MET	2.1
2	C	5	SER	2.1
1	A	296	TYR	2.0
1	A	496	GLN	2.0
1	B	128	LEU	2.0
1	A	185	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DAB	C	2	7/8	0.58	0.19	52,57,65,70	0
2	DAB	E	2	7/8	0.84	0.13	34,40,43,45	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	504	6/6	0.64	0.17	50,55,61,65	0
3	MYA	A	501	63/63	0.81	0.14	21,51,64,76	0
4	GOL	A	503	6/6	0.82	0.12	41,45,47,47	0
4	GOL	B	503	6/6	0.85	0.11	33,42,46,46	0
4	GOL	A	502	6/6	0.95	0.06	26,30,32,36	0
3	MYA	B	501	63/63	0.95	0.07	21,34,43,47	0
5	CL	A	505	1/1	0.95	0.08	49,49,49,49	0
4	GOL	B	502	6/6	0.96	0.07	31,35,39,44	0
5	CL	A	504	1/1	0.98	0.10	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.